

Sevilla Carpets of Deterministic Non-cooperative P Systems

ARTIOM ALHAZOV^{1,2}, SERGEY VERLAN³

¹Institute of Mathematics and Computer Science
Academy of Sciences of Moldova
E-mail: artiom@math.md

²Research Group on Mathematical Linguistics
Rovira i Virgili University, Tarragona, Spain
E-mail: artiome.alhazov@estudiants.urv.es

³Laboratoire d'Informatique Theorique et Appliquee
Universite de Metz, France
E-mail: verlan@lita.univ-metz.fr

Abstract

A Sevilla carpet is a graphical representation of the sequence of multisets of rules applied at every step of the evolution of a P system. This paper is devoted to the research of the shape of Sevilla carpets. The case of deterministic non-cooperative systems is considered, and a characterization of Sevilla carpets is given.

1 Introduction

P systems were introduced by Gh. Păun in [2] as distributed parallel computing devices of biochemical inspiration. These systems are inspired from the structure and the functioning of a living cell. The cell is considered as a set of compartments (membranes) nested one in another and which contain objects and evolution rules. The base model specifies neither the nature of these objects, nor the nature of rules. Numerous variants specify these two parameters by obtaining a lot of different models of computing, see [5] for a comprehensive bibliography.

Various approaches are used to study these systems. Like in the classical theory of formal languages, first, the computational power with respect to different parameters is studied. There are many such decidability and undecidability results of for most of the classes of P systems. After that, complexity classes are introduced and investigated.

Recently, efforts to make models of different phenomena based on P systems were made. In this case, the evolution dynamics of the system is studied and no more the generated language. Sevilla carpets permit to have a first overview of the dynamic evolution of the system [1]. More exactly, a Sevilla carpet is a time/frequency diagram which shows how many times a rule is applied at some moment of time. Potentially, Sevilla carpets may be used to locate critical sections of the execution and to facilitate an optimisation of the system.

In this article, we consider a particular class of P systems, *deterministic non-cooperative P systems*, and we study Sevilla carpets associated to this model. In this case, we show that we may reduce the problem to growth functions of DOL systems. We give several

examples that illustrate this reduction and we present at the end some remarks and open questions.

2 Preliminaries

We will use the following notations. For a string $w \in O^*$ and a symbol $a \in O$ we will denote the number of occurrences of a in w by $|w|_a$ and the length of w by $|w|$; for a string w and a number i , $1 \leq i \leq |w|$, we will denote the i -th symbol of w by $w(i)$. We will write the components of vector $v \in \mathbb{N}^k$ as $v[i]$, $1 \leq i \leq k$.

3 Reducing the Problem to L Systems

Consider a deterministic non-cooperative P system. Without restricting the generality, we can assume that all rules act either on different objects, or in different regions; hence, every object-region pair is associated to at most one rule. Notice that the objects produced in the regions where they do not evolve do not influence the carpet, so assuming no idle objects does not restrict the generality.

Take such a P system $\Pi = (O, \mu, w_1, \dots, w_m, R_1, \dots, R_m)$, assume that the rules R_1, \dots, R_m are uniquely labeled, and the set of labels is V . Let us define a mapping $\rho: \{1, \dots, m\} \times O \rightarrow V \cup \{\lambda\}$ as

$$\rho(i, a) = \begin{cases} \lambda & \text{if } R_i \text{ has no rule } a \rightarrow \lambda \\ l & \text{if } l \text{ is a label of some rule } (a \rightarrow \lambda) \in R_i \end{cases}$$

(recall that there exists at most one).

Construct a deterministic L system $G = (V, w_r, P)$, where

$$w_r = \prod_{i=1}^m \prod_{j=1}^{|w_i|} \rho(i, w_i(j)),$$

$$P = \{l \rightarrow h(l) \mid l = rho(i, a) \neq \lambda, h(l) = \prod_{j=1}^{|\alpha|} \rho(i, \alpha(j)),$$

$$(a \rightarrow \alpha) \in R_i, a \in O, 1 \leq i \leq m\}.$$

Consider the evolutions of Π and G ; notice that each (copy of an) object in G corresponds to (application of a) rule in Π . In this way, in the deterministic non-cooperative case the characterization of Sevilla carpets can be reduced to the study of the (Parikh vector) growth functions of $D0L$ systems. The converse is quite simple: given a $D0L$ system, a P system with the same behavior (carpet) can be obtained just by putting the $D0L$ in one membrane.

For the rest of the paper, let us introduce another useful representation. Suppose that the elements of V are ordered ($|V| = k$, $V = \{l_1, \dots, l_k\}$). Let $v = (|w_r|_{l_1}, \dots, |w_r|_{l_k})$ be the vector of initial multiplicities. Let $M = (n_{i,j})_{i,j \in \{1, \dots, k\}}$ be the transition matrix. The elements of M are defined as $n_{i,j} = |h(l_i)|_{l_j}$.

Notice that $Ps(w_r) = v$, $Ps(h(w_r)) = vM$, etc., so $Ps(h^s(w_r)) = vM^s$, $s \geq 0$. Recall that the elements of Sevilla carpet of the original system (number of applications of rule l_i at time $s \geq 0$) correspond to the components of vM^s (i.e., to numbers $(vM^s)[i]$, $1 \leq i \leq k$, $s \geq 0$).

4 Vector Growth Functions

This approach can be found in some university algebra textbook, or in [4]. For a transition matrix M , the characteristic equation $\det(M - pE)$ needs to be solved to obtain the eigenvalues p_1, \dots, p_e with multiplicities m_1, \dots, m_e , $m_1 + \dots + m_e = k$. The general solution of the growth function is of the form $vM^s = (v_{1,1} + \dots + v_{1,m_1} s^{m_1-1})p_1^s + \dots + (v_{e,1} + \dots + v_{e,m_e} s^{m_e-1})p_e^s$. The coefficients $v_{1,1}, \dots, v_{e,m_e}$ can be determined from the equations obtained from the first k steps ($0 \leq s \leq k-1$).

5 Example of Two Rules

Consider a P system with two objects: $\Pi = (\{a, b\}, [1]_1, a^{v[1]}b^{v[2]}, R)$, $R = \{p_1 : a \rightarrow a^{n_{1,1}}b^{n_{1,2}}, p_2 : b \rightarrow a^{n_{2,1}}b^{n_{2,2}}\}$. It corresponds to a DOL system where P consists of rules of R , where a is replaced by p_1 and b is replaced by p_2 . (Notice that $\Pi' = (\{a\}, [1]_2 [2]_2]_1, a^{v[1]}, a^{v[2]}, R)$ with $R_1 = \{p_1 : a \rightarrow a^{n_{1,1}}a_{in}^{n_{1,2}}\}$, $R_2 = \{p_2 : a \rightarrow a_{out}^{n_{2,1}}a^{n_{2,2}}\}$ would correspond to a DOL system where P consists of rules of R , where a in region 1 is replaced by p_1 and a in region 2 is replaced by p_2 . The DOL system corresponding to Π and Π' is the same: $G = (\{p_1, p_2\}, w, P)$, $P = \{p_1 \rightarrow p_1^{n_{1,1}}p_2^{n_{1,2}}, p_2p_1^{n_{2,1}}p_2^{n_{2,2}}\}$.) The transition matrix is $(n_{i,j})_{i,j \in \{1,2\}}$.

The characteristic equation is $p^2 - (n_{1,1} + n_{2,2})p + (n_{1,1}n_{2,2} - n_{1,2}n_{2,1}) = 0$. There are two cases, depending on whether $D = (n_{1,1} + n_{2,2})^2 - 4(n_{1,1}n_{2,2} - n_{1,2}n_{2,1})$ is zero or not. Notice that $D = (n_{1,1} + n_{2,2})^2 - 4(n_{1,1}n_{2,2} - n_{1,2}n_{2,1}) = (n_{1,1} - n_{2,2})^2 + 4(n_{1,2}n_{2,1})$ cannot be negative with nonnegative values of coefficients $n_{i,j}$, so we will always obtain the real roots of the characteristic equation.

Case 1: $p_1 = p_2 = p = (n_{1,1} + n_{2,2})/2$, $vM^s = (v_{1,1} + v_{1,2}s)p^s$. Finding the coefficients: $v_{1,1} = v$, $p(v_{1,1} + v_{1,2}) = vM$, thus $v_{1,1} = v$, $v_{1,2} = v(M - pE)/p$ ($p_1 = p_2 = 0$ is a special case, but in this case everything is erased in one step).

Putting it all together, if $n_{1,1} = n_{2,2}$ and $n_{1,2}n_{2,1} = 0$, then

$$vM^s = v \left(1 + \frac{2M - (n_{1,1} + n_{2,2})E}{n_{1,1} + n_{2,2}} s \right) \left(\frac{n_{1,1} + n_{2,2}}{2} \right)^s$$

Componentwise:

$$\begin{aligned} (vM^s)[1] &= \left(v[1] + \frac{2v[1]n_{1,1} + 2v[2]n_{1,2} - (n_{1,1} + n_{2,2})}{n_{1,1} + n_{2,2}} s \right) \left(\frac{n_{1,1} + n_{2,2}}{2} \right)^s \\ (vM^s)[2] &= \left(v[2] + \frac{2v[1]n_{2,1} + 2v[2]n_{2,2} - (n_{1,1} + n_{2,2})}{n_{1,1} + n_{2,2}} s \right) \left(\frac{n_{1,1} + n_{2,2}}{2} \right)^s \end{aligned}$$

Example 1 $\Pi = (\{a, b\}, [1]_1, abbbb, R)$, $R = \{p_1 : a \rightarrow aa, p_2 : b \rightarrow bb\}$. $p_1 = p_2 = p = 2$. $vM^s = (v_{1,1} + v_{1,2}s)2^s$. *Coefficients:* $v_{1,1} = (1, 5)$, $v_{1,2} = (1, 5) * (2E - 2E) = (0, 0)$. *Answer:*

$$vM^s = (2^s, 5 \cdot 2^s).$$

Example 2 (*Polynomial growth*) $\Pi = (\{a, b\}, [1]_1, aabbb, R)$, $R = \{p_1 : a \rightarrow abbbb, p_2 : b \rightarrow b\}$. $p_1 = p_2 = p = 1$. $vM^s = (v_{1,1} + v_{1,2}s)1^s$. *Coefficients:* $v_{1,1} = (2, 3)$, $v_{1,1} + v_{1,2} = (2, 11)$, so $v_{1,1} = (0, 4)$ *Answer:*

$$vM^s = (2, 3 + 4s).$$

Case 2: $p_1, p_2 = \frac{n_{1,1} + n_{2,2} \pm \sqrt{(n_{1,1} + n_{2,2})^2 - 4(n_{1,1}n_{2,2} - n_{1,2}n_{2,1})}}{2}$, $vM^s = v_{1,1}p_1^s + v_{2,1}p_2^s$. Coefficients: $v_{1,1} + v_{2,1} = v$, $p_1v_{1,1} + p_2v_{2,1} = vM$. (Notice that coefficients 1, 1, p_1, p_2 are the same for solving for $v_{i,1}[1]$ and for solving for $v_{i,1}[2]$) Solution: $\delta = p_2 - p_1$, $D_1 = -v(M - p_2E)$, $D_2 = v(M - p_1E)$, so $v_{1,1} = D_1/\delta$, $v_{2,1} = D_2/\delta$.

Putting it all together, if $n_{1,1} \neq n_{2,2}$ or $n_{1,2}n_{2,1} \neq 0$, then

$$\begin{aligned} vM^s &= \frac{v}{4\sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})}} \times \\ &\left(\left(2M - (n_{1,1} + n_{2,2} - \sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})})E \right) \right. \\ &\quad \left. \left(\frac{n_{1,1} + n_{2,2} + \sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})}}{2} \right)^s \right. \\ &\quad - \left. \left(2M - (n_{1,1} + n_{2,2} + \sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})})E \right) \right. \\ &\quad \left. \left(\frac{n_{1,1} + n_{2,2} - \sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})}}{2} \right)^s \right) \end{aligned}$$

Componentwise:

$$\begin{aligned} (vM^s)[i] &= \frac{1}{4\sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})}} \times \\ &\left(\left(2v[1]n_{i,1} + 2v[2]n_{i,2} - (n_{1,1} + n_{2,2} - \sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})}) \right) \right. \\ &\quad \left. \left(\frac{n_{1,1} + n_{2,2} + \sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})}}{2} \right)^s \right. \\ &\quad - \left. \left(2v[1]n_{i,1} + 2v[2]n_{i,2} - (n_{1,1} + n_{2,2} + \sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})}) \right) \right. \\ &\quad \left. \left(\frac{n_{1,1} + n_{2,2} - \sqrt{(n_{1,1} - n_{2,2})^2 - 4(n_{1,2}n_{2,1})}}{2} \right)^s \right), \quad 1 \leq i \leq 2. \end{aligned}$$

Example 3 $\Pi = (\{a, b\}, [1]_1, a, R)$, $R = \{p_1 : a \rightarrow bb, p_2 : b \rightarrow aaa\}$.

$D = -4(-2 \cdot 3)$. $p_1, p_2 = \pm 2\sqrt{6}/2$. $vM^s = v_{1,1}\sqrt{6}^s + v_{2,1}(-\sqrt{6})^s$. Coefficients: $v_{1,1} + v_{2,1} = (1, 0)$, $(v_{1,1} - v_{2,1})\sqrt{6} = (0, 2)$, hence, $v_{1,1} = (1/2, \sqrt{6}/3)$, $v_{2,1} = (1/2, -\sqrt{6}/3)$, Answer:

$$vM^s = (\sqrt{6})^s((1/2, \sqrt{6}/3) + (-1)^n(1/2, -\sqrt{6}/3)).$$

Thus, at step s rule p_1 will be applied $6^{s/2}(1 + (-1)^n)/2$ times, and rule p_2 will be applied $6^{(s-1)/2}(1 - (-1)^n)$ times.

Example 4 $\Pi = (\{a, b\}, [1]_1, a, R)$, $R = \{p_1 : a \rightarrow abb, p_2 : b \rightarrow aabbb\}$.

$D = (1 - 3)^2 + 4(2 \cdot 2) = 20$. $p_1, p_2 = 2 \pm \sqrt{5}$. $vM^s = v_{1,1}(2 + \sqrt{5})^s + v_{2,1}(2 - \sqrt{5})^s$. Coefficients: $v_{1,1} + v_{2,1} = (1, 0)$, $(2 + \sqrt{5})v_{1,1} + (2 - \sqrt{5})v_{2,1} = (0, 2)$, hence, $v_{1,1} = (\frac{1}{2} - \frac{1}{2\sqrt{5}}, \frac{1}{2} + \frac{1}{2\sqrt{5}})$, $v_{2,1} = (\frac{1}{\sqrt{5}}, -\frac{1}{\sqrt{5}})$, Answer:

$$vM^s = (2 + \sqrt{5})^s \left(\frac{1}{2} - \frac{1}{2\sqrt{5}}, \frac{1}{2} + \frac{1}{2\sqrt{5}} \right) + (2 - \sqrt{5})^s \left(\frac{1}{\sqrt{5}}, -\frac{1}{\sqrt{5}} \right).$$

These numbers actually correspond a subsequence of the famous Fibonacci numbers ($F(0) = 0$, $F(1) = 1$, $F(i + 2) = F(i + 1) + F(i)$, $i \geq 0$): $vM^s = (F(3s), F(3s + 1))$. The transition matrix is actually a third power of the Fibonacci one (corresponding to rules $a \rightarrow b$, $b \rightarrow ab$).

6 Conclusions

The Sevilla carpet of a deterministic non-cooperative P system is a numerical function (application multiplicity) with one argument (what rule rule) ranging over a finite set and the other argument (at what step) being numerical, from a potentially infinite set. With respect to the time, this function can have polynomial or exponential growth.

Considering non-deterministic systems is more difficult, since then one system defines a family of carpets (each corresponding to a computation). The behavior of cooperative systems is much more difficult to describe or to predict. Nevertheless, studying some properties of them is an interesting research topics.

Acknowledgements The first author is supported by the project TIC2002-04220-C03-02 of the Research Group on Mathematical Linguistics, Tarragona. The first author also acknowledges the Moldovan Research and Development Association (MRDA) and the U.S. Civilian Research and Development Foundation (CRDF), Award No. MM2-3034 for providing a challenging and fruitful framework for cooperation.

References

- [1] G. Ciobanu, Gh. Păun, Gh. Ștefănescu, Sevilla Carpets Associated with P Systems, *Brainstorming Week on Membrane Computing* (M. Cavaliere, C. Martn-Vide, Gh. Paun, eds.) GRLMC Report 26/03, Tarragona, 2003, 135–140.
- [2] G. Păun. Computing with membranes. *Journal of Computer and System Sciences*, 1(61):108–143, 2000. Also TUCS Report No. 208, 1998.
- [3] Gh.Păun, *Membrane computing. An Introduction*, Springer-Verlag, 2002.
- [4] G. Rozenberg, A. Salomaa, *The mathematical theory of L systems*, Academic Press, New York, 1980.
- [5] The P systems web page. <http://psystems.disco.unimib.it/>.