

SRT₂² DOES NOT IMPLY COH IN ω -MODELS

BENOIT MONIN AND LUDOVIC PATEY

ABSTRACT. In this article, we prove that every Δ_2^0 set has an infinite subset in it or its complement whose Turing jump is not of PA degree relative to \emptyset' . We use the relativized statement to build an ω -model of stable Ramsey's theorem for pairs (SRT₂²) which is not a model of the cohesiveness principle (COH). This answers a question of Chong, Slaman and Yang. The proof is an elaboration of Liu's theorem together with the second jump control of Cholak, Jockusch and Slaman.

1. INTRODUCTION

In this article, we prove that stable Ramsey's theorem for pairs does not imply the cohesiveness principle over ω -models¹. This answers questions of Cholak, Jockusch and Slaman [2] and Chong, Slaman and Yang [4]. Our main motivation comes from *reverse mathematics*.

1.1. Reverse mathematics and Ramsey's theorem

Reverse mathematics is a foundational program started by Harvey Friedman in 1975, whose goal is to find the weakest axioms needed to prove ordinary theorems. It uses the framework of second-order arithmetics, with a base theory, RCA₀, capturing “computable mathematics”. The early study of reverse mathematics revealed the existence of four linearly ordered big systems WKL, ACA, ATR, and Π_1^1 CA (in increasing order), such that, given an ordinary theorem, it is very likely either to be provable in RCA₀, or provably equivalent to one of the four systems in RCA₀. These systems together with RCA₀ are known as the “Big Five”. By its success in finding the exact axioms needed for the large majority of theorems and its foundational consequences, in particular its partial answer to Hilbert's program of finitistic reductionism [25], reverse mathematics is cited among the 100 key breakthroughs in mathematics [10]. Among the Big Five, WKL stands for “weak König's lemma”, and asserts that every infinite binary tree admits an infinite path, while ACA is the comprehension axiom restricted to arithmetical formulas. WKL can be thought of as capturing compactness arguments, and ACA is equivalent to the existence, for every set X , of the halting set relative to X . See Simpson [26] for an introduction to reverse mathematics.

Among the theorems studied in reverse mathematics, Ramsey's theorem received a special attention from the community, since Ramsey's theorem for pairs historically was the first theorem known to escape the Big Five phenomenon. Given a set of integers X , $[X]^n$ denotes the set of all n -tuples over X . For a coloring $f : [\omega]^n \rightarrow k$, a set of integers H is *homogeneous* if f is constant over $[H]^n$.

Statement (Ramsey's theorem). RT_kⁿ: “Every k -coloring of $[\omega]^n$ admits an infinite homogeneous set”.

In particular, RT_k¹ is the infinite pigeonhole principle for k -partitions. Ramsey's theorem and its consequences are notoriously hard to analyse from a computability-theoretic viewpoint. Jockusch [14] proved that RT_kⁿ is equivalent to ACA whenever $n \geq 3$, thereby showing that RT_kⁿ satisfies the Big Five phenomenon. The question whether RT_k² implies ACA was a longstanding open question, until Seetapun [24] proved that RT_k² is strictly weaker than ACA. Later, Jockusch [14, 15] and Liu [17] showed that RT_k² is incomparable with WKL, and therefore that RT_k² is not even linearly ordered with the Big Five. See Hirschfeldt [11] for an introduction to the reverse mathematics of Ramsey's theorem.

¹The authors are thankful to Damir Dzhafarov for insightful comments and discussions.

1.2. Stable Ramsey's theorem for pairs and cohesiveness

In order to understand better the computational and proof-theoretic content of Ramsey's theorem for pairs, Cholak, Jockusch and Slaman [2] decomposed it into two statements, namely, stable Ramsey's theorem for pairs, and cohesiveness. A coloring of pairs $f : [\omega]^2 \rightarrow k$ is *stable* if for every $x \in \omega$, $\lim_y f(\{x, y\})$ exists. An infinite set C is *cohesive* for a countable sequence of sets R_0, R_1, \dots if $C \subseteq^* R_i$ or $C \subseteq^* \bar{R}_i$ for every $i \in \omega$, where \subseteq^* means inclusion but for finitely many elements.

Statement (Stable Ramsey's theorem for pairs). SRT_k^2 : "Every stable k -coloring of $[\omega]^2$ admits an infinite homogeneous set".

Statement (Cohesiveness). COH : "Every countable sequence of sets has a cohesive set".

Cholak, Jockusch and Slaman [2] and Mileti [19] proved the equivalence over RCA_0 between RT_k^2 and $\text{SRT}_k^2 \wedge \text{COH}$. They naturally wondered whether this decomposition is non-trivial, in the sense that both statements SRT_k^2 and COH are strictly weaker than RT_k^2 . Hirschfeldt, Jockusch, Kjosso-Hanssen, Lempp and Slaman [13] partially answered the question by proving that COH is strictly weaker than RT_2^2 over RCA_0 . The question whether SRT_2^2 implies RT_2^2 over RCA_0 remained a long-standing open question. Since RT_2^2 is equivalent to $\text{SRT}_2^2 \wedge \text{COH}$, this is equivalent to the question whether SRT_2^2 implies COH over RCA_0 .

From a computability-theoretic viewpoint, stable Ramsey's theorem for pairs and two k colors is equivalent to combinatorially simpler statement called D_k^2 .

Statement. D_k^n : "For every Δ_n^0 k -partition of ω , there is an infinite subset of one of the parts".

Chong, Lempp and Yang [3], proved that the computable equivalence between RT_k^2 and D_k^2 also holds over RCA_0 . The cohesiveness principle also admits a nice computability-theoretic characterization. Jockusch and Stephan [16] proved that the sequence of all primitive recursive sets is a maximally difficult instance of COH . The cohesive sets for this sequence are called *p-cohesive* and their Turing degrees are precisely the ones whose jump is PA over \emptyset' , that is, the degrees whose jump can compute a path through any Δ_2^0 infinite binary tree. The following computability-theoretic question is therefore closely related to the previous question.

Question 1.1. Does every Δ_2^0 set has an infinite subset in it or its complement whose jump is not of PA degree over \emptyset' ?

One natural approach to separate SRT_2^2 from RT_2^2 would be to prove that every Δ_2^0 set admits an infinite subset G in it or its complement of low degree, that is, $G' \leq \emptyset'$. However, Downey, Hirschfeldt, Lempp and Solomon [6] constructed a Δ_2^0 set with no low infinite subset of it or its complement. Very surprisingly, Chong, Slaman and Yang [4] answered the SRT_2^2 vs COH question by constructing a model of $\text{RCA}_0 + \text{SRT}_2^2$ with only low sets, which is not a model of COH . The solution to this apparent paradox was the use of a non-standard model of RCA_0 in which Σ_2^0 induction fails. The sets of this model are low *within the model*, but not low in the meta-theory. The construction of Downey, Hirschfeldt, Lempp and Solomon [6] requires Σ_2^0 induction to be carried out.

Although the proof of Chong, Slaman and Yang [4] formally separated SRT_2^2 from COH over RCA_0 , the separation was not fully satisfactory, for two reasons. First, it leaves open the question whether $(\forall k) \text{SRT}_k^2$ implies COH which as also asked by Cholak, Jockusch and Slaman [2]. Indeed, $(\forall k) \text{SRT}_k^2$ implies Σ_2^0 induction, and therefore cannot have any models with only low sets. The second reasons is that the separations of Chong, Slaman and Yang [4] exploits the failure of a property which is known to hold in standard models. A structure in second-order arithmetics is a tuple $(M, \mathcal{S}, 0, 1, +, \times, <)$ where M denotes the set of integers, together with some constants 0 and 1, some binary operations $+$ and \times and an order relation $<$. \mathcal{S} is a collection of subsets of M representing the second-order part. Among these structures, we are particularly interested in those whose first-order part consists of the standard integers, together with their natural operations. These structures are called ω -structures, and are fully specified by their second-order part \mathcal{S} . Chong, Slaman and Yang [4] naturally asked the following question:

Question 1.2. Is every ω-model of $RCA_0 \wedge SRT_2^2$ a model of COH?

This question had an important impact in the development of reverse mathematics, and computability theory in general, not only by its self interest, but also by range of related questions, new techniques and intellectual emulation it generated in the community. Several articles are dedicated to this question [1, 3, 5, 8, 7, 9, 12, 20, 21, 22] and led to the rediscovery of Weihrauch degrees by Dorais, Dzhafarov, Hirst, Mileti and Shafer [5], and the design of the computable reduction by Dzhafarov [8]. Dzhafarov [8, 9] obtained partial separations by proving that COH is neither Weihrauch reducible, nor strongly computably reducible to SRT_2^2 . The most recent improvement is a proof by Dzhafarov and Patey [7] proving that COH is not Weihrauch reducible to SRT_2^2 even when a finitely many Turing functionals are allowed.

In this article, we answer the question by separating $RCA_0 \wedge SRT_2^2$ from COH on ω-models. For this, we prove the following main theorem.

Theorem 1.3 For every set Z whose jump is not of PA degree over \emptyset' and every $\Delta_2^{0,Z}$ set A , there is an infinite subset $G \subseteq A$ or $G \subseteq \bar{A}$ such that $(G \oplus Z)'$ is not of PA degree over \emptyset' .

This theorem is based on a second-jump control initially developed by Cholak, Jockusch and Slaman [2], and then successively refined by Wang [27], Patey [22] and Monin and Patey [20]. The techniques are combined with combinatorial ideas of Liu [17, 18]. Theorem 1.3 can be iterated to construct an ω-model of $RCA_0 \wedge SRT_2^2$ containing no set whose jump is of PA degree over \emptyset' , from which we deduce the following theorem.

Theorem 1.4 There is an ω-model of $RCA_0 \wedge SRT_2^2$ which is not a model of COH.

This answers a question of Chong, Slaman and Yang [4], but also of Cholak, Jockusch and Slaman [2] since any ω-model of SRT_2^2 is a model of $(\forall k) SRT_k^2$.

By being an adaptation and generalization of the first and second-jump control of Cholak, Jockusch and Slaman, the nature of the proof of Theorem 1.3 comforts the idea that this framework is the appropriate one for the computable and proof-theoretic analysis of Ramsey-like theorems. This gives a reasonable hope to prove general decidability theorems on Ramsey's theorem, in the spirit of [23]




FIGURE 1. Summary diagram of implications between statements over RCA_0 , and over ω-models. All the implications are strict, and the missing implications are separations.

1.3. Definitions and notation

A *binary string* is an ordered tuple of bits $a_0, \dots, a_{n-1} \in \{0, 1\}$. The empty string is written ϵ . A *binary sequence* (or a *real*) is an infinite listing of bits a_0, a_1, \dots . Given $s \in \omega$, 2^s is the set of binary strings of length s and $2^{<s}$ is the set of binary strings of length $< s$. As well, $2^{<\omega}$ is the set of binary strings and 2^ω is the set of binary sequences. Given a string $\sigma \in 2^{<\omega}$, we use $|\sigma|$ to denote its length. Given two strings $\sigma, \tau \in 2^{<\omega}$, σ is a *prefix* of τ (written $\sigma \preceq \tau$) if there exists a string $\rho \in 2^{<\omega}$ such that $\sigma\rho = \tau$. Given a sequence X , we write $\sigma \prec X$ if $\sigma = X \upharpoonright n$ for some $n \in \omega$. A binary string σ can be interpreted as a finite set $F_\sigma = \{x < |\sigma| : \sigma(x) = 1\}$. We write $\sigma \subseteq \tau$ for $F_\sigma \subseteq F_\tau$. We write $\#\sigma$ for the size of F_σ .

A *binary tree* is a set of binary strings $T \subseteq 2^{<\omega}$ which is closed downward under the prefix relation. A *path* through T is a binary sequence $P \in 2^\omega$ such that every initial segment belongs to T .

A *Turing ideal* \mathcal{I} is a collection of sets which is closed downward under the Turing reduction and closed under the effective join, that is, $(\forall X \in \mathcal{I})(\forall Y \leq_T X)Y \in \mathcal{I}$ and $(\forall X, Y \in \mathcal{I})X \oplus Y \in \mathcal{I}$, where $X \oplus Y = \{2n : n \in X\} \cup \{2n+1 : n \in Y\}$. A *Scott set* is a Turing ideal \mathcal{I} such that every infinite binary tree $T \in \mathcal{I}$ has a path in \mathcal{I} . In other words, a Scott set is the second-order part of an ω -model of $\text{RCA}_0 + \text{WKL}$. A countable Turing ideal \mathcal{M} is *coded* by a set X if $\mathcal{M} = \{X_n : n \in \omega\}$ with $X = \bigoplus_n X_n$. A formula is $\Sigma_1^0(\mathcal{M})$ (resp. $\Pi_1^0(\mathcal{M})$) if it is $\Sigma_1^0(X)$ (resp. $\Pi_1^0(X)$) for some $X \in \mathcal{M}$.

Given two sets A and B , we denote by $A < B$ the formula $(\forall x \in A)(\forall y \in B)[x < y]$. We write $A \subseteq^* B$ to mean that $A - B$ is finite, that is, $(\exists n)(\forall a \in A)(a \notin B \rightarrow a < n)$. A *k-cover* of a set X is a sequence of sets Y_0, \dots, Y_{k-1} such that $X \subseteq Y_0 \cup \dots \cup Y_{k-1}$.

2. BACKGROUND AND SKETCH OF THE PROOF

In this section, we give a sketch of the proof that every Δ_2^0 set A admits an infinite subset in it or its complement, whose jump is not of PA degree relative to \emptyset' . Many claims are formally proven in their full generality in Section 3. The proof is done by a variant of Mathias forcing with an effective second-jump control, that is, a notion of forcing whose forcing relation for Σ_2^0 and Π_2^0 formulas is Σ_2^0 and Π_2^0 , respectively. In the remainder of this section, fix a Δ_2^0 set A and let $A^0 = A$ and $A^1 = \bar{A}$.

2.1. First-jump control

Cholak, Jockusch and Slaman [2] designed a notion of forcing for constructing subsets of A^0 or A^1 , with a good first-jump control. This notion of forcing is a variant of Mathias forcing whose conditions are tuples (σ^0, σ^1, X) where $\sigma^0 \subseteq A^0$ and $\sigma^1 \subseteq A^1$ are finite strings representing the stem of the two sets $G^0 \subseteq A^0$ and $G^1 \subseteq A^1$ that we are building. The set $X \subseteq \omega$ is an infinite set belonging to some fixed Scott set \mathcal{M} , and serves as a reservoir of elements to add to the stems σ^0 and σ^1 . For example, by the low basis theorem, X can be chosen to be of low degree. We furthermore require that $\min X > \max(\sigma^0, \sigma^1)$. According to the intuition, a condition $d = (\tau^0, \tau^1, Y)$ *extends* a condition $c = (\sigma^0, \sigma^1, X)$ (written $d \leq c$) if the reservoir gets more restrictive, that is, $Y \subseteq X$, and if the stems are extended only with elements coming from the reservoir X , that is, $\sigma^i \preceq \tau^i$ and $\tau^i - \sigma^i \subseteq X \cap A^i$. The combinatorics of CJS provide a way to decide Σ_1^0 formulas without referring to the set A which is computationally too complex for the question.

Definition 2.1. Let $c = (\sigma^0, \sigma^1, X)$ be a condition, $i < 2$ and $\Phi_e(G, x)$ be a Δ_0 formula.

- (a) $c \Vdash^i (\exists x)\Phi_e(G, x)$ if there is some $x \in \omega$ such that $\Phi_e(\sigma^i, x)$ holds.
- (b) $c \Vdash^i (\forall x)\neg\Phi_e(G, x)$ if for every $x \in \omega$ and every $\rho \subseteq X$, $\neg\Phi_e(\sigma^i \cup \rho, x)$ holds.

Note that the set A does not appear in the definition of the forcing relation for Π_1^0 formulas. The forcing relation for Σ_1^0 and Π_1^0 formulas is therefore $\Sigma_1^0(X)$ and $\Pi_1^0(X)$, respectively, where X is the reservoir of the condition. There is no reason to consider that either a Σ_1^0 formula or its negation can be forced on each side of a condition. However, the following forcing question

which is at the heart of the CJS combinatorics ensures that this can be achieved on at least one side.

Definition 2.2. Given a condition $c = (\sigma^0, \sigma^1, X)$ and two Δ_0 formulas $\Phi_{e_0}(G, x)$ and $\Phi_{e_1}(G, x)$, define $c \text{ ?}\vdash (\exists x)\Phi_{e_0}(G^0, x) \vee (\exists x)\Phi_{e_1}(G^1, x)$ to hold if for every 2-cover $Z^0 \cup Z^1 = X$, there is some side $i < 2$, some $x \in \omega$ and some finite set $\rho \subseteq Z^i$ such that $\Phi_{e_i}(\sigma^i \cup \rho, x)$ holds.

The forcing question for Σ_1^0 formulas is also $\Sigma_1^0(X)$, and satisfies the following property.

Lemma 2.3 (Cholak, Jockusch and Slaman [2]) Let c be a condition, and $\Phi_{e_0}(G, x)$ and $\Phi_{e_1}(G, x)$ be two Δ_0 formulas.

- (a) If $c \text{ ?}\vdash (\exists x)\Phi_{e_0}(G^0, x) \vee (\exists x)\Phi_{e_1}(G^1, x)$, then there is some $d \leq c$ and some $i < 2$ such that $d \Vdash^i (\exists x)\Phi_{e_i}(G, x)$.
- (b) If $c \text{ ?}\not\vdash (\exists x)\Phi_{e_0}(G^0, x) \vee (\exists x)\Phi_{e_1}(G^1, x)$, then there is some $d \leq c$ and some $i < 2$ such that $d \Vdash^i (\forall x)\neg\Phi_{e_i}(G, x)$.

Proof. Suppose $c \text{ ?}\vdash (\exists x)\Phi_{e_0}(G^0, x) \vee (\exists x)\Phi_{e_1}(G^1, x)$ holds. Then letting $Z^0 = X \cap A^0$ and $Z^1 = X \cap A^1$, there is some side $i < 2$, some $x \in \omega$ and some finite set $\rho \subseteq X \cap A^i$ such that $\Phi_{e_i}(\sigma^i \cup \rho, x)$ holds. The condition $d = (\sigma^i \cup \rho, \sigma^{1-i}, X \cap (\max \rho, \infty))$ is an extension of c such that $d \Vdash^i (\exists x)\Phi_{e_i}(G, x)$.

Suppose now that $c \text{ ?}\not\vdash (\exists x)\Phi_{e_0}(G^0, x) \vee (\exists x)\Phi_{e_1}(G^1, x)$. Let \mathcal{P} be the collection of all the sets $Z^0 \oplus Z^1$ with $Z^0 \cup Z^1 = X$ such that for every $i < 2$, every $x \in \omega$ and every finite set $\rho \subseteq Z^i$, $\Phi_{e_i}(\sigma^i \cup \rho, x)$ does not hold. \mathcal{P} is a non-empty $\Pi_1^{0,X}$ class, so since X belongs to the Scott set \mathcal{M} , there is some 2-cover $Z^0 \cup Z^1 = X$ such that $Z^0 \oplus Z^1 \in \mathcal{P} \cap \mathcal{M}$. Let $i < 2$ be such that Z^i is infinite. Then the condition $d = (\sigma^0, \sigma^1, Z^i)$ is an extension of c such that $d \Vdash^i (\forall x)\neg\Phi_{e_i}(G, x)$. \square

By a pairing argument (if for every pair $m, n \in \omega$, $m \in A$ or $n \in B$, then $A = \omega$ or $B = \omega$), if a filter \mathcal{F} is sufficiently generic, there is some side i such that for every Σ_1^0 formula $\varphi(G)$, there is some $c \in \mathcal{F}$ such that $c \Vdash^i \varphi(G)$ or $c \Vdash^i \neg\varphi(G)$. Note that in the proof of Lemma 2.3, the new reservoir refining X is either X truncated by finitely many elements, or in the form $X \cap Z^0$ or $X \cap Z^1$ for some 2-cover $Z^0 \cup Z^1 = \omega$ such that $Z^0 \oplus Z^1 \in \mathcal{M}$.

2.2. Second-jump control

The forcing relation for Σ_2^0 formulas can be defined inductively by stating that $c \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ if $c \Vdash^i (\forall y)\Phi_e(G, x, y)$ for some $x \in \omega$. The relation $c \Vdash^i (\forall y)\Phi_e(G, x, y)$ is $\Pi_1^0(X)$ where X is the reservoir of c . In particular, whenever X is low, then the relation $c \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ is Σ_2^0 .

The definition of a forcing relation for Π_2^0 formulas is more problematic. A Π_2^0 formula $(\forall x)(\exists y)\neg\Phi_e(G, x, y)$ can be seen as a collection of Σ_1^0 properties $\langle (\exists y)\neg\Phi_e(G, x, y) : x \in \omega \rangle$ which all need to be forced. It is usually impossible to force infinitely many Σ_1^0 properties simultaneously, and one has to satisfy them one by one, by proving that for every $x \in \omega$, the set of collections forcing $(\exists y)\neg\Phi_e(G, x, y)$ is dense below c . The forcing relation $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$ is therefore naturally defined by the statement

$$(\forall x \in \omega)(\forall c_1 \leq c)(\exists c_2 \leq c_1)c_2 \Vdash^i (\exists y)\neg\Phi_e(G, x, y)$$

This definition of the forcing relation for Π_2^0 formulas is however too complex from a computational viewpoint. The main complexity comes from the description of the reservoir refinement, saying “there exists an infinite set $Y \in \mathcal{M}$ such that $Y \subseteq X$.”

Cholak, Jockusch and Slaman [2] went around this problem by observing that the only operations needed on the reservoirs to provide a good first-jump control were truncation and splitting. In some sense, the notion of forcing should not be considered as a variant of Mathias forcing since the refinement operation of the reservoirs does not need to be the arbitrary subset relation. The relation $c = (\sigma^0, \sigma^1, X) \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$ is then translated into the sentence “For every $x \in \omega$, every τ^0 and τ^1 extending σ^0 and σ^1 with elements from $X \cap A^0$ and $X \cap A^1$, respectively, the collection of reservoirs Y such that $(\tau^0, \tau^1, Y) \not\vdash (\forall y)\Phi_e(G, x, y)$

is *large*”, for some notion of largeness which enables truncation and splitting. We now give a modern presentation of the notion of forcing designed by Cholak, Jockusch and Slaman with a good second-jump control.

Definition 2.4. A class $\mathcal{A} \subseteq 2^\omega$ is a *largeness class* if it satisfies the following properties:

- (1) For every $X \in \mathcal{A}$ and $Y \supseteq X$, $Y \in \mathcal{A}$
- (2) For every $k \in \omega$ and every $X_0 \cup \dots \cup X_{k-1} \supseteq \omega$, there is some $j < k$ such that $X_j \in \mathcal{A}$

Fix a countable Scott set $\mathcal{M} = \{X_0, X_1, \dots\}$ coded by a set M of low degree, that is, $M = \bigoplus_i X_i$ and $M' \leq_T \emptyset'$. Such a Scott set exists by the Low Basis theorem [15]. Fix a uniformly M -computable enumeration $\mathcal{U}_0, \mathcal{U}_1, \dots$ of all the upward-closed $\Sigma_1^{0,X}$ sub-classes of 2^ω for every $X \in \mathcal{M}$. We are particularly interested in largeness classes in the form $\mathcal{U}_C = \bigcap_{e \in C} \mathcal{U}_e$ for some Δ_2^0 set of indices $C \subseteq \omega$. Given an infinite set $X \in \mathcal{M}$, we let \mathcal{L}_X be the largeness class of all $Z \subseteq \omega$ such that $Z \cap X$ is infinite.

Let us enrich the previous notion of forcing with a Δ_2^0 set of indices $C \subseteq \omega$ representing a largeness class of the form \mathcal{U}_C and to which the reservoir must belong. A forcing condition is therefore a tuple $(\sigma^0, \sigma^1, X, C)$ where $\sigma^0 \subseteq A^0$, $\sigma^1 \subseteq A^1$, X is an infinite set belonging to \mathcal{M} (in particular of low degree), $C \subseteq \omega$ is a Δ_2^0 set such that \mathcal{U}_C is a largeness class, and $\mathcal{U}_C \subseteq \mathcal{L}_X$.

Remark 2.5. Instead of asking $X \in \mathcal{U}_C$, we require the stronger fact that $\mathcal{U}_C \subseteq \mathcal{L}_X$. Since X is of low degree, \mathcal{L}_X can be put in the form \mathcal{U}_C for some Δ_2^0 set $C \subseteq \omega$. The requirement $X \in \mathcal{U}_C$ is not strong enough, as there might be some cover $Y_0 \cup \dots \cup Y_{k-1} \supseteq X$ such that $Y_j \notin \mathcal{U}_C$ for every $j < k$. There might also be some $D \supseteq C$ such that \mathcal{U}_D is a largeness sub-class of \mathcal{U}_C , but $X \notin \mathcal{U}_D$. By requiring that $\mathcal{U}_C \subseteq \mathcal{L}_X$, we ensure that every largeness subclass $\mathcal{U}_D \subseteq \mathcal{U}_C$ is a largeness class *within* X , that is, for every k and every k -cover $Y_0 \cup \dots \cup Y_{k-1} \supseteq X$, there is some $j < k$ such that $Y_j \in \mathcal{U}_D$. In particular $X \in \mathcal{U}_D$.

A condition (τ^0, τ^1, Y, D) extends $(\sigma^0, \sigma^1, X, C)$ if (τ^0, τ^1, Y) extends (σ^0, σ^1, X) as before, except that $D \supseteq C$, which means that $\mathcal{U}_D \subseteq \mathcal{U}_C$. The forcing relation for Σ_1^0 and Π_1^0 formulas is left unchanged. In particular, it does not depend on the largeness class \mathcal{U}_C .

Definition 2.6. Given a Δ_0 formula $\Phi_e(G, x, y)$, a finite set $\sigma \in 2^{<\omega}$ and some integer $x \in \omega$, let $\zeta(e, \sigma, x)$ be an index of the Σ_1^0 class

$$\mathcal{U}_{\zeta(e, \sigma, x)} = \{X : (\exists \rho \subseteq X)(\exists y \in \omega) \neg \Phi_e(\sigma \cup \rho, x, y)\}$$

In other words, $\mathcal{U}_{\zeta(e, \sigma, x)}$ is the collection of all reservoirs X such that the Mathias condition (σ, X) does not force $(\forall y)\Phi_e(G, x, y)$. We can now define a forcing relation for Σ_2^0 and Π_2^0 formulas whose complexities are Σ_2^0 and Π_2^0 , respectively.

Definition 2.7. Let $c = (\sigma^0, \sigma^1, X, C)$ be a condition, $i < 2$ and $\Phi_e(G, x, y)$ be a Δ_0 formula.

- (a) $c \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ if there is some $x \in \omega$ such that $c \Vdash^i (\forall y)\Phi_e(G, x, y)$ holds.
- (b) $c \Vdash^i (\forall x)(\exists y)\neg \Phi_e(G, x, y)$ if for every $x \in \omega$ and every $\rho \subseteq X \cap A^i$, $\zeta(e, \sigma^i \cup \rho, x) \in C$.

The interpretation of the forcing relation for Σ_2^0 formulas is immediate. The forcing relation for a Π_2^0 formula $(\forall x)(\exists y)\neg \Phi_e(G, x, y)$ ensures that for every $x \in \omega$ and every extension $d \leq c$, $d \not\Vdash^i (\forall y)\Phi_e(G, x, y)$. Indeed, if $d = (\tau^0, \tau^1, Y, D)$, then for every $x \in \omega$, $Y \in \mathcal{U}_D \subseteq \mathcal{U}_C \subseteq \mathcal{U}_{\zeta(e, \tau^i, x)}$.

Remark 2.8. Note that if a condition c forces a Σ_2^0 formula on a side $i < 2$, then the formula will hold on $G^i = \bigcup \{\sigma^i : (\sigma^0, \sigma^1, X, C) \in \mathcal{F}\}$ for every filter \mathcal{F} containing c . On the other hand, if c forces a Π_2^0 formula on side i , then the filter \mathcal{F} must be sufficiently generic for the property to hold on G^i . More precisely, the forcing relation for a formula $(\forall x)(\exists y)\neg \Phi_e(G, x, y)$ states that for every $x \in \omega$, the formula $(\forall y)\Phi_e(G, x, y)$ will never be forced. One can deduce that $(\exists y)\neg \Phi_e(G^i, x, y)$ whenever the side i is 1-generic, meaning that every Σ_1^0 formula or its negation is forced on side i . However, the disjunctive nature of the first-jump control guarantees only that at least one of the sides will be 1-generic. Therefore, one can ensure only on one side that the forced Π_2^0 formulas will actually hold.

Because of the assumption that A is Δ_2^0 , we can design a non-disjunctive forcing question for Σ_2^0 formulas, which will be Σ_2^0 . This enables us to prove that for *each* side, the set of conditions forcing a Σ_2^0 formula or its negation is dense. However, by the previous remark, the forced properties are only guaranteed to hold on one side.

Definition 2.9. Let $c = (\sigma^0, \sigma^1, X, C)$ be a condition, $i < 2$ and $\Phi_e(G, x, y)$ be a Δ_0 formula. Let $c \text{ ?}\vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ hold if

$$\mathcal{U}_C \cap \{\mathcal{U}_{\zeta(e, \sigma^i \cup \rho, x)} : x \in \omega, \rho \subseteq X \cap A^i\}$$

is not a largeness class.

As we will see in Lemma 3.2, the forcing question holds if and only if there is a finite set $F \subseteq X$ and some $n \in \omega$ such that the class $\mathcal{U}_F \cap \{\mathcal{U}_{\zeta(e, \sigma^i \cup \rho, x)} : x < n, \rho \subseteq X \cap A^i \upharpoonright n\}$ is not a largeness class. Note that the class is $\Sigma_1^{0, Z}$ for some $Z \in \mathcal{M}$. A complexity analysis for the forcing question shows that not being a largeness class for a $\Sigma_1^{0, Z}$ class is $\Sigma_2^{0, Z}$ (see Lemma 3.3), hence Σ_2^0 whenever Z is low. The forcing relation enjoys the following lemma, which shows in particular that every Σ_2^0 formula or its complement can be forced in *each* side.

Lemma 2.10 Let c be a condition, $i < 2$ and $\Phi_e(G, x, y)$ be a Δ_0 formula.

- (1) If $c \text{ ?}\vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$, then there is a $d \leq c$ such that $d \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$.
- (2) If $c \text{ ?}\not\vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$, then there is a $d \leq c$ such that $d \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$.

Proof. Say $c = (\sigma^0, \sigma^1, X, C)$.

Case 1: $c \text{ ?}\vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$. Then there is a finite set $F \subseteq C$ and some $n \in \omega$ such that the class $\mathcal{U}_F \cap \{\mathcal{U}_{\zeta(e, \sigma^i \cup \rho, x)} : x < n, \rho \subseteq X \cap A^i \upharpoonright n\}$ is not a largeness class. Since the class is $\Sigma_1^{0, Z}$ for some Z belonging to the Scott set \mathcal{M} , there is a k -cover $Y_0 \cup \dots \cup Y_{k-1} = \omega$ belonging to \mathcal{M} such that for every $j < k$, $Y_j \notin \mathcal{U}_F \cap \{\mathcal{U}_{\zeta(e, \sigma^i \cup \rho, x)} : x < n, \rho \subseteq X \cap A^i \upharpoonright n\}$. Let $j < k$ be such that $\mathcal{U}_C \cap \mathcal{L}_{X \cap Y_j}$ is a largeness class (see Lemma 3.7). In particular, there is some $x \in \omega$ and some $\rho \subseteq X \cap A^i$ such that $Y_j \notin \mathcal{U}_{\zeta(e, \sigma^i \cup \rho, x)}$, hence $(\sigma^i \cup \rho, x) \Vdash (\forall y)\Phi_e(G, x, y)$. Let $D \supseteq C$ be a Δ_2^0 set such that $\mathcal{U}_D = \mathcal{U}_C \cap \mathcal{L}_{X \cap Y_j}$. The condition $(\sigma^i \cup \rho, \sigma^{1-i}, X \cap Y_j - \{0, \dots, \max \rho\}, D)$ is the desired extension.

Case 2: $c \text{ ?}\not\vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$. Then let $D = C \cup \{\zeta(e, \sigma^i \cup \rho, x) : x \in \omega, \rho \subseteq X \cap A^i\}$. The condition $(\sigma^0, \sigma^1, X, D)$ is the desired extension. \square

One can use the combinatorics of largeness classes to provide a more direct proof that all the forced Π_2^0 formulas must hold on one of the sides for every sufficiently generic filter. We say that a condition $c = (\sigma^0, \sigma^1, X, C)$ is *i-valid* for some $i < 2$ if $X \cap A^i \in \mathcal{U}_C$. Since either $X \cap A^0$ or $X \cap A^1$ belongs to \mathcal{U}_C , every condition is *i-valid* for at least one $i < 2$. Moreover, *i-validity* is upward-closed under the extension relation, that is, if a condition d is *i-valid* and $d \leq c$, then c is *i-valid* as well. Therefore, for every filter \mathcal{F} , there is at least one side $i < 2$ such that every condition is *i-valid*. When a condition c is *i-valid*, one can make some progress on Σ_1^0 formulas on the *i*-th side, as states the following lemma.

Lemma 2.11 For every *i-valid* condition $c = (\sigma^0, \sigma^1, X, C)$ and every $\zeta(e, \sigma^i, x) \in C$, there is an extension $d \leq c$ such that $d \Vdash^i (\exists y)\neg\Phi_e(G, x, y)$.

Proof. Since c is *i-valid*, then $X \cap A^i \in \mathcal{U}_C \subseteq \mathcal{U}_{\zeta(e, \sigma^i, x)}$. Thus there is some $y \in \omega$ and some $\rho \subseteq X \cap A^i$ such that $\neg\Phi_e(\sigma^i \cup \rho, x, y)$ holds. The condition $d = (\sigma^i \cup \rho, \sigma^{1-i}, X - \{0, \dots, \max \rho\}, C)$ is the desired extension. \square

2.3. Forcing a jump of non-PA degree over \emptyset'

Now the main combinatorics of the second-jump control have been introduced, let us explain the core argument of forcing the jump of a solution not to be of PA degree over \emptyset' . A degree is PA over \emptyset' if and only if it computes a $\{0, 1\}$ -valued completion of $n \mapsto \Phi_n^{\emptyset'}(n)$.

Suppose we want to prove that for every Δ_2^0 set A , there is an infinite set $G \subseteq A$ or $G \subseteq \bar{A}$ such that G' does not compute a $\{0, 1\}$ -valued completion of $n \mapsto \Phi_n^{\theta'}$. We need the following notion of valuation.

Definition 2.12. A *valuation* is a partial function $p : \subseteq \omega \rightarrow 2$. A valuation is θ' -correct if $p(n) = \Phi_n^{\theta'} \downarrow$ for all $n \in \text{dom}(p)$. Two valuations p, q are *incompatible* if there is an $n \in \omega$ such that $p(n) \neq q(n)$.

Fix condition c , a side $i < 2$ and a Turing functional Γ . In order to force $\Gamma^{G'}$ not to be a completion of $n \mapsto \Phi_n^{\theta'}$, it is sufficient to find an extension $d \leq c$ such that one of the following holds:

- (1) $d \Vdash^i \Gamma^{G'} \not\subseteq p$ for some θ' -correct valuation p
- (2) $d \Vdash^i \Gamma^{G'} \subseteq p_0$ and $d \Vdash^i \Gamma^{G'} \subseteq p_1$ for two incompatible valuations p_0 and p_1

where $\Gamma^{G'} \not\subseteq p$ is the Σ_2^0 formula $(\exists n \in \text{dom } p) \Gamma^{G'}(n) \downarrow \neq p(n)$, and $\Gamma^{G'} \subseteq p$ is the Π_2^0 formula $(\forall n \in \text{dom } p) \Gamma^{G'}(n) \uparrow \vee \Gamma^{G'}(n) \downarrow = p(n)$. In particular, forcing the Σ_2^0 formula for a θ' -correct valuation ensures that $\Gamma^{G'}(n) \downarrow \neq \Phi_n^{\theta'} \downarrow$, while forcing the Π_2^0 formula for two incompatible valuations forces the partiality of $\Gamma^{G'}$. In both cases, we force $\Gamma^{G'}$ not to be a completion of $n \mapsto \Phi_n^{\theta'}$. The following lemma is adapted from a combinatorial lemma proven by Liu (Lemma 6.6 in [17]) and will be proven in Lemma 3.12.

Lemma 2.13 (Liu [17]) Let W be a θ' -c.e. set of valuations. Either W contains a θ' -correct valuation, or for every k , there are k pairwise incompatible valuations outside W .

Let us apply Lemma 2.13 on the following θ' -c.e. set of valuations

$$W = \{p : c \text{ ?}\Vdash^i \Gamma^{G'} \not\subseteq p\}$$

Case 1: $p \in W$ for some θ' -correct valuation p . Then $c \text{ ?}\Vdash^i \Gamma^{G'} \not\subseteq p$. By Lemma 2.10, there is an extension $d \leq c$ such that $d \Vdash^i \Gamma^{G'}(n) \not\subseteq p$.

Case 2: $W \cap \{p_0, p_1, p_2\} = \emptyset$ for three pairwise incompatible valuations p_0, p_1 and p_2 . Say $c = (\sigma^0, \sigma^1, X, C)$. By definition of W , $c \text{ ?}\Vdash^i \Gamma^{G'} \not\subseteq p_j$ for every $j < 3$. Unfolding the definition of the forcing question, for every $j < 3$, the class

$$\mathcal{U}_C \bigcap \{\mathcal{U}_{\zeta(e_j, \sigma^i \cup \rho, x)} : x \in \omega, \rho \subseteq X \cap A^i\}$$

is a largeness class, where $e_j \in \omega$ is an index of the Σ_2^0 formula $(\exists x)(\forall y) \Phi_{e_j}(G, x, y)$ which holds if $\Gamma^{G'} \not\subseteq p_j$. Let $C_j = C \cup \{\langle \zeta(e_j, \sigma^i \cup \rho, x), 0 \rangle : x \in \omega, \rho \subseteq X \cap A^i\}$. Although \mathcal{U}_{C_j} is a largeness class for every $j < 3$, in general, it is not true that there are $j_0 < j_1 < 3$ such that $\mathcal{U}_{C_{j_0}} \cap \mathcal{U}_{C_{j_1}}$ is a largeness class. However, the product $\mathcal{U}_{C_{j_0}} \times \mathcal{U}_{C_{j_1}}$ is a largeness class in the following generalized sense:

Definition 2.14. A class $\mathcal{A} \subseteq 2^\omega \times 2^\omega$ is a *largeness class* if

- (1) For every $\langle X_0, X_1 \rangle \in \mathcal{A}$ and $Y_0 \supseteq X_0$ and $Y_1 \supseteq X_1$, $\langle Y_0, Y_1 \rangle \in \mathcal{A}$
- (2) For every $k, \ell \in \omega$ and every $X_0 \cup \dots \cup X_{k-1} \supseteq \omega$, and $Y_0 \cup \dots \cup Y_{\ell-1} \supseteq \omega$, there is some $r < k$ and $s < \ell$ such that $\langle X_s, Y_\ell \rangle \in \mathcal{A}$.

Note that by taking the common refinement of the two covers, one can replace item (2) by “For every $k \in \omega$ and every $X_0 \cup \dots \cup X_{k-1} \supseteq \omega$, there is some $j_0, j_1 < k$ such that $\langle X_{j_0}, X_{j_1} \rangle \in \mathcal{A}$.” Again, given some pair $\langle X_0, X_1 \rangle$, we let $\mathcal{L}_{\langle X_0, X_1 \rangle}$ be the class of all pairs $\langle Y_0, Y_1 \rangle$ such that $Y_0 \cap X_0$ and $Y_1 \cap X_1$ are both infinite. The notion of largeness class over an arbitrary product and the \mathcal{L} notation is defined accordingly.

The cartesian product of two largeness classes is again a largeness class. However, some largeness classes over $2^\omega \times 2^\omega$ cannot be expressed as a cartesian product of two largeness classes over 2^ω , as witnessed by the class $\{\langle X, Y \rangle : |X \cap Y| = \infty\}$. The extension of c forcing partiality of $\Gamma^{G'}$ on side i is of the following type:

$$c = (\sigma_{j_0, j_1}^0, \sigma_{j_0, j_1}^1, X_0, X_1, X_2, D : j_0 < j_1 < 3)$$

where $\sigma_{j_0, j_1}^i \subseteq A^i$ for every $j_0 < j_1 < 3$ and $i < 2$, X_0, X_1, X_2 are sets of low degree with $\max(\sigma_{j_0, j_1}^0, \sigma_{j_0, j_1}^1) < \min(X_{j_0}, X_{j_1})$ for every $j_0 < j_1 < 3$. Moreover, D is a Δ_2^0 set of indices such that \mathcal{U}_D^3 is a largeness class over $2^\omega \times 2^\omega \times 2^\omega$, such that $\mathcal{U}_D^3 \subseteq \mathcal{L}_{\langle X_0, X_1, X_2 \rangle}$.

One can think of such a condition c from a partial order \mathbb{P} as three parallel conditions $c^{\{0,1\}}$, $c^{\{0,2\}}$ and $c^{\{1,2\}}$ from a partial order \mathbb{Q} where

$$c^{\{j_0, j_1\}} = (\sigma_{j_0, j_1}^0, \sigma_{j_0, j_1}^1, X_{j_0}, X_{j_1}, \pi_{\{j_0, j_1\}}(\mathcal{U}_D^3))$$

with $\pi_{\{j_0, j_1\}}(\mathcal{U}_D^3) = \{\langle Y_{j_0}, Y_{j_1} \rangle : \langle Y_0, Y_1, Y_2 \rangle \in \mathcal{U}_D^3\}$. Any such \mathbb{Q} -condition $c^{\{j_0, j_1\}}$ has two reservoirs X_{j_0} and X_{j_1} , both of which σ_{j_0, j_1}^0 and σ_{j_0, j_1}^1 take elements from. The forcing relation over \mathbb{Q} is defined as follows:

Definition 2.15. Given a Δ_0 formula $\Phi_e(G, x, y)$, a finite set $\sigma \in 2^{<\omega}$ and some integer $x \in \omega$, let $\zeta_2(e, \sigma, x)$ be an index of the Σ_1^0 class

$$\mathcal{U}_{\zeta_2(e, \sigma, x)}^2 = \{\langle X_0, X_1 \rangle : (\exists \rho \subseteq X_0 \cup X_1)(\exists y \in \omega) \neg \Phi_e(\sigma \cup \rho, x, y)\}$$

Definition 2.16. Let $c = (\sigma^0, \sigma^1, X_0, X_1, \mathcal{A})$ be a \mathbb{Q} -condition, $i < 2$ and $\Phi_e(G, x, y)$ be a Δ_0 formula.

- (a) $c \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ if there is some $x \in \omega$ such that $c \Vdash^i (\forall y)\Phi_e(G, x, y)$ holds.
- (b) $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$ if for every $x \in \omega$ and every $\rho \subseteq (X_0 \cup X_1) \cap A^i$, $\mathcal{A} \subseteq \mathcal{U}_{\zeta_2(e, \sigma^i \cup \rho, x)}^2$.

We need again to define a notion of i -validity for a \mathbb{Q} -condition to ensure that the forced Π_2^0 formulas hold for any sufficiently generic set. A \mathbb{Q} -condition $c = (\sigma^0, \sigma^1, X_0, X_1, \mathcal{A})$ is i -valid if $\langle X_0 \cap A^i, X_1 \cap A^i \rangle \in \mathcal{A}$. There is no reason to believe that every \mathbb{Q} -condition must have a valid side. However, by the pigeonhole principle, for every \mathbb{P} -condition $c = (\sigma_{j_0, j_1}^0, \sigma_{j_0, j_1}^1, X_0, X_1, X_2, D : j_0 < j_1 < 3)$, there is some $j_0 < j_1 < 3$ and some $i < 2$ such that $c^{\{j_0, j_1\}}$ is an i -valid \mathbb{Q} -condition. Indeed, since $\mathcal{U}_C^3 \subseteq \mathcal{L}_{\langle X_0, X_1, X_2 \rangle}$ is a largeness class, there are some $i_0, i_1, i_2 < 2$ such that $\langle X_0 \cap A^{i_0}, X_1 \cap A^{i_1}, X_2 \cap A^{i_2} \rangle \in \mathcal{U}_C^3$. In particular, there are some $j_0 < j_2 < 3$ be such that $i_{j_0} = i_{j_2}$. The \mathbb{Q} -condition $c^{\{j_0, j_2\}}$ is i_{j_0} -valid. The existence of a valid side is the reason we pick three pairwise incompatible valuations.

Having made the necessary definitions, consider the \mathbb{P} -condition

$$d = (\sigma_{j_0, j_1}^0, \sigma_{j_0, j_1}^1, X_0, X_1, X_2, D : j_0 < j_1 < 3)$$

where $\sigma_{j_0, j_1}^0 = \sigma^0$, $\sigma_{j_0, j_1}^1 = \sigma^1$, $X_0 = X_1 = X_2 = X$ and D is such that $\mathcal{U}_D^3 = \mathcal{U}_{C_0} \times \mathcal{U}_{C_1} \times \mathcal{U}_{C_2}$. The condition d is an extension of c such that for every $j_0 < j_1 < 3$

$$d^{\{j_0, j_1\}} \Vdash^i \Gamma^{G'}(n) \subseteq p_{j_0} \text{ and } d^{\{j_0, j_1\}} \Vdash^i \Gamma^{G'}(n) \subseteq p_{j_1}$$

We need to define a new forcing question for the generalized notion of \mathbb{P} -condition, in order to satisfy more requirements. The new notion of \mathbb{P} -condition admits multiple branches, namely, the \mathbb{Q} -conditions. Only one side of one branch is guaranteed to be valid. We therefore need to force the requirements on each side of each branch.

Given a \mathbb{P} -condition c and some side $i < 2$, we let $H(c, i)$ be the set of branches (here the set of pairs $\{j_0, j_1\} \in \{[0, 1, 2]\}^2$) such that the requirement is not forced on the side i of $c^{\{j_0, j_1\}}$. We design a forcing question parameterized by the set $H(c, i)$ such that if $c \Vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y)$ holds, then there is an extension $d \leq c$ which does not increase the number of branches, and such that $d^{\{j_0, j_1\}} \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ for some $\{j_0, j_1\} \in H$. On the other hand, if $c \not\Vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y)$ for sufficiently many e_j (which depends on the number of branches), then there is an extension $d \leq c$ which increases the number of branches, but such that for every new branch ν refining a branch $\{j_0, j_1\} \in H$, there are two indices $r \neq s$ such that $d^{\nu} \Vdash^i (\forall x)(\exists y)\neg\Phi_{e_r}(G, x, y)$ and $d^{\nu} \Vdash^i (\forall x)(\exists y)\neg\Phi_{e_s}(G, x, y)$ simultaneously.

In both cases, the number of branches for which the requirement is not forced decreases. In the first case, one more branch satisfies the Σ_2^0 outcome. In the second case, all the remaining branches satisfy the Π_2^0 outcome. Each time the Π_2^0 outcome occurs, then the number of branches increases.

Definition 2.17. Let $c = (\sigma_{j_0, j_1}^0, \sigma_{j_0, j_1}^1, X_0, X_1, X_2, C : j_0 < j_1 < 3)$ be a \mathbb{P} -condition, $i < 2$ and $\Phi_e(G, x, y)$ be a Δ_0 formula. Fix $H \subseteq \{0, 1, 2\}^2$. Let $c \text{ ? } \vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y)$ hold if

$$\mathcal{U}_C^3 \bigcap_{\{j_0, j_1\} \in H} \{ \langle X_0, X_1, X_2 \rangle : \langle X_{j_0}, X_{j_1} \rangle \in \mathcal{U}_{\zeta_2(e, \sigma_{j_0, j_1}^i \cup \rho, x)} : x \in \omega, \rho \subseteq (X_{j_0} \cap X_{j_1}) \cap A^i \}$$

is not a largeness class.

The \mathbb{P} and \mathbb{Q} notions of forcing have to be generalized to conditions with arbitrary many branches and reservoirs. The general case is formally defined and proven in the next section.

Remark 2.18. One important obstacle when using a variant of Mathias forcing to separate SRT_2^2 from COH on ω -models is that every sufficiently generic filter produces a solution to SRT_2^2 which is r-cohesive *as a set*. Indeed, given a condition (σ^0, σ^1, X) and one computable set R , either $X \cap R$ or $X \cap \overline{R}$ is infinite (or belongs to some largeness class). Then either $(\sigma^0, \sigma^1, X \cap R)$ or $(\sigma^0, \sigma^1, X \cap \overline{R})$ is an extension forcing both sets to be either almost included in R or in \overline{R} . In our situation, we overcome the problem by using increasingly many reservoirs simultaneously. Indeed, consider a condition $(\sigma^0, \sigma^1, X_0, X_1)$ where σ^0 and σ^1 take their elements from $X_0 \cup X_1$, and both X_0 and X_1 are required to be infinite. Then if only $X_0 \cap R$ and $X_1 \cap \overline{R}$ are infinite, the generic sets will have an infinite intersection with R and \overline{R} .

3. JUMP PA AVOIDANCE

In this section, we give a formal proof of the following main theorem outlined in Section 2:

Theorem 1.3. For every set Z whose jump is not of PA degree over \emptyset' and every Δ_2^0, Z set A , there is an infinite subset $G \subseteq A$ or $G \subseteq \overline{A}$ such that $(G \oplus Z)'$ is not of PA degree over \emptyset' .

Before proving Theorem 1.3, we prove its main consequence, namely, the separation of SRT_2^2 from COH on ω -models.

Theorem 1.4. There is an ω -model of $\text{RCA}_0 \wedge \text{SRT}_2^2$ which is not a model of COH .

Proof. By Theorem 1.3, there is a countable sequence of sets Z_0, Z_1, \dots such that for every $s \in \omega$, the jump of $Z_0 \oplus \dots \oplus Z_s$ is not of PA degree over \emptyset' , and for every $\Delta_2^0(Z_0 \oplus \dots \oplus Z_s)$ set A , there is some $t \in \omega$ such that $Z_t \subseteq A$ or $Z_t \subseteq \overline{A}$. Let $\mathcal{I} = \{X \in 2^\omega : (\exists s) X \leq_T Z_0 \oplus \dots \oplus Z_s\}$. The collection \mathcal{I} is a Turing ideal. Let \mathcal{M} be the ω -structure whose second-order part is \mathcal{I} . Every instance of SRT_2^2 in \mathcal{I} has a solution in \mathcal{I} , so \mathcal{M} is an ω -model of SRT_2^2 . Moreover, \mathcal{I} does not contain any set whose jump is of PA degree over \emptyset' . By Jockusch and Stephan [16], \mathcal{I} does not contain any p-cohesive set, so \mathcal{M} is not a model of COH . \square

The remainder of this section is devoted to the proof of Theorem 1.3. In what follows, fix a Δ_2^0 set A , and let $A^0 = A$ and $A^1 = \overline{A}$. Fix also a countable Scott set \mathcal{M} , coded by a low set M , as in the previous section.

3.1. Largeness classes

The following notion of largeness class was introduced by the authors in [20] to design a notion of forcing controlling the second jump of solutions to the pigeonhole principle. In what follows, given two sets A and B , we denote by $A \rightarrow B$ the class of all functions from A to B .

Definition 3.1. Fix a finite set $I \subseteq \omega^{<\omega}$. A *largeness class* is a collection of sets $\mathcal{A} \subseteq I \rightarrow 2^\omega$ such that

- (a) If $\langle X_\nu : \nu \in I \rangle \in \mathcal{A}$ and $Y_\nu \supseteq X_\nu$ for every $\nu \in I$, then $\langle Y_\nu : \nu \in I \rangle \in \mathcal{A}$
- (b) For every k -cover Y_0, \dots, Y_{k-1} of ω , there is some $\langle j_\nu < k : \nu \in I \rangle$ such that $\langle Y_{j_\nu} : \nu \in I \rangle \in \mathcal{A}$.

Whenever $I = \{\epsilon\}$, we identify the class $\{\epsilon\} \rightarrow 2^\omega$ with the class 2^ω . This yields a notion of largeness class for subsets of 2^ω . The collection of all the infinite sets is a largeness class. Moreover, any superclass of a largeness class is again a largeness class.

Given $I \subseteq 2^{<\omega}$, we fix a uniformly M -computable enumeration $\mathcal{U}_0^I, \mathcal{U}_1^I, \dots$ of all the $\Sigma_1^{0,Z}$ subclasses of $I \rightarrow 2^\omega$, upward-closed under the superset relation, where $Z \in \mathcal{M}$. Here, the upward-closure means that if $\langle X_\nu : \nu \in I \rangle \in \mathcal{U}_e^I$ and $Y_\nu \supseteq X_\nu$ for every $\nu \in I$, then $\langle Y_\nu : \nu \in I \rangle \in \mathcal{U}_e^I$. Given a set $C \subseteq \omega$, we write

$$\mathcal{U}_C^I = \bigcap_{e \in C} \mathcal{U}_e^I$$

If C is Δ_2^0 , then \mathcal{U}_C^I is Π_2^0 .

Lemma 3.2 Suppose $\mathcal{A}_0 \supseteq \mathcal{A}_1 \supseteq \dots$ is a decreasing sequence of largeness classes. Then $\bigcap_s \mathcal{A}_s$ is a largeness class.

Proof. If $\langle X_\nu : \nu \in I \rangle \in \bigcap_s \mathcal{A}_s$ and $Y_\nu \supseteq X_\nu$ for every $\nu \in I$, then for every s , since \mathcal{A}_s is a largeness class, $\langle Y_\nu : \nu \in Y \rangle \in \mathcal{A}_s$, so $\langle Y_\nu : \nu \in Y \rangle \in \bigcap_s \mathcal{A}_s$. Let Y_0, \dots, Y_{k-1} be a k -cover of ω . For every $s \in \omega$, there is some $\langle j_\nu < k : \nu \in I \rangle$ such that $\langle Y_{j_\nu} : \nu \in I \rangle \in \mathcal{A}_s$. By the infinite pigeonhole principle, there is some $\langle j_\nu < k : \nu \in I \rangle$ such that $\langle Y_{j_\nu} : \nu \in I \rangle \in \mathcal{A}_s$ for infinitely many s . Since $\mathcal{A}_0 \supseteq \mathcal{A}_1 \supseteq \dots$ is a decreasing sequence, $\langle Y_{j_\nu} : \nu \in I \rangle \in \bigcap_s \mathcal{A}_s$. \square

Lemma 3.2 has several very useful consequences. In particular, if \mathcal{U}_C is not a largeness class, then by Lemma 3.2, there is a finite set $F \subseteq C$ such that the class \mathcal{U}_F is not a largeness class. The set F being finite, the class \mathcal{U}_F is $\Sigma_1^{0,Z}$ for some $Z \in \mathcal{M}$, so since \mathcal{M} is a Scott set, there is a k -cover $X_0 \cup \dots \cup X_{k-1} = \omega$ in \mathcal{M} such that for every $j < k$, $X_j \notin \mathcal{U}_F \supseteq \mathcal{U}_C$. Therefore we can always find a k -cover belonging to \mathcal{M} , witnessing that \mathcal{U}_C is not a largeness class, whatever the complexity of the set C . Another consequence is the following lemma.

Lemma 3.3 Let \mathcal{A} be a Σ_1^0 class. The sentence “ \mathcal{A} is a largeness class” is Π_2^0 .

Proof. Say $\mathcal{A} = \{ \langle X_\nu : \nu \in I \rangle : (\exists \vec{\sigma} \preceq \vec{X}) \varphi(\vec{\sigma}) \}$ where φ is a Σ_1^0 formula. By compactness, \mathcal{A} is a largeness class iff for every $\vec{\sigma} = \langle \sigma_\nu : \nu \in I \rangle$ and $\vec{\tau} = \langle \tau_\nu : \nu \in I \rangle$ such that $\sigma_\nu \subseteq \tau_\nu$ for every $\nu \in I$ and $\varphi(\vec{\sigma})$ holds, $\varphi(\vec{\tau})$ holds, and for every k , there is some $n \in \omega$ such that for every $\sigma_0 \cup \dots \cup \sigma_{k-1} = \{0, \dots, n\}$, there is some $\langle j_\nu < k : \nu \in I \rangle$ such that $\varphi(\langle \sigma_{j_\nu} : \nu \in I \rangle)$ holds. \square

Definition 3.4. Given $\langle X_\nu : \nu \in I \rangle$, we let

$$\mathcal{L}_{\langle X_\nu : \nu \in I \rangle} = \{ \langle Y_\nu : \nu \in I \rangle : (\forall \nu \in I) |Y_\nu \cap X_\nu| = \infty \}$$

The following trivial lemma is very useful.

Lemma 3.5 Let $\langle X_\nu : \nu \in I \rangle$ and $\langle Y_\nu : \nu \in I \rangle$ be such that $X_\nu =^* Y_\nu$ for every $\nu \in I$. Then $\mathcal{L}_{\langle X_\nu : \nu \in I \rangle} = \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$.

Proof. By symmetry, it suffices to prove that $\mathcal{L}_{\langle X_\nu : \nu \in I \rangle} \subseteq \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$. Fix $\langle Z_\nu : \nu \in I \rangle \in \mathcal{L}_{\langle X_\nu : \nu \in I \rangle}$. Then for every $\nu \in I$, $Z_\nu \cap X_\nu$ is infinite. Since $X_\nu =^* Y_\nu$, then $Z_\nu \cap Y_\nu$ is infinite, so $\langle Z_\nu : \nu \in I \rangle \in \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$. \square

Lemma 3.6 Let \mathcal{A} be a largeness class. The class

$$\mathcal{L}(\mathcal{A}) = \{ \langle X_\nu : \nu \in I \rangle \in \mathcal{A} : \mathcal{A} \cap \mathcal{L}_{\langle X_\nu : \nu \in I \rangle} \text{ is a largeness class} \}$$

is a largeness subclass of \mathcal{A} .

Proof. The class $\mathcal{L}(\mathcal{A})$ is trivially a subclass of \mathcal{A} . Moreover, $\mathcal{L}(\mathcal{A})$ is upward-closed. Suppose for the contradiction that $\mathcal{L}(\mathcal{A})$ is not a largeness class. Then there is a cover $X_0 \cup \dots \cup X_{k-1} = \omega$ such that for every $\langle j_\nu < k : \nu \in I \rangle$, $\langle X_{j_\nu} : \nu \in I \rangle \notin \mathcal{L}(\mathcal{A})$. In other words, for every $\langle j_\nu < k : \nu \in I \rangle$, $\mathcal{A} \cap \mathcal{L}_{\langle X_{j_\nu} : \nu \in I \rangle}$ is not a largeness class. Thus for every $\langle j_\nu < k : \nu \in I \rangle$, there is a cover $Y_0 \cup \dots \cup Y_{\ell-1} = \omega$ such that for every $\langle i_\nu < \ell : \nu \in I \rangle$, $\langle Y_{i_\nu} : \nu \in I \rangle \notin \mathcal{A} \cap \mathcal{L}_{\langle X_{j_\nu} : \nu \in I \rangle}$. By taking the common refinement of all these covers, there is a cover $Z_0 \cup \dots \cup Z_{n-1} = \omega$ such that for every $\langle j_\nu < k : \nu \in I \rangle$, $\langle Z_{j_\nu} : \nu \in I \rangle \notin \mathcal{A} \cap \mathcal{L}_{\langle X_{j_\nu} : \nu \in I \rangle}$. Since \mathcal{A} is a largeness class,

there is some $\langle j_\nu < k : \nu \in I \rangle$ such that $\langle Z_{j_\nu} : \nu \in I \rangle \in \mathcal{A}$. Then $\langle Z_{j_\nu} : \nu \in I \rangle \in \mathcal{L}_{\langle Z_{j_\nu} : \nu \in I \rangle}$. Contradiction. \square

Lemma 3.7 If $\mathcal{A} \cap \mathcal{L}_{\langle X_\nu : \nu \in I \rangle} \cap \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$ is a largeness class, then so is $\mathcal{A} \cap \mathcal{L}_{\langle X_\nu \cap Y_\nu : \nu \in I \rangle}$.

Proof. $\mathcal{A} \cap \mathcal{L}_{\langle X_\nu \cap Y_\nu : \nu \in I \rangle}$ is upward-closed. Let $Z_0 \cup \dots \cup Z_{k-1} = \omega$. By refining the covering, we can assume that for every $j < k$ and $\nu \in I$, $Z_j \subseteq X_\nu$ or $Z_j \cap X_\nu = \emptyset$, and $Z_j \subseteq Y_\nu$ or $Z_j \cap Y_\nu = \emptyset$. Since $\mathcal{A} \cap \mathcal{L}_{\langle X_\nu : \nu \in I \rangle} \cap \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$ is a largeness class, there is some $\langle j_\nu < k : \nu \in I \rangle$ such that $\langle Z_{j_\nu} : \nu \in I \rangle \in \mathcal{A} \cap \mathcal{L}_{\langle X_\nu : \nu \in I \rangle} \cap \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$. We claim that $Z_{j_\nu} \subseteq X_\nu \cap Y_\nu$ for every $\nu \in I$. Indeed, since $\langle Z_{j_\nu} : \nu \in I \rangle \in \mathcal{L}_{\langle X_\nu : \nu \in I \rangle}$, $Z_{j_\nu} \cap X_\nu \neq \emptyset$, so $Z_{j_\nu} \subseteq X_\nu$. Similarly, since $\langle Z_{j_\nu} : \nu \in I \rangle \in \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$, $Z_{j_\nu} \subseteq Y_\nu$. Thus $\langle Z_{j_\nu} : \nu \in I \rangle \in \mathcal{A} \cap \mathcal{L}_{\langle X_\nu \cap Y_\nu : \nu \in I \rangle}$. \square

Definition 3.8. Given a class $\mathcal{A} \subseteq I \rightarrow 2^\omega$ and a set $J \subseteq I$, define the projection $\pi_J(\mathcal{A})$ be the set of all $\langle X_\nu : \nu \in J \rangle$ such that the class

$$\{\langle X_\nu : \nu \in I - J \rangle : \langle X_\nu : \nu \in I \rangle \in \mathcal{A}\}$$

is a largeness class.

Lemma 3.9 If $\mathcal{A} \subseteq I \rightarrow 2^\omega$ is a largeness class and $J \subseteq I$, then $\pi_J(\mathcal{A})$ is a largeness class.

Proof. The class $\pi_J(\mathcal{A})$ is upward-closed by upward-closure of \mathcal{A} . Let $Y_0 \cup \dots \cup Y_{k-1} = \omega$. Suppose for the contradiction that for every $\langle j_\nu : \nu \in J \rangle$, $\langle Y_{j_\nu} : \nu \in J \rangle \notin \pi_J(\mathcal{A})$. Thus for every $\langle j_\nu : \nu \in J \rangle$, the class

$$\{\langle X_\nu : \nu \in I - J \rangle : \langle X_\nu : \nu \in I - J \rangle \cup \langle Y_{j_\nu} : \nu \in J \rangle \in \mathcal{A}\}$$

is not a largeness class. By taking the common refinement of $Y_0 \cup \dots \cup Y_{k-1} = \omega$ with all the covers of ω witnessing that the classes above are not largeness classes, we obtain a cover $Z_0 \cup \dots \cup Z_{\ell-1} = \omega$ witnessing that \mathcal{A} is not a largeness class. \square

Lemma 3.10 Let $\mathcal{U}_C^I \subseteq I \rightarrow 2^\omega$ be a largeness class for some Δ_2^0 set C , and $\mathcal{A} \subseteq \pi_J(\mathcal{U}_C^I)$ be a Π_2^0 largeness class. Then there is a Δ_2^0 set $D \supseteq C$ such that $\mathcal{U}_D^I \subseteq \mathcal{U}_C^I$ is a largeness class and $\pi_J(\mathcal{U}_D^I) = \mathcal{A}$.

Proof. Let \mathcal{U}_D^I be the class of all $\langle X_\nu : \nu \in I \rangle \in \mathcal{U}_C^I$ such that $\langle X_\nu : \nu \in J \rangle \in \mathcal{A}$. Since \mathcal{A} is a Π_2^0 class, then so is \mathcal{U}_D^I , and therefore D can be chosen to be Δ_2^0 . Furthermore we can assume without loss of generality that $D \supseteq C$. By construction, $\mathcal{U}_D^I \subseteq \mathcal{U}_C^I$.

We claim that \mathcal{U}_D^I is a largeness class. \mathcal{U}_D^I is upward-closed since both \mathcal{U}_C^I and \mathcal{A} are. Let $Y_0 \cup \dots \cup Y_{k-1} = \omega$. Since \mathcal{A} is a largeness subclass of $J \rightarrow 2^\omega$, there is some $\langle j_\nu : \nu \in J \rangle$ such that $\langle Y_{j_\nu} : \nu \in J \rangle \in \mathcal{A}$. Since $\mathcal{A} \subseteq \pi_J(\mathcal{U}_C^I)$, the collection

$$\{\langle X_\nu : \nu \in I - J \rangle : \langle X_\nu : \nu \in I - J \rangle \cup \langle Y_{j_\nu} : \nu \in J \rangle \in \mathcal{U}_C^I\}$$

is a largeness class. Therefore, there is some $\langle j_\nu : \nu \in I - J \rangle$ such that $\langle Y_{j_\nu} : \nu \in I \rangle \in \mathcal{U}_C^I$. In particular, $\langle Y_{j_\nu} : \nu \in I \rangle \in \mathcal{U}_D^I$. This proves that \mathcal{U}_D^I is a largeness class.

Last, it is immediate to see that $\pi_J(\mathcal{U}_D^I) = \mathcal{A}$. \square

3.2. Valuation

The notion of valuation is a combinatorial trick of Liu [17] to obtain, whenever the Σ_2^0 outcome cannot be satisfied, arbitrarily many Π_2^0 formulas such that forcing any two of them is sufficient to force partiality of a Turing functional. We now define the notion of valuation and prove Liu's combinatorial lemma in its full generality (Lemma 3.12).

Definition 3.11. A *valuation* is a partial function $p \subseteq \omega \rightarrow 2$. A valuation is *X-correct* if $p(n) = \Phi_n^X(n) \downarrow$ for all $n \in \text{dom}(p)$. Two valuations p, q are *incompatible* if there is an $n \in \omega$ such that $p(n) \neq q(n)$.

The following lemma is adapted from Lemma 6.6 in Liu [17].

Lemma 3.12 (Liu [17]) Fix X and $Y \geq_T X$ such that Y is not of PA degree relative to X . Let W be a Y -c.e. set of valuations. Either W contains an X -correct valuation, or for every k , there are k pairwise incompatible valuations outside W .

Proof. Suppose W does not contain any X -correct valuation, otherwise we are done. Let S be the collection of all finite sets F such that for each $n \notin F$, either $\Phi_n^X(n) \downarrow$ or there is a valuation $p \in W$ such that $F \cup \{n\} \subseteq \text{dom } p$ and for every $m \in \text{dom } p \setminus (F \cup \{n\})$, we have $p(m) = \Phi_m^X(m) \downarrow$. If $F \notin S$, there there is at least one $n \notin F$ such that the above does not hold. We say that any such n witnesses $F \notin S$.

First suppose for the contradiction that $\emptyset \in S$. Then for each n , either $\Phi_n^X(n) \downarrow$ or there is a valuation $p \in W$ such that $n \in \text{dom } p$, and for every $m \in \text{dom } p \setminus \{n\}$, $\Phi_m^X(m) \downarrow$. We can then define a Y -computable completion h of $n \mapsto \Phi_n^X(n)$ as follows. Given n , wait until either $\Phi_n^X(n) \downarrow$, in which case let $h(n) = \Phi_n^X(n)$, or a p as above enters W , in which case we let $h(n) = 1 - p(n)$. Since W does not contain any valuation, in the latter case, if $\Phi_n^X(n) \downarrow$ then $\Phi_n^X(n) \neq p(n)$, so $h(n) = \Phi_n^X(n)$. Since Y is not of PA degree over X , this case cannot occur, so $\emptyset \notin S$.

Let n_0 witness the fact that $\emptyset \notin S$. Given n_0, \dots, n_j , if $\{n_0, \dots, n_j\} \notin S$, then let n_{j+1} witness this fact. Note that if n_j is defined, then $\Phi_{n_j}^X(n_j) \uparrow$.

Suppose for the contradiction that $\{n_0, \dots, n_j\} \in S$. Then $\{n_0, \dots, n_{j-1}\} \notin S$, otherwise n_j would not be defined. We can then define a Y -computable completion h of $n \mapsto \Phi_n^X(n)$ as follows. First, let $h(n_\ell) = 0$ for $\ell \leq j$. Given $n \notin \{n_0, \dots, n_j\}$, we wait until either $\Phi_n^X(n) \downarrow$, in which case we let $h(n) = \Phi_n^X(n)$, or a valuation p enters W such that $\{n_0, \dots, n_j, n\} \subseteq \text{dom } p$ and for every $m \in \text{dom } p \setminus \{n_0, \dots, n_j, n\}$, $p(m) = \Phi_m^X(m) \downarrow$. If $\Phi_n^X(n) \uparrow$, then the latter case must occur, since $\{n_0, \dots, n_j\} \in S$. In this case, we cannot have $p(n) = \Phi_n^X(n)$, as then p would be a counter-example to the fact that n_j witnesses $\{n_0, \dots, n_{j-1}\} \notin S$. Thus we can let $h(n) = 1 - p(n)$. We again have a contradiction since Y is not of PA degree over X .

Thus $\{n_0, \dots, n_j\} \notin S$ for all j . There are 2^{j+1} pairwise incompatible valuations with domain $\{n_0, \dots, n_j\}$. None of them can be in W , as this would contradict the fact that n_j witnesses $\{n_0, \dots, n_{j-1}\} \notin S$. This completes the proof. \square

Lemma 3.13 Let G be a set such that for every Δ_0 formula $\Phi_e(G, x, y, p)$, where x and y are integer variables and p is a valuation variable, either $(\exists x)(\forall y)\Phi_e(G, x, y, p)$ holds for some \emptyset' -correct valuation p , or $(\forall x)(\exists y)\neg\Phi_e(G, x, y, p_0)$ and $(\forall x)(\exists y)\neg\Phi_e(G, x, y, p_1)$ hold for two incompatible valuations p_0 and p_1 . Then G' is not PA over \emptyset' .

Proof. Suppose for the contradiction that G' is of PA degree over \emptyset' . In particular, there is a Turing functional Γ such that $\Gamma^{G'}$ is a $\{0, 1\}$ -valued completion of $n \mapsto \Phi_n^{\emptyset'}(n)$. Given $n, s \in \omega$, we denote by $\Gamma^{G'}(n)[s]$ the G -computable s -approximation of $\Gamma^{G'}(n)$. Let $\Phi_e(G, x, y, p)$ hold if there is some $n \in \text{dom } p$ such that if $\Gamma^{G'}(n)[x + y] \downarrow$ then $\Gamma^{G'}(n)[x + y] \neq p(n)$. We have two cases:

Case 1: $(\exists x)(\forall y)\Phi_e(G, x, y, p)$ holds for some \emptyset' -correct valuation p . Then there is some $n \in \text{dom}(p)$ such that $\Gamma^{G'}(n) \uparrow$ or $\Gamma^{G'}(n) \downarrow \neq p(n)$. By definition of a \emptyset' -correct valuation, $\Phi_n^{\emptyset'}(n) \downarrow = p(n)$ for every $n \in \text{dom}(p)$. Thus $\Gamma^{G'}$ is not a completion of $n \mapsto \Phi_n^{\emptyset'}(n)$.

Case 2: $(\forall x)(\exists y)\neg\Phi_e(G, x, y, p_0)$ and $(\forall x)(\exists y)\neg\Phi_e(G, x, y, p_1)$ hold for two incompatible valuations p_0 and p_1 . Then $\Gamma^{G'} \upharpoonright \text{dom } p_0 \subseteq p_0$ and $\Gamma^{G'} \upharpoonright \text{dom } p_1 \subseteq p_1$. Since p_0 and p_1 are incompatible, then $\Gamma^{G'}$ is partial. \square

3.3. Index set

We now define an ordered structure of sets of indices to simplify branches refinement whenever the Π_2^0 outcome occurs. Define the sequence of integers u_0, u_1, \dots inductively by $u_0 = 1$ and $u_{n+1} = \binom{2u_n+1}{2} u_n$.

Definition 3.14. Given $n \in \omega$, the n -index set is defined inductively as follows. The 0-index set \mathcal{I}_0 is a the singleton empty string $\{\epsilon\}$. Let \mathcal{I}_n be the n -index set. The $(n + 1)$ -index set is

the set

$$\mathcal{I}_{n+1} = (2u_n + 1) \times \mathcal{I}_n = \{x \hat{\ } \nu : x \leq 2u_n, \nu \in \mathcal{I}_n\}$$

Definition 3.15. Given $n \in \omega$, an n -index is defined inductively as follows. The unique 0-index is the singleton empty string $\{\epsilon\}$. Given an n -index $I \subseteq \mathcal{I}_n$, an $(n+1)$ -index is a set $\{x \hat{\ } \nu : \nu \in I\} \cup \{y \hat{\ } \nu : \nu \in I\}$ for some $x < y \leq 2u_n$.

Note that in particular, an n -index is a subset of \mathcal{I}_n . We write $I \triangleleft \mathcal{I}_n$ to say that I is an n -index.

Lemma 3.16 For every $n \in \omega$, $|\{I \subseteq \mathcal{I}_n : I \triangleleft \mathcal{I}_n\}| = u_n$.

Proof. By induction over n . Case $n = 0$. There is only one 0-index and $u_0 = 1$. Suppose $|\{I \subseteq \mathcal{I}_n : I \triangleleft \mathcal{I}_n\}| = u_n$. Then $|\{J \subseteq \mathcal{I}_{n+1} : J \triangleleft \mathcal{I}_{n+1}\}| = \binom{2u_n+1}{2} \cdot |\{I \subseteq \mathcal{I}_n : I \triangleleft \mathcal{I}_n\}| = \binom{2u_n+1}{2} u_n = u_{n+1}$. \square

The following lemma is the main combinatorial lemma of indices, which will be used in Lemma 3.32 to prove that every \mathbb{P} -condition admits a branch with a valid side.

Lemma 3.17 For every $n \in \omega$ and every 2-cover $B_0 \cup B_1 = \mathcal{I}_n$, there is an n -index $I \triangleleft \mathcal{I}_n$ and some $i < 2$ such that $I \subseteq B_i$.

Proof. By induction on n . The case $n = 0$ is trivial. Assume it holds for n . We prove it for $n+1$. For every $x \leq 2u_n$ and $i < 2$, let $B_{x,i} = \{\nu : x \hat{\ } \nu \in B_i\}$. By induction hypothesis, there is some $I_x \triangleleft \mathcal{I}_n$ and $i_x < 2$ such that $I_x \subseteq B_{x,i_x}$. By Lemma 3.16, $|\{I \subseteq \mathcal{I}_n : I \triangleleft \mathcal{I}_n\}| = u_n$ so by the pigeonhole principle, there is some $x < y \leq 2u_n$ such that $I_x = I_y$ and $i_x = i_y$. The $(n+1)$ -index $\{x \hat{\ } \nu : \nu \in I_x\} \cup \{y \hat{\ } \nu : \nu \in I_x\}$ is included in B_{i_x} . \square

Definition 3.18. Fix $m \geq n$, $J \triangleleft \mathcal{I}_m$ and $I \triangleleft \mathcal{I}_n$.

- (1) Define a partial order $J \leq I$ inductively on $m - n$ as follows: If $m = n$, then $J \leq I$ if $J = I$. If $m > n$, then $J \leq I$ if $K \leq I$ for some $K \triangleleft \mathcal{I}_{m-1}$ and there are some $x < y \leq 2u_{m-1}$ such that $J = \{x \hat{\ } \nu : \nu \in K\} \cup \{y \hat{\ } \nu : \nu \in K\}$.
- (2) Let $J \bowtie I$ be the set of all $\mu \in \omega^{<\omega}$ such that $I = \{\nu : \mu \hat{\ } \nu \in J\}$.
- (3) Given a class $\mathcal{A} \subseteq I \rightarrow 2^\omega$, let $J \otimes \mathcal{A}$ be the subclass of $J \rightarrow 2^\omega$ of all $\langle X_\nu^\mu : \nu \in I, \mu \in J \bowtie I \rangle$ such that for every $\mu \in J \bowtie I$, $\langle X_\nu^\mu : \nu \in I \rangle \in \mathcal{A}$.
- (4) Given a class $\mathcal{A} \subseteq I \rightarrow 2^\omega$, let $\mathcal{I}_n \odot \mathcal{A}$ be the subclass of $\mathcal{I}_n \rightarrow 2^\omega$ of all $\langle X_\nu : \nu \in \mathcal{I}_n \rangle$ such that $\langle X_\nu : \nu \in I \rangle \in \mathcal{A}$.
- (5) Let $\mathcal{A} \subseteq I \rightarrow 2^\omega$ and $\mathcal{B} \subseteq J \rightarrow 2^\omega$. We write $\mathcal{B} \leq \mathcal{A}$ if $J \leq I$ and $\mathcal{B} \subseteq J \otimes \mathcal{A}$.

One can easily prove that the relations $J \leq I$ and $\mathcal{B} \leq \mathcal{A}$ are partial orders.

Lemma 3.19 Suppose $J \leq I$. Then $J = \{\mu \hat{\ } \nu : \mu \in J \bowtie I, \nu \in I\}$.

Proof. Say $J \triangleleft \mathcal{I}_n$ and $I \triangleleft \mathcal{I}_n$ with $m \geq n$. We prove the lemma by induction over $m - n$. Suppose $m = n$. Then $I = J$, so $J \bowtie I = \{\epsilon\}$. In particular, $J = \{\epsilon \hat{\ } \nu : \nu \in I\}$. Suppose $m > n$. By definition of $J \leq I$, there is some $K \triangleleft \mathcal{I}_{m-1}$ and $x < y \leq 2u_{m-1}$ such that $K \leq I$ and $J = \{x \hat{\ } \nu : \nu \in K\} \cup \{y \hat{\ } \nu : \nu \in K\}$. By induction hypothesis, $K = \{\mu \hat{\ } \nu : \mu \in K \bowtie I, \nu \in I\}$. Thus $J \bowtie I = \{x \hat{\ } \mu : \mu \in K \bowtie I\} \cup \{y \hat{\ } \mu : \mu \in K \bowtie I\}$, and $J = \{x \hat{\ } \mu \hat{\ } \nu : \mu \in K \bowtie I, \nu \in I\} \cup \{y \hat{\ } \mu \hat{\ } \nu : \mu \in K \bowtie I, \nu \in I\} = \{\mu \hat{\ } \nu : \mu \in J \bowtie I, \nu \in I\}$. \square

3.4. \mathbb{Q} -forcing

We now define the partial order of \mathbb{Q} -conditions, which represent branches of \mathbb{P} -conditions.

Definition 3.20. A \mathbb{Q}_n -condition is a tuple $(\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I)$ where

- (1) $\sigma^i \subseteq A^i$ for each $i < 2$; I is an n -index
- (2) $\mathcal{A} \subseteq I \rightarrow 2^\omega$ is a largeness subclass of $\mathcal{L}_{\langle X_\nu : \nu \in I \rangle}$
- (3) $X_\nu \in \mathcal{M}$ for each $\nu \in I$ and \mathcal{A} is Π_2^0

We let $\mathbb{Q} = \bigcup_n \mathbb{Q}_n$.

Definition 3.21. The partial order on \mathbb{Q} is defined by

$$(\tau^0, \tau^1, Y_\mu, \mathcal{B} : \mu \in J) \leq (\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I)$$

if $J \leq I$, for every $\mu \in J$ and $\nu \in I$ such that ν is a suffix of μ , $Y_\mu \subseteq X_\nu$, $\mathcal{B} \leq \mathcal{A}$, and for every $i < 2$, $\sigma^i \preceq \tau^i$ and $\tau^i - \sigma^i \subseteq \bigcup_{\nu \in I} X_\nu$.

Lemma 3.22 Let $c = (\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I) \in \mathbb{Q}_n$ and $d = (\tau^0, \tau^1, Y_\mu, \mathcal{B} : \mu \in J) \in \mathbb{Q}_m$ with $m \geq n$ be such that $d \leq c$. Then for every $i < 2$, $(\tau^i, \bigcup_{\mu \in J} Y_\mu)$ Mathias extends $(\sigma^i, \bigcup_{\nu \in I} X_\nu)$.

Proof. Since $J \leq I$, then by Lemma 3.19, $J = \{\rho \hat{\ } \nu : \rho \in J \bowtie I, \nu \in I\}$. It follows that for every $\mu \in J$, there is some $\nu \in I$ such that ν is a suffix of μ , and by definition of $d \leq c$, $Y_\mu \subseteq X_\nu$. Therefore $\bigcup_{\mu \in J} Y_\mu \subseteq \bigcup_{\nu \in I} X_\nu$. Since $\tau^i \succeq \sigma^i$ and $\tau^i - \sigma^i \subseteq \bigcup_{\nu \in I} X_\nu$, then $(\tau^i, \bigcup_{\mu \in J} Y_\mu)$ Mathias extends $(\sigma^i, \bigcup_{\nu \in I} X_\nu)$. \square

3.5. Forcing relation

Definition 3.23. Let (σ, X) be a Mathias condition and $\Phi_e(G, x)$ be a Δ_0 formula with an integer variable x .

- (1) $(\sigma, X) \Vdash (\exists x)\Phi_e(G, x)$ if there is some $x \in \omega$ such that $\Phi_e(\sigma, x)$ holds.
- (2) $(\sigma, X) \Vdash (\forall x)\neg\Phi_e(G, x)$ if for every $x \in \omega$ and $\rho \subseteq X$, $\Phi_e(\sigma, x)$ does not hold.

Definition 3.24. Given some n -index I , let ζ_I be the function which takes as a paramter an index e of a Δ_0 formula $\Phi_e(G, x, y)$, a finite set $\sigma \in 2^{<\omega}$ and some integer $x \in \omega$, and returns a code for the Σ_1^0 class

$$\mathcal{U}_{\zeta_I(e, \sigma, x)}^I = \{ \langle X_\nu : \nu \in I \rangle : (\sigma, \bigcup_{\nu \in I} X_\nu) \Vdash (\forall y)\Phi_e(G, x, y) \}$$

Definition 3.25. Let $c = (\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I)$ be a \mathbb{Q}_n -condition, $\Phi_e(G, x, y)$ be a Δ_0 formula with free integer variables x and y , and let $i < 2$.

- (1) $c \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ if there is some $x \in \omega$ such that $(\sigma^i, \bigcup_{\nu \in I} X_\nu) \Vdash (\forall y)\Phi_e(G, x, y)$
- (2) $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$ if for every $x \in \omega$, every $\rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu$, $\mathcal{A} \subseteq \mathcal{U}_{\zeta_I(e, \sigma^i \cup \rho, x)}^I$

Lemma 3.26 Let c, d be two \mathbb{Q} -conditions such that $d \leq c$, and $\Phi_e(G, x, y)$ be a Δ_0 formula.

- (1) If $c \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ then so does d .
- (2) If $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$ then so does d .

Proof. Say $c = (\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I) \in \mathbb{Q}_n$ and $d = (\tau^0, \tau^1, Y_\mu, \mathcal{B} : \mu \in J) \in \mathbb{Q}_m$ with $m \geq n$.

(1) Suppose $c \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$. Then there some $x \in \omega$ such that $(\sigma^i, \bigcup_{\nu \in I} X_\nu) \Vdash (\forall y)\Phi_e(G, x, y)$. By Lemma 3.22, $(\tau^i, \bigcup_{\mu \in J} Y_\mu)$ Mathias extends $(\sigma^i, \bigcup_{\nu \in I} X_\nu)$, so $(\tau^i, \bigcup_{\mu \in J} Y_\mu) \Vdash (\forall y)\Phi_e(G, x, y)$. Folding the definition, $d \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$.

(2) Fix some $x \in \omega$ and some $\rho \subseteq A^i \cap (\bigcup_{\mu \in J} Y_\mu)$. Since $d \leq c$, then there is some $\rho_0 \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu$ such that $\tau^i = \sigma^i \hat{\ } \rho_0$. Moreover, by Lemma 3.22, $(\tau^i, \bigcup_{\mu \in J} Y_\mu)$ Mathias extends $(\sigma^i, \bigcup_{\nu \in I} X_\nu)$, so $\bigcup_{\mu \in J} Y_\mu \subseteq \bigcup_{\nu \in I} X_\nu$. Therefore $\rho_0 \cup \rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu$. By applying the definition of $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$ to x and $\rho_0 \cup \rho$, $\mathcal{A} \subseteq \mathcal{U}_{\zeta_I(e, \sigma^i \cup \rho_0 \cup \rho, x)}^I = \mathcal{U}_{\zeta_I(e, \tau^i \cup \rho, x)}^I$.

We claim that $\mathcal{B} \subseteq \mathcal{U}_{\zeta_J(e, \tau^i \cup \rho, x)}^J$. Since $\mathcal{B} \leq \mathcal{A}$, $\mathcal{B} \subseteq J \otimes \mathcal{A}$. Fix some $\langle Z_\nu^\mu : \nu \in I, \mu \in J \bowtie I \rangle$ such that for every $\mu \in J \bowtie I$, $\langle Z_\nu^\mu : \nu \in I \rangle \in \mathcal{A}$. Since $\mathcal{A} \subseteq \mathcal{U}_{\zeta_I(e, \tau^i \cup \rho, x)}^I$, for every $\mu \in J \bowtie I$, $(\tau^i, \bigcup_{\nu \in I} Z_\nu^\mu) \Vdash (\forall y)\Phi_e(G, x, y)$. Therefore $(\tau^i, \bigcup_{\nu \in I, \mu \in J \bowtie I} Z_\nu^\mu) \Vdash (\forall y)\Phi_e(G, x, y)$. Thus $\langle Z_\nu^\mu : \nu \in I, \mu \in J \bowtie I \rangle \in \mathcal{U}_{\zeta_J(e, \tau^i \cup \rho, x)}^J$. So $\mathcal{B} \subseteq \mathcal{U}_{\zeta_J(e, \tau^i \cup \rho, x)}^J$. It follows that $d \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$. \square

3.6. \mathbb{P} -forcing

Definition 3.27. A \mathbb{P}_n -condition is a tuple $(\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$ where

- (1) $\sigma_I^i \subseteq A^i$ for each $i < 2$ and $I \triangleleft \mathcal{I}_n$

- (2) $\mathcal{U}_C^{\mathcal{I}_n} \subseteq \mathcal{I}_n \rightarrow 2^\omega$ is a largeness subclass of $\mathcal{L}_{\langle X_\nu : \nu \in \mathcal{I}_n \rangle}$
(3) $X_\nu \in \mathcal{M}$ for each $\nu \in \mathcal{I}_n$; C is Δ_2^0

A \mathbb{P}_n -condition $c = (\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$ represents u_n many parallel \mathbb{Q}_n -conditions defined for each $I \triangleleft \mathcal{I}_n$ by

$$c^{[I]} = (\sigma_I^0, \sigma_I^1, X_\nu, \pi_I(\mathcal{U}_C) : \nu \in I)$$

We let $\mathbb{P} = \bigcup_n \mathbb{P}_n$.

Definition 3.28. The partial order on \mathbb{P} is defined by

$$(\tau_J^0, \tau_J^1, Y_\mu, D : J \triangleleft \mathcal{I}_m, \mu \in \mathcal{I}_m) \leq (\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$$

if $m \geq n$, and for every $J \triangleleft \mathcal{I}_m$ and $I \triangleleft \mathcal{I}_n$ such that $J \leq I$

$$(\tau_J^0, \tau_J^1, Y_\mu, \pi_J(\mathcal{U}_D) : \mu \in J) \leq (\sigma_I^0, \sigma_I^1, X_\nu, \pi_I(\mathcal{U}_C) : \nu \in I)$$

Lemma 3.29 Fix a \mathbb{P}_n -condition c and some $I \triangleleft \mathcal{I}_n$. For every \mathbb{Q}_n -condition $d \leq c^{[I]}$, then there is a \mathbb{P}_n -condition $e \leq c$ such that $e^{[I]} = d$.

Proof. Say $c = (\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$ and $d = (\tau_I^0, \tau_I^1, Y_\nu, \mathcal{A} : \nu \in I)$. By Lemma 3.10, there is some Δ_2^0 set $D \supseteq C$ such that $\mathcal{U}_D^{\mathcal{I}_n} \subseteq \mathcal{U}_C^{\mathcal{I}_n}$ is a largeness class and $\pi_I(\mathcal{U}_D^{\mathcal{I}_n}) = \mathcal{A}$. For every $J \triangleleft \mathcal{I}_n$ with $J \neq I$, let $\tau_J^0 = \sigma_J^0$ and $\tau_J^1 = \sigma_J^1$. For $\nu \in \mathcal{I}_n - I$, let $Y_\nu = X_\nu$. The \mathbb{P}_n -condition $e = (\tau_J^0, \tau_J^1, Y_\nu, D : J \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$ is an extension of c such that $e^{[I]} = d$. \square

3.7. Validity

As explained in Section 2, the forcing relation for Π_2^0 formulas relies on the 1-genericity of the filter for the properties to actually hold. We define the notion of validity so that the forced Π_2^0 formulas will be satisfied on the valid sides.

Definition 3.30. A \mathbb{Q}_n -condition $(\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I)$ is *i-valid* for $i < 2$ if $\langle X_\nu \cap A^i : \nu \in I \rangle \in \mathcal{A}$.

The following lemma ensures that whenever a Π_2^0 formula is forced on a valid side, then seeing the formula as a collection of Σ_1^0 formulas, one can satisfy each of them independently.

Lemma 3.31 Let c be an *i-valid* \mathbb{Q}_n -condition, and let $\Phi_e(G, x, y)$ be a Δ_1^0 formula. If $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$ then for every $x \in \omega$ there is some $d = (\tau^0, \tau^1, Y_\nu, \mathcal{A} : \nu \in I) \in \mathbb{Q}_n$ extending c such that $(\tau^i, \bigcup_{\nu \in I} Y_\nu) \Vdash (\exists y)\neg\Phi_e(G, x, y)$.

Proof. Say $c = (\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I)$ and fix $x \in \omega$. Since $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$, then $\mathcal{A} \subseteq \mathcal{U}_{\zeta_I(e, \sigma^i, x)}^I$. Since c is *i-valid*, then $\langle X_\nu \cap A^i : \nu \in I \rangle \in \mathcal{A}$, then $(\sigma^i, A^i \cap \bigcup_{\nu \in I} X_\nu) \not\Vdash (\forall y)\Phi_e(G, x, y)$. Therefore there is some $\rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu$ and some $y \in \omega$ such that $\neg\Phi_e(\sigma^i \cup \rho, x, y)$ holds. Let $\tau^i = \sigma^i \cup \rho$ and $\tau^{1-i} = \sigma^{1-i}$. For every $\nu \in I$, let $Y_\nu = X_\nu - \{0, \dots, \max \rho\}$. By Lemma 3.5, $\mathcal{A} \subseteq \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$. The tuple $d = (\tau^0, \tau^1, Y_\nu, \mathcal{A} : \nu \in I)$ is a \mathbb{Q}_n -condition extending c . Moreover $(\tau^i, \bigcup_{\nu \in I} Y_\nu) \Vdash (\exists y)\neg\Phi_e(G, x, y)$. \square

The following two lemmas state that every \mathbb{P} -filter induces as tree of valid sides.

Lemma 3.32 For every \mathbb{P}_n -condition c , there is some $I \triangleleft \mathcal{I}_n$ and some $i < 2$ such that $c^{[I]}$ is *i-valid*.

Proof. Say $c = (\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$. Since $A^0 \cup A^1 = \omega$ and by Lemma 3.6, $\mathcal{L}(\mathcal{U}_C^{\mathcal{I}_n})$ is a largeness class, then there is some $\langle i_\nu < 2 : \nu \in \mathcal{I}_n \rangle$ such that $\langle A^{i_\nu} : \nu \in \mathcal{I}_n \rangle \in \mathcal{L}(\mathcal{U}_C^{\mathcal{I}_n})$. Thus $\mathcal{U}_C^{\mathcal{I}_n} \cap \mathcal{L}_{\langle X_\nu : \nu \in \mathcal{I}_n \rangle} \cap \mathcal{L}_{\langle A^{i_\nu} : \nu \in \mathcal{I}_n \rangle}$ is a largeness class, so by Lemma 3.7, $\mathcal{U}_C^{\mathcal{I}_n} \cap \mathcal{L}_{\langle X_\nu \cap A^{i_\nu} : \nu \in \mathcal{I}_n \rangle}$ is a largeness class.

Let $B_0 = \{\nu \in \mathcal{I}_n : i_\nu = 0\}$ and $B_1 = \{\nu \in \mathcal{I}_n : i_\nu = 1\}$. Since $B_0 \cup B_1 = \mathcal{I}_n$, by Lemma 3.17, there is some $I \triangleleft \mathcal{I}_n$ and some $i < 2$ such that $I \subseteq B_i$. Since $\mathcal{U}_C^{\mathcal{I}_n} \cap \mathcal{L}_{\langle X_\nu \cap A^{i_\nu} : \nu \in \mathcal{I}_n \rangle}$ is a largeness class, then $\langle X_\nu \cap A^{i_\nu} : \nu \in I \rangle \in \pi_I(\mathcal{U}_C^{\mathcal{I}_n} \cap \mathcal{L}_{\langle X_\nu \cap A^{i_\nu} : \nu \in \mathcal{I}_n \rangle})$. Moreover $\pi_I(\mathcal{U}_C^{\mathcal{I}_n} \cap \mathcal{L}_{\langle X_\nu \cap A^{i_\nu} : \nu \in \mathcal{I}_n \rangle}) \subseteq$

$\pi_I(\mathcal{U}_C^{\mathcal{I}_n})$, then $\langle X_\nu \cap A^{i\nu} : \nu \in I \rangle \in \pi_I(\mathcal{U}_C^{\mathcal{I}_n})$. As $I \subseteq B_i$, $\langle X_\nu \cap A^i : \nu \in I \rangle = \langle X_\nu \cap A^{i\nu} : \nu \in I \rangle \in \pi_I(\mathcal{U}_C^{\mathcal{I}_n})$. Thus the \mathbb{Q}_n -condition $c^{[I]}$ is i -valid. \square

Lemma 3.33 Let $d, c \in \mathbb{Q}$ be such that $d \leq c$. If d is i -valid, then so is c .

Proof. Say $c = (\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I) \in \mathbb{Q}_n$ and $d = (\tau^0, \tau^1, Y_\mu, \mathcal{B} : \mu \in J) \in \mathbb{Q}_m$ with $m \geq n$. Since d is i -valid, $\langle Y_\mu \cap A^i : \mu \in J \rangle \in \mathcal{B}$. Since $d \leq c$, then $J \leq I$ and $\mathcal{B} \leq \mathcal{A}$. By definition of $\mathcal{B} \leq \mathcal{A}$, $\mathcal{B} \subseteq J \otimes \mathcal{A}$, thus letting $\rho \in J \otimes I$, and $Z_\nu = Y_{\rho \sim \nu}$, $\langle Z_\nu \cap A^i : \nu \in I \rangle \in \mathcal{A}$. By upward-closure of \mathcal{A} , $\langle X_\nu \cap A^i : \nu \in I \rangle \in \mathcal{A}$. Thus c is i -valid. \square

The following lemma states that the generic sets corresponding to valid sides are infinite.

Lemma 3.34 For every i -valid \mathbb{Q}_n -condition $c = (\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in I)$, there is a \mathbb{Q}_n -condition $d = (\tau^0, \tau^1, Y_\nu, \mathcal{A} : \nu \in I) \leq c$ such that $\#\tau^i > \#\sigma^i$.

Proof. By definition of i -validity of c , $\langle X_\nu \cap A^i : \nu \in I \rangle \in \mathcal{A} \subseteq \mathcal{L}_{\langle X_\nu : \nu \in I \rangle}$. So in particular, $X_\nu \cap A^i$ is infinite. Pick any $x \in \bigcup_{\nu \in I} X_\nu \cap A^i$, and let $Y_\nu = X_\nu - \{0, \dots, x\}$. By Lemma 3.5, $\mathcal{A} \subseteq \mathcal{L}_{\langle Y_\nu : \nu \in I \rangle}$. Then $d = (\sigma^i \cup \{x\}, \sigma^{1-i}, Y_\nu, \mathcal{A} : \nu \in I)$ is the desired extension. \square

3.8. Forcing question

As explained in Section 2, a \mathbb{P} -condition representing multiple parallel \mathbb{Q} -condition, only one of which being valid on one side, we need to force the requirements on each side of each branch. In the following forcing question, the finite set H is intended to be the set of all branches which have not been forced yet.

Definition 3.35. Let $c = (\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n) \in \mathbb{P}_n$, let $H \subseteq \{I \triangleleft \mathcal{I}_n\}$, let $\Phi_e(G, x, y)$ be a Δ_0 formula with free variables x and y and let $i < 2$. Define the relation $c \text{?}\vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y)$ to hold if

$$\mathcal{U}_C^{\mathcal{I}_n} \cap \left\{ \mathcal{I}_n \odot \mathcal{U}_{\zeta_I(e, \sigma_I^i \cup \rho, x)}^I : I \in H, x \in \omega, \rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu \right\}$$

is not a largeness class

Lemma 3.36 Let $c \in \mathbb{P}_n$, let $H \subseteq \{I \triangleleft \mathcal{I}_n\}$, let $\Phi_e(G, x, y)$ be a Δ_0 formula with free variables x and y , and let $i < 2$. The relation $c \text{?}\vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y)$ is $\Sigma_1^{0, \emptyset'}$.

Proof. By Lemma 3.2, $c \text{?}\vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y)$ holds if there is a finite set $F \subseteq C$, and some $t \in \omega$ such that the following class

$$\mathcal{U}_F^{\mathcal{I}_n} \cap \left\{ \mathcal{I}_n \odot \mathcal{U}_{\zeta_I(e, \sigma_I^i \cup \rho, x)}^I : I \in H, x < t, \rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu \upharpoonright t \right\}$$

is not a largeness class. Note that this class is $\Sigma_1^{0, Z}$ for some $Z \in \mathcal{M}$. By Lemma 3.3, not being a largeness class for a $\Sigma_1^{0, Z}$ class is $\Sigma_2^{0, Z}$, hence $\Sigma_1^0(\emptyset')$ whenever Z is low. Thus, the whole formula is $\Sigma_1^0(A^i \oplus C \oplus \emptyset')$. Since A^i and C are Δ_2^0 , the formula is $\Sigma_1^0(\emptyset')$. \square

The following lemma states that in the Σ_2^0 outcome, one can find an extension forcing the Σ_2^0 formula on one branch of H , which is the set of branches not having been satisfied yet.

Lemma 3.37 Let $c \in \mathbb{P}_n$, let $H \subseteq \{I \triangleleft \mathcal{I}_n\}$, let $\Phi_e(G, x, y)$ be a Δ_0 formula with free variables x and y and let $i < 2$. Suppose

$$c \text{?}\vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y)$$

then there is some $d \in \mathbb{P}_n$ with $d \leq c$ and some $I \in H$ such that $d^{[I]} \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$.

Proof. Say $c = (\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$. Since $c \text{ ? } \vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y)$, then by Lemma 3.2, there is a finite set $F \subseteq C$, and some $t \in \omega$ such that the following class

$$\mathcal{U}_F^{\mathcal{I}_n} \cap \{\mathcal{I}_n \odot \mathcal{U}_{\zeta_I(e, \sigma_I^i \cup \rho, x)}^I : I \in H, x < t, \rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu \upharpoonright t\}$$

is not a largeness class. Since the class is $\Sigma_1^{0, Y}$ for some $Y \in \mathcal{M}$ and since \mathcal{M} is a Scott set, there is a cover $Z_0 \cup \dots \cup Z_{k-1} = \omega$ in \mathcal{M} such that for every $j < k$, $Z_j \notin \mathcal{U}_F^{\mathcal{I}_n} \cap \{\mathcal{I}_n \odot \mathcal{U}_{\zeta_I(e, \sigma_I^i \cup \rho, x)}^I : I \in H, x < n, \rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu \upharpoonright n\}$.

By Lemma 3.6, $\mathcal{L}(\mathcal{U}_C^{\mathcal{I}_n})$ is a largeness class, then there is some $\langle j_\nu : \nu \in \mathcal{I}_n \rangle$ such that $\langle Z_{j_\nu} : \nu \in \mathcal{I}_n \rangle \in \mathcal{L}(\mathcal{U}_C^{\mathcal{I}_n})$. Thus $\mathcal{U}_C^{\mathcal{I}_n} \cap \mathcal{L}_{\langle X_\nu : \nu \in \mathcal{I}_n \rangle} \cap \mathcal{L}_{\langle Z_{j_\nu} : \nu \in \mathcal{I}_n \rangle}$ is a largeness class, so by Lemma 3.7, the class $\mathcal{U}_C^{\mathcal{I}_n} \cap \mathcal{L}_{\langle X_\nu \cap Z_{j_\nu} : \nu \in \mathcal{I}_n \rangle}$ is a largeness class. In particular $\langle X_\nu \cap Z_{j_\nu} : \nu \in \mathcal{I}_n \rangle \in \mathcal{U}_C^{\mathcal{I}_n}$, so there is some $I \in H$, some $x < t$ and some $\rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu \upharpoonright t$ such that

$$\langle X_\nu \cap Z_{j_\nu} : \nu \in \mathcal{I}_n \rangle \notin \mathcal{I}_n \odot \mathcal{U}_{\zeta_I(e, \sigma_I^i \cup \rho, x)}^I$$

Let $D \supseteq C$ be such that $\mathcal{U}_D^{\mathcal{I}_n} = \mathcal{U}_C^{\mathcal{I}_n} \cap \mathcal{L}_{\langle X_\nu \cap Z_{j_\nu} : \nu \in \mathcal{I}_n \rangle}$. For every $\nu \in \mathcal{I}_n$, let $Y_\nu : (X_\nu \cap Z_{j_\nu}) - \{0, \dots, t\}$. In particular, $\mathcal{U}_D^{\mathcal{I}_n} \subseteq \mathcal{L}_{\langle Y_\nu : \nu \in \mathcal{I}_n \rangle}$. Let $\tau_I^i = \sigma_I^i \cup \rho$, and $\tau_I^{1-i} = \sigma_I^{1-i}$. For every $J \triangleleft \mathcal{I}_n$ with $J \neq I$, let $\tau_J^0 = \sigma_J^0$ and $\tau_J^1 = \sigma_J^1$. The \mathbb{P}_n -condition $d = (\tau_J^0, \tau_J^1, Y_\nu, D : J \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$ is an extension of c such that $d \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$ with $I \in H$. \square

The following lemma states that whenever sufficiently many formulas have satisfied the Π_2^0 outcome, then one can find an extension with more branches, such that any branch refining a branch in H will force at least two of the Π_2^0 formulas. Letting H be the set of branches for which the requirement has not been forced yet, one obtain an extension on which the requirement is forced on all branches simultaneously.

Lemma 3.38 Let $c \in \mathbb{P}_n$, let $H \subseteq \{I \triangleleft \mathcal{I}_n\}$, let $\Phi_{e_0}(G, x, y), \dots, \Phi_{e_{2u_n}}(G, x, y)$ be $2u_n + 1$ many Δ_0 formulas with free variables x and y and let $i < 2$. Suppose that for every $j \leq 2u_n$,

$$c \text{ ? } \Vdash_H^i (\exists x)(\forall y)\Phi_{e_j}(G, x, y)$$

Then there is some $d \in \mathbb{P}_{n+1}$ with $d \leq c$ such that for every $I \in H$ and $J \triangleleft \mathcal{I}_{n+1}$ such that $J \leq I$, there are some $a < b \leq 2u_n$ such that

$$d \Vdash^{[J]} (\forall x)(\exists y) \neg \Phi_{e_a}(G, x, y) \quad \text{and} \quad d \Vdash^{[J]} (\forall x)(\exists y) \neg \Phi_{e_b}(G, x, y)$$

Proof. Say $c = (\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n)$. For every $j \leq 2u_n$, the class

$$\mathcal{A}_j = \mathcal{U}_C^{\mathcal{I}_n} \cap \{\mathcal{I}_n \odot \mathcal{U}_{\zeta_I(e_j, \sigma_I^i \cup \rho, x)}^I : I \in H, x \in \omega, \rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu\}$$

is a largeness class. Let $D \subseteq \omega$ be a Δ_2^0 set such that $\mathcal{U}_D^{\mathcal{I}_{n+1}}$ is the class of all $\langle Z_{j \frown \nu} : j \leq 2u_n, \nu \in \mathcal{I}_n \rangle$ such that for every $j \leq 2u_n$, $\langle Z_{j \frown \nu} : \nu \in \mathcal{I}_n \rangle \in \mathcal{A}_j$. In particular, $\mathcal{U}_D^{\mathcal{I}_{n+1}}$ is a largeness class. For every $j \frown \nu \in \mathcal{I}_{n+1}$, let $Y_{j \frown \nu} = X_\nu$. For every $J \triangleleft \mathcal{I}_{n+1}$, let $\tau_J^0 = \sigma_J^0$ and $\tau_J^1 = \sigma_J^1$, where $I \triangleleft \mathcal{I}_n$ is the unique n -index such that $J \leq I$.

Claim 1: $\mathcal{U}_D^{\mathcal{I}_{n+1}} \subseteq \mathcal{L}_{\langle Y_\mu : \mu \in \mathcal{I}_{n+1} \rangle}$. Let $\langle Z_{j \frown \nu} : j \leq 2u_n, \nu \in \mathcal{I}_n \rangle \in \mathcal{U}_D^{\mathcal{I}_{n+1}}$. For every $j \leq 2u_n$, $\langle Z_{j \frown \nu} : \nu \in \mathcal{I}_n \rangle \in \mathcal{A}_j$. Since $\mathcal{A}_j \subseteq \mathcal{L}_{\langle X_\nu : \nu \in \mathcal{I}_n \rangle}$, then $|Z_{j \frown \nu} \cap X_\nu| = \infty$. Since $X_\nu = Y_{j \frown \nu}$, then $|Z_{j \frown \nu} \cap Y_{j \frown \nu}| = \infty$, so $\langle Z_{j \frown \nu} : j \leq 2u_n, \nu \in \mathcal{I}_n \rangle \in \mathcal{L}_{\langle Y_\mu : \mu \in \mathcal{I}_{n+1} \rangle}$. This proves Claim 1.

Claim 2: $\mathcal{U}_D^{\mathcal{I}_{n+1}} \leq \mathcal{U}_C^{\mathcal{I}_n}$. We need to prove that $\mathcal{U}_D^{\mathcal{I}_{n+1}} \subseteq \mathcal{I}_{n+1} \otimes \mathcal{U}_C^{\mathcal{I}_n}$. Fix $\langle Z_{j \frown \nu} : j \leq 2u_n, \nu \in \mathcal{I}_n \rangle \in \mathcal{U}_D^{\mathcal{I}_{n+1}}$. Then for every $j \leq 2u_n$, $\langle Z_{j \frown \nu} : \nu \in \mathcal{I}_n \rangle \in \mathcal{A}_j \subseteq \mathcal{U}_C^{\mathcal{I}_n}$. Thus $\mathcal{U}_D^{\mathcal{I}_{n+1}} \subseteq \mathcal{I}_{n+1} \otimes \mathcal{U}_C^{\mathcal{I}_n}$. This proves Claim 2.

Let $d = (\tau_J^0, \tau_J^1, Y_\mu, D : J \triangleleft \mathcal{I}_{n+1}, \mu \in \mathcal{I}_{n+1})$. In particular d is a \mathbb{P}_{n+1} -condition extending c . Fix $I \in H$ and $J \triangleleft \mathcal{I}_{n+1}$ such that $J \leq I$. In particular, there are some $a < b \leq 2u_n$ such that $J = \{a \frown \nu : \nu \in I\} \cup \{b \frown \nu : \nu \in I\}$.

Claim 3: $d^{[J]} \Vdash^i (\forall x)(\exists y)\neg\Phi_{e_a}(G, x, y)$ and $d^{[J]} \Vdash^i (\forall x)(\exists y)\neg\Phi_{e_b}(G, x, y)$. We prove that $d^{[J]} \Vdash^i (\forall x)(\exists y)\neg\Phi_{e_a}(G, x, y)$. The other case is symmetric. For every $x \in \omega$ and $\rho \subseteq A^i \cap \bigcup_{\mu \in J} Y_\mu$, in particular $\rho \subseteq A^i \cap \bigcup_{\nu \in I} X_\nu$. Fix $\langle Z_\mu : \mu \in J \rangle \in \pi_J(\mathcal{U}_D^{\mathcal{I}_{n+1}})$. In particular $\langle Z_{a^\nu} : \nu \in I \rangle \in \mathcal{A}_a \subseteq \mathcal{U}_{\zeta_I(e_a, \sigma_I^i \cup \rho, x)}^I$. So $(\sigma_I^i \cup \rho, \bigcup_{\nu \in I} Z_{a^\nu}) \Vdash (\forall y)\Phi_{e_a}(G, x, y)$. As $\sigma_I^i = \tau_J^i$ and $\bigcup_{\nu \in I} Z_{a^\nu} \subseteq \bigcup_{\mu \in J} Z_\mu$, then $(\tau_J^i \cup \rho, \bigcup_{\mu \in J} Z_\mu) \Vdash (\forall y)\Phi_{e_a}(G, x, y)$. So $\langle Z_\mu : \mu \in J \rangle \in \mathcal{U}_{\zeta_J(e_a, \tau_J^i \cup \rho, x)}^J$. Thus for every $x \in \omega$ and $\rho \subseteq A^i \cap \bigcup_{\mu \in J} Y_\mu$, $\pi_J(\mathcal{U}_D^{\mathcal{I}_{n+1}}) \subseteq \mathcal{U}_{\zeta_J(e_a, \tau_J^i \cup \rho, x)}^J$. This is the definition of $d^{[J]} \Vdash^i (\forall x)(\exists y)\neg\Phi_{e_a}(G, x, y)$. This proves Claim 3 and Lemma 3.38. \square

3.9. Requirements

We now define the requirements specific to our purpose, namely, obtaining a set whose jump does not compute a $\{0, 1\}$ -valued completion of the partial function $n \mapsto \Phi_n^{\theta'}$.

Definition 3.39. Fix a Δ_0 formula $\Phi_e(G, x, y, p)$ with free integer variables x and y , and free valuation variable p .

- (1) Let $c \in \mathbb{Q}_n$ and $i < 2$. We say that c forces the e -th requirement on side i if $c \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y, p)$ for some θ' -correct valuation p , or $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y, p_0)$ and $c \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y, p_1)$ for two incompatible valuations.
- (2) Let $c \in \mathbb{P}_n$ and $i < 2$. We say that c forces the e -th requirement on side i if $c^{[I]}$ forces the e -th requirement on side i for every $I \triangleleft \mathcal{I}_n$.

Given a condition $c \in \mathbb{P}_n$, $e \in \omega$ and $i < 2$, let $H(c, e, i)$ be the set of $I \triangleleft \mathcal{I}_n$ such that c does not force the e -th requirement on the i -th side.

Lemma 3.40 For every $c \in \mathbb{P}_n$, $i < 2$ and $e \in \omega$ such that $H(c, e, i) \neq \emptyset$, there is \mathbb{P} -condition $d \leq c$ such that $|H(d, e, i)| < |H(c, e, i)|$.

Proof. Let $H = H(c, e, i)$ and W be the set of all valuations p such that $c \Vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y, p)$. By Lemma 3.3, the set W is θ' -c.e., so by Lemma 3.12, we have two cases.

Case 1: $p \in W$ for some θ' -correct valuation p . By definition of W , $c \Vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y, p)$. By Lemma 3.37, there is a \mathbb{P}_n -condition $d \leq c$ such that $|H(d, e, i)| < |H(c, e, i)|$.

Case 2: $p_0, \dots, p_{2u_n} \notin W$ for $2u_n + 1$ pairwise incompatible valuations. So

$$c \not\Vdash_H^i (\exists x)(\forall y)\Phi_e(G, x, y, p_j)$$

for every $j \leq 2u_n$. By Lemma 3.38, there is a \mathbb{P}_{n+1} -condition $d \leq c$ such that $d^{[J]}$ forces the e -th requirement on side i for every $J \triangleleft \mathcal{I}_n$ such that $J \leq I$ for some $I \in H = H(c, e, i)$. Therefore $|H(d, e, i)| = 0 < |H(c, e, i)|$. \square

Lemma 3.41 For every $c \in \mathbb{P}$ and $e \in \omega$, there is \mathbb{P} -condition $d \leq c$ forcing the e -th requirement on both sides.

Proof. Apply iteratively Lemma 3.40 to obtain a condition d_0 such that $H(d_0, e, 0) = \emptyset$. Then apply again iteratively Lemma 3.40 below d_0 to obtain an extension d_1 such that $H(d_1, e, 1) = \emptyset$. The condition d_1 is the desired extension. \square

3.10. Construction

As explained, a \mathbb{P} -condition represents multiple parallel \mathbb{Q} -conditions. By Lemma 3.32, every \mathbb{P} -condition admits a branch with a valid side. Moreover, by Lemma 3.33, the valid sides of \mathbb{Q} -conditions are upward-closed under the extension relation. This motivates the following definition.

Definition 3.42. A path through a \mathbb{P} -filter \mathcal{F} is a pair $\langle P, i \rangle$ where $i < 2$ and for every $n \in \omega$, $P(n) \triangleleft \mathcal{I}_n$ is such that $P(n+1) \leq P(n)$ and for every $c \in \mathcal{F} \cap \mathbb{P}_n$, $c^{[P(n)]}$ is i -valid.

By Lemma 3.32 and Lemma 3.33, every \mathbb{P} -filter admits a path. We then let

$$\mathcal{F}(P, i) = \bigcup \{ \sigma_{P(n)}^i : (\sigma_I^0, \sigma_I^1, X_\nu, C : I \triangleleft \mathcal{I}_n, \nu \in \mathcal{I}_n) \in \mathcal{F} \}$$

We can prove that the forced formulas hold along any path.

Lemma 3.43 Let \mathcal{F} be a sufficiently generic \mathbb{P} -filter, and let $\langle P, i \rangle$ be a path through \mathcal{F} and let $G^i = \mathcal{F}(P, i)$. Let $\Phi_e(G, x, y)$ be a Δ_0 formula and $c \in \mathcal{F}$.

- (1) If $c^{[P(n)]} \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$, then $(\exists x)(\forall y)\Phi_e(G^i, x, y)$ holds.
- (2) If $c^{[P(n)]} \Vdash^i (\forall x)(\exists y)\neg\Phi_e(G, x, y)$, then $(\forall x)(\exists y)\neg\Phi_e(G^i, x, y)$ holds.

Proof. Say $c^{[P(n)]} = (\sigma^0, \sigma^1, X_\nu, \mathcal{A} : \nu \in P(n)) \in \mathbb{Q}_n$.

(1) By definition of $c^{[P(n)]} \Vdash^i (\exists x)(\forall y)\Phi_e(G, x, y)$, then there is some $x \in \omega$ such that $(\sigma^i, \bigcup_{\nu \in P(n)} X_\nu) \Vdash (\forall y)(\Phi_e(G^i, x, y))$. In particular, $\sigma^i \prec G^i$ and $G^i - \sigma^i \subseteq \bigcup_{\nu \in P(n)} X_\nu$, so for every $y \in \omega$, $\neg\Phi(G^i, x, y)$ holds.

(2) By Lemma 3.26, Lemma 3.31 and Lemma 3.29, for every $x \in \omega$, there is some $m \in \omega$ and $d \in \mathcal{F} \cap \mathbb{P}_m$ such that $d^{[P(m)]} = (\tau^0, \tau^1, Y_\mu, \mathcal{B} : \mu \in P(m))$ and $(\tau^i, \bigcup_{\mu \in P(m)} Y_\mu) \Vdash (\exists y)\neg\Phi_e(G, x, y)$. In particular, $\tau^i \prec G^i$, so $(\exists y)\Phi_e(G^i, x, y)$ holds. \square

We are now ready to prove Theorem 1.3.

Proof of Theorem 1.3. We prove the theorem for $Z = \emptyset$, as the whole argument relativizes. Fix a Δ_2^0 set A and let $A^0 = A$ and $A^1 = \bar{A}$. Let \mathcal{F} be a sufficiently generic \mathbb{P} -filter. Let $\langle P, i \rangle$ be a path through \mathcal{F} . Let $G^i = \mathcal{F}(P, i)$. By definition of a \mathbb{P} -condition, $G^i \subseteq A^i$. By Lemma 3.34 and Lemma 3.29, G^i is infinite.

By Lemma 3.41, for every $e \in \omega$, there is some $n \in \omega$ and some $c \in \mathcal{F} \cap \mathbb{P}_n$ such that c forces the e -th requirement both sides. In particular, $c^{[P(n)]}$ forces the e -th requirement on side i . By Lemma 3.43, the e -th requirement holds on G^i . By Lemma 3.13, the jump of G^i is not PA over \emptyset' . This completes the proof of Theorem 1.3. \square

REFERENCES

- [1] Peter A. Cholak, Damir D. Dzhafarov, Denis R. Hirschfeldt, and Ludovic Patey. Some results concerning the SRT_2^2 vs. COH problem, 2019.
- [2] Peter A. Cholak, Carl G. Jockusch, and Theodore A. Slaman. On the strength of Ramsey's theorem for pairs. *Journal of Symbolic Logic*, 66(01):1–55, 2001.
- [3] C. Chong, Steffen Lempp, and Yue Yang. On the role of the collection principle for Σ_2^0 -formulas in second-order reverse mathematics. *Proceedings of the American Mathematical Society*, 138(3):1093–1100, 2010.
- [4] Chitat Chong, Theodore Slaman, and Yue Yang. The metamathematics of stable Ramsey's theorem for pairs. *Journal of the American Mathematical Society*, 27(3):863–892, 2014.
- [5] François G. Dorais, Damir D. Dzhafarov, Jeffrey L. Hirst, Joseph R. Mileti, and Paul Shafer. On uniform relationships between combinatorial problems. *Trans. Amer. Math. Soc.*, 368(2):1321–1359, 2016.
- [6] Rod Downey, Denis R. Hirschfeldt, Steffen Lempp, and Reed Solomon. A Δ_2^0 set with no infinite low subset in either it or its complement. *Journal of Symbolic Logic*, 66(3):1371–1381, 2001.
- [7] Damir Dzhafarov and Ludovic Patey. Coh, srt22, and multiple functionals, 2019.
- [8] Damir D. Dzhafarov. Cohesive avoidance and strong reductions. *Proceedings of the American Mathematical Society*, 143(2):869–876, 2014.
- [9] Damir D. Dzhafarov. Strong reductions between combinatorial principles. *J. Symb. Log.*, 81(4):1405–1431, 2016.
- [10] Richard Elwes. *Math in 100 key breakthroughs*. Quercus, New York, 2013.
- [11] Denis R. Hirschfeldt. *Slicing the truth*, volume 28 of *Lecture Notes Series. Institute for Mathematical Sciences. National University of Singapore*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2015. On the computable and reverse mathematics of combinatorial principles, Edited and with a foreword by Chitat Chong, Qi Feng, Theodore A. Slaman, W. Hugh Woodin and Yue Yang.
- [12] Denis R. Hirschfeldt and Carl G. Jockusch. On notions of computability-theoretic reduction between Π_2^1 principles. *J. Math. Log.*, 16(1):1650002, 59, 2016.
- [13] Denis R. Hirschfeldt, Carl G. Jockusch, Bjørn Kjos-Hanssen, Steffen Lempp, and Theodore A. Slaman. The strength of some combinatorial principles related to Ramsey's theorem for pairs. *Computational Prospects of Infinity, Part II: Presented Talks, World Scientific Press, Singapore*, pages 143–161, 2008.
- [14] Carl G. Jockusch. Ramsey's theorem and recursion theory. *Journal of Symbolic Logic*, 37(2):268–280, 1972.
- [15] Carl G. Jockusch and Robert I. Soare. Π_1^0 classes and degrees of theories. *Transactions of the American Mathematical Society*, 173:33–56, 1972.

- [16] Carl G. Jockusch and Frank Stephan. A cohesive set which is not high. *Mathematical Logic Quarterly*, 39(1):515–530, 1993.
- [17] Lu Liu. RT₂² does not imply WKL₀. *Journal of Symbolic Logic*, 77(2):609–620, 2012.
- [18] Lu Liu. Cone avoiding closed sets. *Transactions of the American Mathematical Society*, 367(3):1609–1630, 2015.
- [19] Joseph Roy Mileti. *Partition theorems and computability theory*. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)—University of Illinois at Urbana-Champaign.
- [20] Benoit Monin and Ludovic Patey. Pigeons do not jump high. To appear. Available at <https://arxiv.org/abs/1803.09771>, 2018.
- [21] Ludovic Patey. The weakness of being cohesive, thin or free in reverse mathematics. *Israel J. Math.*, 216(2):905–955, 2016.
- [22] Ludovic Patey. Controlling iterated jumps of solutions to combinatorial problems. *Computability*, 6(1):47–78, 2017.
- [23] Ludovic Patey. Ramsey-like theorems and moduli of computation, 2019.
- [24] David Seetapun and Theodore A. Slaman. On the strength of Ramsey’s theorem. *Notre Dame Journal of Formal Logic*, 36(4):570–582, 1995.
- [25] Stephen G. Simpson. Partial realizations of Hilbert’s Program. *J. Symbolic Logic*, 53(2):349–363, 1988.
- [26] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Cambridge University Press, 2009.
- [27] Wei Wang. Cohesive sets and rainbows. *Annals of Pure and Applied Logic*, 165(2):389–408, 2014.