

CONTINUOUS HIGHER RANDOMNESS

LAURENT BIENVENU, NOAM GREENBERG, AND BENOIT MONIN

ABSTRACT. We investigate the role of continuous reductions and continuous relativisation in the context of higher randomness. We define a higher analogue of Turing reducibility and show that it interacts well with higher randomness, for example with respect to van-Lambalgen’s theorem and the Miller-Yu / Levin theorem. We study lowness for continuous relativization of randomness, and show the equivalence of the higher analogues of the different characterisations of lowness for Martin-Löf randomness. We also characterise computing higher K -trivial sets by higher random sequences. We give a separation between higher notions of randomness, in particular between higher weak-2-randomness and Π_1^1 -randomness. To do so we investigate classes of functions computable from Kleene’s O based on strong forms of the higher limit lemma.

1. INTRODUCTION

Algorithmic randomness uses the tools of computability theory to give a formal definition of the notion of a random infinite binary sequence, a sequence we would expect be the result of independent coin tosses. Many theorems of probability theory and analysis detail properties of real numbers which are shared by all elements of a set of measure 1. In other words a “typical” – or “random” real satisfies the property. For example, a monotone function is differentiable at almost every real. This fact though does not tell us what “typical reals” are; for every real x there is some monotone function which is not differentiable at x . Restricting ourselves to a computable viewpoint allows us to consider only countably many properties of measure 1. For example we can characterise the collection of reals x at which every computable monotone function is differentiable [BMN16].

Varying the computational strength of the tools involved we obtain in fact a hierarchy of randomness notions. Roughly, the stronger the tools we have the easier it is to detect irregular behaviour and so the harder it is to be considered random. Many of the resulting notions of randomness are robust. The best known notion, due to Martin-Löf [ML66], can be defined by using computably enumerable betting strategies, by the incompressibility of initial segments, and by specifying a natural class of effectively presented, effectively null G_δ sets. The resulting field studies these notions of randomness, investigates questions such as “what does it mean for one sequence to be more random than another?”, measures the computational strength of random oracles, looks at connections to effective analysis, and much more (see [Nie09, DH10]). A particularly deep area of investigation concerns notions opposite to randomness, such as K -triviality, and relates them to computational weakness.

Greenberg was supported by the Marsden Fund, by a Rutherford Discovery Fellowship from the Royal Society of New Zealand, and by a Turing research fellowship from the Templeton Foundation. Monin was partially supported by the Marsden Fund.

While they in some sense formalise the intuitive notion of effective computation (albeit disregarding questions of time and space resources), computability-related notions do not satisfy natural closure properties. For example, the variation function of a computable function of bounded variation need not be computable. As result, even though every function of bounded variation is the difference of two monotone functions, a real number x can be random in the sense that every computable monotone function is differentiable at x , but not in the sense that every computable function of bounded variation is differentiable at x . This is related to the fact that the halting problem is not computable. To overcome similar problems, Martin-Löf himself suggested that the “pattern detection tools” for defining randomness should be taken from a much larger collection. Such collections are given by the closely-related fields of effective descriptive set theory and so-called “higher computability” (see [Sac90]). The collection of Δ_1^1 (or *hyperarithmetical*) sets is the smallest one closed under taking the relativised halting problem and closing downward under Turing reducibility; alternatively, under taking infinite computable Boolean operations. Martin-Löf defined a real to be Δ_1^1 -random if it is an element of every Δ_1^1 set of measure 1. The closure properties of the hyperarithmetical sets result, for example, in the fact that a real x is Δ_1^1 random if and only if every Δ_1^1 monotone function is differentiable at x if and only if every Δ_1^1 function of bounded variation is differentiable at x .

Beyond the desirable closure properties, working with Δ_1^1 and Π_1^1 sets is particularly natural and appealing to computability theorists. This is because one can view these notions as analogues of the fundamental and familiar notions of “computable” and “computably enumerable”, interpreted over an enlarged domain of computation. The theory of admissible computability generalises computability to *admissible ordinals*. The smallest admissible ordinal is ω_1^{ck} , the least ordinal which is not the order-type of a computable well-ordering of the natural numbers. The corresponding domain of computation is $L_{\omega_1^{\text{ck}}}$, the smallest admissible set, which is the initial segment of the constructible universe of height ω_1^{ck} . A real is Δ_1^1 if and only if it is an element of $L_{\omega_1^{\text{ck}}}$. The Spector-Gandy theorem says that the Π_1^1 sets are those which are defined by an existential quantifier ranging over the collection of hyperarithmetical sets. Via coding of structures by reals this shows that the Π_1^1 sets are precisely those which are computably enumerable over the structure $L_{\omega_1^{\text{ck}}}$. Informally, these are the sets that can be enumerated effectively if the enumeration procedure takes ω_1^{ck} many steps. With this viewpoint in mind, many intuitive ideas from traditional “countable” computability (computability over ω), for example reduction and separation theorems (or the fixed-point theorem) extend to the higher setting with precisely the same proofs.

An important advance in the theory of “higher randomness” was made by Hjorth and Nies in [HN07]. They examined the higher analogue of Martin-Löf randomness and also isolated the new, stronger notion of Π_1^1 -randomness. They also looked at the higher analogues of the K -trivial sets. The theory was then further developed by Chong, Nies and Yu [CNY08] and by Chong and Yu [CY15]. One of the projects they are concerned with is the separation of higher notions of randomness. One of the results in this paper is the separation between Π_1^1 -randomness and the higher analogue of weak 2-randomness. We also consider higher K -triviality.

1.1. Randomness and continuity. A main theme of this paper is the centrality of continuous reductions to the theory of randomness. The insight that randomness

and traditional relative hyperarithmetic reducibility do not interact well goes back to Hjorth and Nies [HN07].

As a first motivating example we consider the fact that strong randomness notions are downward closed in the Turing degrees of ML-random sets. For example, Miller and Yu [MY08] showed that if X and Y are ML-random, Y computes X and Y is in addition \mathbf{d} -random (for some Turing degree \mathbf{d}) then X too is \mathbf{d} -random. Similarly, an ML-random set X is weakly 2-random if and only if it forms a minimal pair with \emptyset' [Nie09, Theorem 5.3.15], a property clearly downward closed in the Turing degrees. Another example is difference randomness, which is equivalent to being ML-random and not computing \emptyset' [FN11]. The argument of Miller and Yu's works for almost every randomness notion stronger than Martin-Löf's: suppose that Y computes X and that X is random; say $\Phi(Y) = X$ where Φ is some Turing functional. For a finite binary string σ let $\Phi^{-1}[\sigma]$ be the collection of oracles Z such that $\Phi(Z) \geq \sigma$; we include oracles for which $\Phi(Z)$ is not total. Then $\sigma \mapsto \lambda(\Phi^{-1}[\sigma])$ (here λ denotes Lebesgue measure on Cantor space 2^ω) is a continuous c.e. semimeasure (multiplied by $2^{|\sigma|}$ it is a c.e. supermartingale). Since X is ML-random, $\lambda(\Phi^{-1}[X \upharpoonright_n]) \leq 2^{-n}$. By withholding computations, we can massage the functional Φ so that $\lambda(\Phi^{-1}[\sigma]) \leq 2^{-|\sigma|}$ for all σ (but still $\Phi(Y) = X$). Using the massaged functional we can pull back any strong test $\langle U_n \rangle$ which captures X (a difference test, a weak 2-test, a Demuth test, a \mathbf{d} -ML-test,...) and obtain a similar test which captures Y .

The key to this argument is the continuity of the map Φ on 2^ω . The reducibility \leq_h (relatively hyperarithmetic) is not given by partial continuous functionals. And indeed some of the examples above fail in the higher setting. Hjorth and Nies [HN07] introduced the notion of Π_1^1 -ML-randomness, the higher analogue of ML-randomness; Nies [Nie09, 9.2.17] introduced the notion of strong Π_1^1 -ML-randomness, the higher analogue of weak 2-randomness, studied later by Chong and Yu [CY15]. There are however reals X and Y such that $X \leq_h Y$, Y is strongly Π_1^1 -ML-random, and X is Π_1^1 -ML-random but not strongly so.

Rather than use \leq_h , we need a continuous higher analogue of Turing reducibility. For preciseness, recall that a functional is simply a set of pairs (τ, σ) of finite binary strings. Looking forward, note that we do not require that the functional be consistent; we discuss this shortly. If Φ is a functional then for any $X \in 2^{\leq \omega}$ (finite or infinite) we let

$$\Phi(X) = \bigcup \{ \sigma : (\tau, \sigma) \in \Phi \text{ for some } \tau \leq X \}^1.$$

For $X, Y \in 2^\omega$, $X \leq_T Y$ if and only if $\Phi(Y) = X$ for some c.e. functional Φ . This motivates the following definition:

Definition 1.1. Let $X, Y \in 2^\omega$. X is *higher Turing reducible* to Y if $\Phi(Y) = X$ for some Π_1^1 functional Φ . We write $X \leq_{\omega_1^{\text{ck}_T}} Y$.

With this notion some of the familiar theorems mentioned above generalise to the higher setting. For example, we will show:

Theorem 1.2. *Let X, Y be Π_1^1 -ML-random. Suppose that $X \leq_{\omega_1^{\text{ck}_T}} Y$ and that Y is in fact strongly Π_1^1 -ML-random. Then X too is strongly Π_1^1 -ML-random.*

¹Here the union uses the set-theoretic concept of functions as sets of ordered pairs. So $(x, y) \in \Phi(X)$ if and only if there is some $\tau \leq X$ and σ such that $\sigma(x) = y$ and $(\tau, \sigma) \in \Phi$.

We will also see, for example, that a Π_1^1 -ML-random set X is higher difference random if and only if $O \not\leq_{\omega_1^{\text{ck}}\text{T}} X$, where Kleene's O is the complete Π_1^1 set of numbers. On the other hand, we will see that some results only partially generalise, or completely fail in the higher setting. For example, a Martin-Löf random real is weak-2-random if and only if it forms a minimal pair with \emptyset' , but we show in [GM] that it is not the case that a Π_1^1 -ML-random set is strongly Π_1^1 -ML-random if and only if it forms a minimal pair with Kleene's O in the $\leq_{\omega_1^{\text{ck}}\text{T}}$ -degrees.

Continuity also matters when it comes to relativizing randomness notions. The two-step product theorem (see for example [Jec08] or [Kun11]) says that if \mathbb{P} and \mathbb{Q} are notions of forcing then a filter $G \times H \subset \mathbb{P} \times \mathbb{Q}$ is V -generic if and only if the filter $G \subset \mathbb{P}$ is V -generic and $H \subset \mathbb{Q}$ is $V[G]$ -generic. The theorem has effective analogues. For example, a join $G \oplus H$ is (Cohen) 1-generic if and only if G is 1-generic and H is 1-generic relative to G (see for example [Yu06]). van Lambalgen [vL87] gave an analogous effectivisation for ML-randomness. It fails in the higher setting: there are reals X and Y such that $X \oplus Y$ is Π_1^1 -ML-random, but Y is not $\Pi_1^1(X)$ -ML-random. The reason for this failure is that the relativisation is not continuous: enumerating clopen subsets of a component of a $\Pi_1^1(X)$ -ML-test is not determined by only finitely many bits of X . Similarly to Turing reducibility, we need to define a continuous higher analogue of being computably enumerable relative to an oracle. The treatment is similar. An *enumeration functional* is a set of pairs (τ, m) consisting of a finite binary string and a natural number. If Ψ is an enumeration functional and $X \in 2^{\leq \omega}$ then we let

$$\Psi^X = \{m : (\tau, m) \in \Psi \text{ for some } \tau \leq X\}.$$

A set B is c.e. in X if and only if $B = \Psi^X$ for some c.e. enumeration functional Ψ .

Definition 1.3. Let $X \in 2^\omega$. A set $B \subseteq \omega$ is *higher X -c.e.* if $B = \Psi^X$ for some Π_1^1 enumeration functional Ψ .²

Armed with this definition we can consider higher X -c.e. open sets (sets of the form $\bigcup_{\sigma \in B} [\sigma]$ where B is a higher X -c.e. set of strings), and so higher X -ML-tests and higher X -ML-randomness. Thus Π_1^1 -ML-randomness is simply higher \emptyset -ML-randomness, and so we call it “higher ML-randomness”. We will show that this continuous relativisation satisfies van Lambalgen's theorem.

Theorem 1.4. *Let $X, Y \in 2^\omega$. Then $X \oplus Y$ is higher ML-random if and only if X is higher ML-random and Y is higher X -ML-random.*

The issue of continuous relativisation is directly related to the study of anti-randomness and lowness for randomness. A celebrated result of Nies's (together with work by Hirschfeldt, Nies and Stephan [Nie05, HNS07]) is the coincidence of a number of classes, each formalising a notion of distance from randomness or weakness as an oracle in detecting randomness: the K -trivial sets; the sets which are low for ML-randomness; the sets which are low for K ; and the sets which are a base for ML-randomness. Hjorth and Nies [HN07] showed that this result fails in the higher setting: while there are sets which are higher K -trivial but not hyperarithmetical, every set which is low for Π_1^1 -ML-randomness is hyperarithmetical.

²We remark that we can think of an enumeration functional as an open subset of $2^\omega \times \omega$. If Ψ is such a set then Ψ^X is the X -section of Ψ .

(Higher K -triviality is defined using a Π_1^1 analogue of prefix-free Kolmogorov complexity.) Again this uses the fact that the relativisation of Π_1^1 -ML-randomness used in the definition of lowness for this notion is not continuous. We will show that using continuous relativisation the coincidence does hold:

Theorem 1.5. *The following are equivalent for $A \in 2^\omega$:*

- (1) *A is higher K -trivial.*
- (2) *Every higher ML-random set is also higher A -ML-random.*
- (3) *There is some higher A -ML-random set X such that $A \leq_{\omega_1^{\text{ck}_T}} X$.*

We will also discuss lowness for K .

1.2. A general method for defining higher analogues. The two examples we gave of higher analogues of basic concepts of computability (Turing reducibility and relative computable enumerability) follow a common method which is already implicit in the Chong-Yu work and which we will employ everywhere. We realise that the most fundamental concept of computability theory is computable enumerability. From it, all other notions can be derived: a partial computable function is one with c.e. graph, Turing reducibility is defined using c.e. functionals, etc. Recall again that a set of numbers is Π_1^1 if and only if it is Σ_1 -definable over $L_{\omega_1^{\text{ck}}}$; in the terminology of higher computability, it is ω_1^{ck} -c.e. The method of obtaining higher analogues is to replace every instance of “c.e.” by “ ω_1^{ck} -c.e.”. As we observed, this means that “higher ML-randomness” is the notion of Π_1^1 -ML-randomness defined by Hjorth and Nies; and “higher weak 2-randomness” is the notion of strong Π_1^1 -ML-randomness defined by Chong, Nies and Yu. It is only the basic notion of computable enumerability which is being modified; all other quantifiers range over the natural numbers (rather than ω_1^{ck}), and unlike metarecursion theory, the objects studied are subsets of ω rather than subsets of ω_1^{ck} . For example, a higher ML-test is an ω -sequence of (uniformly) higher c.e. open sets, rather than a sequence of length ω_1^{ck} . The fact though that the basic existential quantifier (the computable unbounded search) ranges over ω_1^{ck} motivates some of our notation (such as $\leq_{\omega_1^{\text{ck}_T}}$).

1.3. Continuity and its discontent. Beyond the inherent interest in higher notions, the study of generalisations of computability sheds light on the familiar notions by separating concepts which “accidentally” coincide in usual computability. An example of such a phenomenon is directly related to the examples of the use of continuity that we discussed above.

Consider the definition of higher Turing reducibility. The definition of Turing reducibility in terms of functionals usually imposes extra requirements of *consistency* on the functional. Namely that if (τ, σ) and (τ', σ') are two “axioms” in the functional Φ and τ and τ' are compatible, then σ and σ' are compatible. Indeed, in [HN07] Hjorth and Nies introduce a continuous reducibility (which they denote by $\leq_{\text{fin-h}}$). Their definition is similar to Definition 1.1 except that they require that the functional Φ be the graph of an order-preserving function from strings to strings and moreover that its domain is closed under taking initial segments. In “traditional” (or “countable”) computability this extra requirement creates no difficulty. Namely $X \leq_T Y$ if and only if $X = \Phi(Y)$ for some c.e. functional Φ ³ if and only if $X = \Phi(Y)$ for some consistent c.e. functional if and only if $X = \Phi(Y)$

³If Φ is an inconsistent Turing functional and two inconsistent axioms in Φ apply to an oracle Y then $\Phi(Y) \notin 2^\omega$ and so Y does not compute anything with the functional Φ .

for some c.e. functional satisfying the definition of Hjorth and Nies. We will show in [BGHM] that the higher analogues of the two first notions are distinct, while the second one coincide with the third one, but not uniformly. In this paper we will argue that among these two distinct reducibilities, the one given by Definition 1.1 is the one which fits best with the general theory of higher randomness.

It may be instructive to see why the argument that traditionally these reducibilities are the same fails in the higher setting. To turn an arbitrary functional into a consistent one (without losing total computations), when an axiom (τ, σ) enters the functional at some stage s , we consider all extensions of τ of length s , and map those among them to σ for which this does not introduce an inconsistency. This argument uses what we call a “time trick”: the fact that the number of stages is the same as the length of the oracle, namely ω . This equality fails in the higher setting, in which we still use oracles of length ω but effective constructions have ω_1^{ck} many stages. Thus any argument that relies on a time trick cannot be simply copied in the higher setting. In some cases the argument can be rectified (an example is the proof of the higher Kraft-Chaitin theorem by Hjorth and Nies). In other cases, such as the equivalence of the three definitions of Turing reducibility, in the higher setting the theorem fails.

To give evidence that Definition 1.1 is more useful than other possible generalisations of Turing reducibility to the higher setting, consider for example one of the most basic properties of relative computability. The following is easily verified using arguments of general computability:

Proposition 1.6. *The following are equivalent for $X, Y \in 2^\omega$:*

- (1) $X \leq_{\omega_1^{\text{ckT}}} Y$.
- (2) *Both X and its complement are higher Y -c.e.*

The proposition fails if we replace $\leq_{\omega_1^{\text{ckT}}}$ by its stricter variant. The difficulty is in the direction (2) \implies (1): suppose that $\Psi_1^Y = X$ and $\Psi_0^Y = \omega - X$. We build a functional Φ with the aim that $\Phi(Y) = X$. When we see strings τ and σ such that $\Psi_0^\tau \supseteq \{n : \sigma(n) = 0\}$ and $\Psi_1^\tau \supseteq \{n : \sigma(n) = 1\}$ we enumerate the axiom (τ, σ) into Φ . It is possible that for other oracles Z , Ψ_1^Z and Ψ_0^Z do not enumerate a set and its complement. But before we see this fact, at earlier stages, computations corresponding to such oracles Z appear to give a set and its complement — inconsistent with σ — and enumerate into Φ axioms (with use extending τ but incomparable with Y) which are inconsistent with (τ, σ) . The current stage may be infinite (a stage $s \in [\omega, \omega_1^{\text{ck}})$), and so such an event could have happened arbitrarily close to Y (i.e. extending longer and longer initial segments τ of Y). Thus even if we take τ to be an arbitrarily long initial segment of Y , enumerating (τ, σ) into Φ makes Φ inconsistent; of course $\Phi(Y)$ does not contain inconsistencies. This shows how the time trick can fail bitterly. In [BGHM] we show how to turn this situation around to prove, for example, that the two generalisations of Turing reducibility are distinct: there exists some X, Y such that $X \leq_{\omega_1^{\text{ckT}}} Y$ but not via a functional which is consistent everywhere.

The utility of Definition 1.1 with respect to randomness is witnessed in Theorem 1.5 (in the notion of a higher base for randomness) and also in the example of difference randomness. The following theorem is the correct generalisation of a theorem of Franklin and Ng’s. We delay the proper definition of “higher ω -computably approximable” to the next subsection.

Theorem 1.7. *The following are equivalent for a higher ML-random set X :*

- (1) $O \not\leq_{\omega_1^{\text{ckT}}} X$ (where O is Kleene's complete Π_1^1 set).
- (2) X avoids all nested tests of the form $\langle U_n \cap P \rangle$ where $\langle U_n \rangle$ are uniformly higher effectively open, P is higher effectively closed (a closed Σ_1^1 set of reals), and $\lambda(U_n \cap P) \leq 2^{-n}$.
- (3) X avoids all nested tests of the form $\langle W_{f(n)} \rangle$ where W_e (for $e < \omega$) is the e^{th} higher effectively open set; $\lambda(W_{f(n)}) \leq 2^{-n}$; f is higher ω -computably approximable, witnessed by $\langle f_s \rangle_{s < \omega_1^{\text{ck}}}$; and if $f_s(n) \neq f_t(n)$ then $W_{f_s(n)}$ and $W_{f_t(n)}$ are disjoint.

The proof is the same as in [FN11]. We note where we use the fact that inconsistent functionals are allowed. In proving (1) \implies (3) the functional Γ which we build determines that $\Gamma(\tau)(n) = O_s(n)$ where s is the least such that $[\tau] \subseteq W_{f(n)}[s]$. On the elements of the Solovay test $\{W_{f(n)}[s] : n \in O_{s+1} - O_s\}$ this functional may be inconsistent. In fact, in [BGHM] we show that there is a higher ML-random sequence which is higher Turing above O , but is not fin-h above O .

Similarly, our definition of the relativisation of higher ML-randomness runs into consistency problems when we try to construct a uniform universal test. Classically there is a sequence $\langle U_n \rangle$ of enumeration operators such that for all Z , $\langle U_n^Z \rangle$ is a universal Z -ML-test. This fails in the higher setting. The point is that we cannot take a higher c.e. operator (a Π_1^1 enumeration functional) U and produce another such functional V such that $\lambda(V^Z) \leq \varepsilon$ for all Z (for some fixed ε), and such that $U^Z = V^Z$ if $\lambda(U^Z) \leq \varepsilon$. Again a time trick fails. In a sense it is a topological problem. In standard computability, at a finite stage s the collection of reals for which an axiom of U_s applies is clopen. When the axiom (τ, σ) enters U (indicating that $[\sigma] \subseteq U^Z$ for all $Z \in [\tau]$) we let $C = \{Z \in 2^\omega : \lambda([\sigma] \cup V_s^Z) > \varepsilon\}$; this set is clopen and so we can let V enumerate $[\sigma]$ with oracles in the clopen set $[\tau] - C$. In the higher setting, C is open but may fail to be clopen, as s may be infinite. Indeed C could be dense. There may be no way to add $[\sigma]$ to V^Z for reals Z outside C without making $\lambda(V^Z) > \varepsilon$ for some reals $Z \in C$.

This is an issue we will need to monitor; in some cases we can find work-arounds to get analogues of “lower” results. In other cases this is impossible. In [BGHM] we not only show that there is no uniform universal oracle higher ML-test; indeed we construct an oracle A for which there is no universal higher A -ML-test.

Similarly we can construct an oracle relative to which there is no optimal higher discrete c.e. semimeasure, and so an oracle relative to which higher prefix-free Kolmogorov complexity K^A is not defined. Thus we need to modify the definition of “higher low for K ”, to say that every higher discrete A -c.e. semimeasure is dominated by the optimal higher c.e. one. Of course if A is low for higher K then higher K^A exists, namely it is higher K . We will show that this notion coincides with higher K -triviality as well.

1.4. The higher limit lemma. The analysis of functions approximable by hyperarithmetic functions corresponds to that given to Δ_2^0 sets and functions by Shoenfield's limit lemma. Here Kleene's O plays the role of the halting problem \emptyset' . This analysis will help us separate notions of higher randomness.

Recall that a sequence $\langle f_s \rangle_{s < \omega_1^{\text{ck}}}$ is ω_1^{ck} -computable if it is Σ_1 -definable over $L_{\omega_1^{\text{ck}}}$. Such a sequence is a ω_1^{ck} -approximation of a function $f \in \omega^\omega$ if for all n there is

some $s < \omega_1^{\text{ck}}$ such that $f_t(n) = f(n)$ for all $t \in [s, \omega_1^{\text{ck}})$. In Section 6.1 we shall prove:

Proposition 1.8. *The following are equivalent for $f \in \omega^\omega$:*

- (1) $f \leq_{\omega_1^{\text{ck}}\text{T}} O$;
- (2) $f \leq_{\text{T}} O$;
- (3) f has an ω_1^{ck} -computable approximation.

Since a subset of ω is c.e. in O if and only if it is Σ_2 definable over $L_{\omega_1^{\text{ck}}}$, the functions computable from Kleene's O are the functions which are Δ_2 -definable over $L_{\omega_1^{\text{ck}}}$. Thus we call such functions ‘‘higher Δ_2^0 ’’. We will investigate subclasses of the collection of all higher Δ_2^0 functions (such as the higher ω -c.a. functions which we define below). These classes are related to notions of randomness in two ways:

A. In the style of Demuth, we can use higher Δ_2^0 functions to give indices for higher effectively open components of tests: tests of the form $\langle W_{f(n)} \rangle$ where W_e is the e^{th} higher c.e. open set. The strongest such notion is higher $\text{MLR}[O]$, for which we use all functions $f \leq_{\text{T}} O$. Unlike lower computability, this is strictly stronger than the higher version of weak 2-randomness; indeed strictly stronger than Π_1^1 -randomness.

B. We can study Δ_2^0 properties according to their approximability. In the lower setting this involves classes determined by bounding the number of mind changes. For example in [FHM⁺15] Figueira et al. show that while there is a ML-random sequence with an approximation whose first n bits change at most 2^n many times, no such random can be superlow. The general theme is that among random sequences, approximations with few changes correspond to computational strength.

In the higher setting we identify a number of classes of functions lying between the higher ω -c.a. functions and all higher Δ_2^0 functions. In some sense they too are described by conditions about how often the approximation changes. However these conditions are qualitative rather than quantitative. Thus these classes have no lower analogues.

Definition 1.9. Let $\langle f_s \rangle$ be an ω_1^{ck} -computable approximation of a higher Δ_2^0 function f . For $n < \omega$ let $s(n)$ be the least stage $s < \omega_1^{\text{ck}}$ such that $f_s \upharpoonright_n = f \upharpoonright_n$. The approximation $\langle f_s \rangle$ is *collapsing* if $\sup_n s(n) = \omega_1^{\text{ck}}$. Equivalently, $\langle f_s \rangle$ is collapsing if for all $s < \omega_1^{\text{ck}}$, f does not belong to the closure of the set $\{f_t : t < s\}$.

Gandy's basis theorem implies that there is an O -computable Π_1^1 -random sequence, and so a Π_1^1 -random sequence with some ω_1^{ck} -computable approximation. However no such random sequence can have a collapsing approximation, since the sequence $\langle s(n) \rangle$ is Σ_1 -definable over $(L_{\omega_1^{\text{ck}}}, f)$, and so if f has a collapsing approximation then $\omega_1^f > \omega_1^{\text{ck}}$ (f collapses ω_1^{ck}). Roughly, the intuition here is that an approximation of a Π_1^1 -random sequence X must change so much so that all initial segments of X appear long before the end of the approximation. We note though that there are sets $X \leq_{\text{T}} O$ which collapse ω_1^{ck} but do not have a collapsing approximation.

Some of the classes we consider are defined by topological conditions. For example:

Definition 1.10. An ω_1^{ck} -computable approximation $\langle f_s \rangle$ of a function f is *compact* if the set $\{f_s : s < \omega_1^{\text{ck}}\} \cup \{f\}$ is a compact subset of Baire space ω^ω .

Of course if we approximate an element of Cantor space we may assume that all elements of the approximation are also elements of Cantor space. In that case an approximation is compact if and only if it is closed (for the usual topology).

Lemma 1.11. Suppose that $\langle f_s \rangle$ is a compact approximation of a function $f \notin \Delta_1^1$. Then $\langle f_s \rangle$ is a collapsing approximation.

Proof. Let $s(n)$ be defined as above. Suppose that $s(\omega) = \sup_n s(n)$ is a computable ordinal. Consider the closure A of the set $\{f_t : t < s(\omega)\}$. The function f is an element of A . However A is countable, as it is contained in the compact set $\{f_t : t < \omega_1^{\text{ck}}\} \cup \{f\}$. Further, A is the set of paths of a finitely branching hyperarithmetic tree with a hyperarithmetic bound on its branching. Running the Cantor-Bendixon analysis of closed sets within $L_{\omega_1^{\text{ck}}}$ we see that every element of A is hyperarithmetic, and so f is. \square

We will show that no higher weakly 2-random set can have a closed approximation. Thus, to separate Π_1^1 -randomness from higher weak 2-randomness we will need to find a class strictly between compact approximations and collapsing approximations.

Narrowing our classes further we return to the idea of counting the number of changes. Finite-change approximations have been implicitly used by Yu [Yu11].

Definition 1.12. An ω_1^{ck} -computable approximation $\langle f_s \rangle$ is a *finite-change* approximation if for no n is there an increasing infinite sequence $\langle t(i) \rangle_{i < \omega}$ of stages such that $f_{t(i+1)}(n) \neq f_{t(i)}(n)$ for all $i < \omega$.

Note that it is not enough to require that there are only finitely many stages s such that $f_{s+1}(n) \neq f_s(n)$. For it is possible that there are limit stages s at which a new value is given. On the other hand, if $\langle f_s \rangle$ changes only finitely often then for all limit s , $\lim_{t \rightarrow s} f_t$ exists. Since this limit is ω_1^{ck} -computable from s , we may assume that for all limit s , $f_s = \lim_{t \rightarrow s} f_t$. In this case, we can indeed define the *number of changes* on n to be the number of stages s such that $f_{s+1}(n) \neq f_s(n)$. Without this assumption we can define the number of changes to be the longest length of any increasing sequence $\langle t(i) \rangle$ of stages such that $f_{t(i+1)}(n) \neq f_{t(i)}(n)$. To pay a debt, we mention the definition of higher ω -c.a. functions.

Definition 1.13. A *higher ω -computable approximation* is a finite-change ω_1^{ck} -computable approximation $\langle f_s \rangle$ for which the number of changes is bounded by a hyperarithmetic function.

Like its lower analogue, a function has a higher ω -computable approximation if and only if it is higher truth-table reducible to O .

Suppose that $\langle f_s \rangle$ is a finite-change approximation which has been modified so that $f_s = \lim_{t \rightarrow s} f_t$ for all limit ordinals s . Then the set $\{f_s : s < \omega_1^{\text{ck}}\} \cup \{f\}$ is a closed subset of Baire space. Further, because this is a finite-change approximation, it is contained in the set of paths of a finitely branching subtree of $\omega^{<\omega}$, which is compact. Hence:

Lemma 1.14. If f has a finite-change approximation then it has a compact approximation.

The simplest finite-change approximation is an ω_1^{ck} -enumeration of a Π_1^1 set, or a monotone approximation of a higher left-c.e. (left- Π_1^1) real. Chong and Yu showed [CY15] that a higher left-c.e. sequence cannot be higher weak 2-random. Their proof used the Lebesgue density theorem. Lemma 1.14 and Proposition 5.1 give a new proof of their result. They also answer Yu's question whether the two halves of higher Ω are Π_1^1 -random or not. Since they both have a finite-change approximation, they are not even higher weakly 2-random. Indeed this gives us a separation of higher weak 2-randomness from higher difference randomness, since the two halves of higher Ω do not higher compute each other and so are $\leq_{\omega_1^{\text{ck}}\text{T}}$ -incomplete.

2. EXTREMES OF HIGHER TURING AND HIGHER C.E.

Before we discuss randomness we investigate the notions of higher relative computability and enumeration, in particular when they coincide with familiar notions. With very strong oracles they collapse to the familiar notions of Turing reducibility and relative computable enumerability. With weak oracles they coincide with relative Δ_1^1 and Π_1^1 .

2.1. Higher computability and strong oracles. The fact that all Π_1^1 sets are O -computable implies the following:

Proposition 2.1. *A set is higher O -c.e. if and only if it is O -c.e.; and so a set is higher O -computable if and only if it is O -computable. Furthermore, these equivalences hold when O is replaced by any oracle $Y \geq_{\text{T}} O$.⁴*

As mentioned in the introduction, there is an effective higher enumeration of all Π_1^1 sets, and so we can define an effective higher enumeration of all higher enumeration functionals. We will use the familiar notation $\langle W_e \rangle$ to denote such an enumeration. We will never use both c.e. sets and Π_1^1 sets in the same context so no confusion should arise. The enumeration gives rise to a higher jump operator $Y \mapsto J^Y = \bigoplus_{e < \omega} W_e^Y$, for which we easily verify $Y <_{\omega_1^{\text{ck}}\text{T}} J^Y$ for every Y . Since Proposition 2.1 is uniform in the indices for O -c.e. and higher O -c.e. sets, we see that in the particular case where $Y \geq_{\text{T}} O$, the higher jump J^Y and the standard Turing jump J' are recursively isomorphic. On the other hand, O is recursively isomorphic to J^\emptyset .

Reals which have collapsing approximations (Definition 1.9) are computationally strong in that they compute a copy of ω_1^{ck} . Recall that $X \leq_{\text{fin-h}} Y$ if $X \leq_{\omega_1^{\text{ck}}\text{T}} Y$ via a strongly consistent functional: one whose graph is a monotone function from strings to strings, whose domain is closed under taking initial segments.

Proposition 2.2. *Suppose that $Y \in 2^\omega$ has a collapsing approximation. Then for every higher Y -computable set X we actually have $X \leq_{\text{fin-h}} Y$.*

⁴ A more elaborate proof of this Proposition will be used later, so we detail it here. Kleene's O computes a bijection between ω and ω_1^{ck} , and so relative to O , quantifiers ranging over ω_1^{ck} can be transformed to quantifiers ranging over ω . Formally, there is an O -computable binary relation $E \subset \omega^2$ such that $(\omega, E) \cong (L_{\omega_1^{\text{ck}}}, \epsilon)$, and further, such that $f \upharpoonright_\omega$ is O -computable, where $f: L_{\omega_1^{\text{ck}}} \rightarrow (\omega, E)$ is the unique isomorphism. Every set which is higher O -c.e. is $\Sigma_1((L_{\omega_1^{\text{ck}}}, \epsilon), O)$ -definable, and so, if X is higher O -c.e. then $X = f^{-1}Z$ where Z is Σ_1 -definable in the structure (ω, E) and so Z (and so X) is O -c.e.

Proof. Let Φ be a higher Turing functional such that $\Phi(Y) = X$, and let $\langle Y_s \rangle$ be a collapsing approximation for Y . We may assume that for all $\sigma \in 2^{<\omega}$, $|\Phi(\sigma)| \leq |\sigma|$. We define a fin-h functional Ψ by recursion, by selectively copying Φ -computations. At stage s let Ψ_s consist of all the axioms already enumerated into Ψ by stage s . For every $n < \omega$, if:

- $Y_s \upharpoonright_n$ is not in the domain of Ψ_s ; and
- $\Phi_s(Y_s \upharpoonright_n)$ is consistent,

then we enumerate an axiom mapping $Y_s \upharpoonright_n$ to $\Phi_s(Y_s \upharpoonright_n)$ into Ψ_{s+1} . Then Ψ is a fin-h functional. It suffices to show that $\Psi(Y)$ is total. Let $n < \omega$ and let $s(n)$ be the least s such that $Y_s \upharpoonright_n = Y \upharpoonright_n$. Since the approximation is collapsing, there is some $k \geq n$ such that $\Phi_{s(k)}(Y \upharpoonright_n) = \Phi(Y \upharpoonright_n)$. Also, $Y \upharpoonright_k$ is not in the domain of $\Psi_{s(k)}$, and $\Phi(Y \upharpoonright_k)[s(k)]$ is consistent and extends $\Phi(Y \upharpoonright_n)$. It follows that $\Psi_{s(k)+1}(Y) \supseteq \Phi(Y \upharpoonright_n)$. \square

On the other hand we know that there are O -computable sets X and Y such that $X \leq_{\omega_1^{\text{ck}} \text{T}} Y$ but $X \not\leq_{\text{fin-h}} Y$, so some assumption on the nature of the approximations is necessary.

2.2. Higher computability and relative Π_1^1 .

Proposition 2.3. *Suppose that Y preserves ω_1^{ck} (that is, $\omega_1^Y = \omega_1^{\text{ck}}$). Then for all X , $X \leq_{\omega_1^{\text{ck}} \text{T}} Y$ if and only if $X \leq_{\text{T}} Y \oplus H$ for some hyperarithmetic set H .*

Proof. If H is hyperarithmetic and $X \leq_{\text{T}} Y \oplus H$ then we can easily devise a hyperarithmetic functional Φ such that $\Phi(Y) = X$, and so $X \leq_{\omega_1^{\text{ck}} \text{T}} Y$.

In the other direction, suppose that Φ is a Π_1^1 functional, $\Phi(Y) = X$ and $\omega_1^Y = \omega_1^{\text{ck}}$. Let $\langle \Phi_s \rangle_{s < \omega_1^{\text{ck}}}$ be an effective enumeration of Φ . Define $f: \omega \rightarrow \omega_1^{\text{ck}}$ by letting $f(n)$ be the least stage $s < \omega_1^{\text{ck}}$ such that $\Phi_s(Y)$ extends $X \upharpoonright_n$. The function f is Δ_1 -definable over $L_{\omega_1^{\text{ck}}}(Y)$; since Y preserves ω_1^{ck} , f is bounded below ω_1^{ck} . Let $s < \omega_1^{\text{ck}}$ bound the range of f . Then $\Phi_s(Y) = X$ and so $X \leq_{\text{T}} Y \oplus \Phi_s$; and Φ_s is hyperarithmetic. \square

For $Y \in 2^\omega$ we let $\Delta_1^1 \oplus Y$ be the class of sets Turing reducible to $H \oplus Y$ for some hyperarithmetic set H . Thus Proposition 2.3 says that if Y preserves ω_1^{ck} then $\Delta_1^1 \oplus Y$ is the class of sets higher Turing reducible to Y . Unfortunately the proposition cannot be reversed. This can be seen by considering the Borel rank of the set of oracles for which $\Delta_1^1 \oplus Y$ equals the collection of sets higher Turing reducible to Y , which is fairly low, whereas the Borel rank of the reals which collapse ω_1^{ck} is high (precisely $\Sigma_{\omega_1^{\text{ck}}+2}^0$ [Ste78]). Alternatively we can observe that if $Y \geq_{\text{T}} O$, then $\Delta_1^1 \oplus Y$ is of course the collection of Y -computable sets, which by Proposition 2.1 equals the collection of sets higher Turing reducible to Y .

Remark 2.4. Let Ω denote the higher version of Chaitin's left-c.e. random number. A standard argument shows that $\Omega \equiv_{\omega_1^{\text{ck}} \text{T}} O$, indeed the equivalence is higher weak-truth-table. However since higher Ω is Δ_1^1 -random it does not (Turing) compute any noncomputable hyperarithmetic set, let alone Kleene's O , nor is O Turing reducible to $\Omega \oplus H$ for any hyperarithmetic set H . This shows that the conclusion of Proposition 2.3 fails for the oracle Ω .

It is well-known that for sufficiently Cohen generic, sufficiently random and sufficiently Sacks generic (with respect to forcing with hyperarithmetic perfect sets)

sets Y , $\Delta_1^1(Y) = \Delta_1^1 \oplus Y$; we discuss this shortly. Note that this equality does imply that Y preserves ω_1^{ck} ; if $\omega_1^Y > \omega_1^{\text{ck}}$ then $Y^{(\omega_1^{\text{ck}})}$ is not in $\Delta_1^1 \oplus Y$. Thus, if $\Delta_1^1(Y) = \Delta_1^1 \oplus Y$ then for all X , $X \leq_h Y$ if and only if $X \leq_{\omega_1^{\text{ck}}T} Y$.

We require a notion of uniformity for this equality. First we settle some notation.

Notation 2.5. We sometimes blur the distinction between notations for ordinals and the ordinals they denote: if $\alpha \in O$ then we let α denote also the ordinal $|\alpha|_O$; we let $\alpha + 1$ be the notation for the successor of α , and so on. For $\alpha \in O^Y$, we let $Y^{(\alpha)} = H_\alpha^Y$ be the iteration of the Turing jump along α .

Definition 2.6. Let $Y \in 2^\omega$. We say that $\Delta_1^1(Y) = \Delta_1^1 \oplus Y$ *uniformly in Y* if there is a Turing functional Ψ and a higher Y -partial computable function g (a function whose graph is higher Y -c.e.) such that for all $\alpha \in O^Y$, $g(\alpha) \in O$ and $Y^{(\alpha)} = \Psi(Y, \emptyset^{g(\alpha)}, \alpha)$.

Recall that a Y -hyperarithmetical index for a set $A \in \Delta_1^1(Y)$ is a pair (e, α) where $\alpha \in O^Y$ and $A = \Phi_e(Y^{(\alpha)})$ (where here Φ_e is the e^{th} (*lower*) Turing functional). Similarly, a $\Delta_1^1 \oplus Y$ -index for a set A is a pair (e, a) where a is a hyperarithmetical index for a set $H \in \Delta_1^1$ and $A = \Phi_e(H, Y)$. Then $\Delta_1^1(Y) = \Delta_1^1 \oplus Y$ uniformly in Y if there is a higher Y -partial computable method of transforming a Y -hyperarithmetical index for a set $A \in \Delta_1^1(Y)$ to a $\Delta_1^1 \oplus Y$ -index for the same set. (The reverse direction is uniform for all oracles.)

Proposition 2.7. *The following are equivalent for $Y \in 2^\omega$:*

- (1) *A set is higher Y -c.e. if and only if it is $\Pi_1^1(Y)$.*
- (2) *$\Delta_1^1(Y) = \Delta_1^1 \oplus Y$ uniformly in Y .*

Proof. Assume (1). Note that since there are universal $\Pi_1^1(Y)$ and higher Y -c.e. sets, the equivalence is uniform: there are computable functions translating between $\Pi_1^1(Y)$ -indices and higher Y -c.e. indices. Given this, we see that the proof of Proposition 2.3 can be performed effectively in Y , as follows. Given $\alpha \in O^Y$ we obtain indices for higher enumeration functionals which with oracle Y enumerate $A = Y^{(\alpha)}$ and its complement. As a result we obtain an index for a higher Turing functional Φ such that $A = \Phi(Y)$ (Proposition 1.6 is uniform). The relation “ $\Phi_s(Y)$ is total” is Δ_1 -definable over $L_{\omega_1^{\text{ck}}}(Y)$ (uniformly in Φ and $s < \omega_1^{\text{ck}}$); the argument of Proposition 2.3 gives us a function g satisfying $Y^{(\alpha)} = \Psi(Y, \emptyset^{g(\alpha)}, \alpha)$ which is $\Pi_1^1(Y)$ -definable. Applying (1) again, we see that g is higher Y -partial computable.

Assume (2), and let g witness the uniformity. We recall that we can view O as a subset of O^Y (as the set of notations in O^Y which hereditarily do not look at the oracle Y when computing increasing sequences of notations). Uniformly in $\alpha \in O$ we can get a $\Delta_1^1(Y)$ -index for

$$O_\alpha^Y = \{\beta \in O^Y : \beta < \alpha\};$$

and the point is that $O^Y = \bigcup_{\alpha \in O} O_\alpha^Y$, as Y preserves ω_1^{ck} . Using g and varying over $\alpha \in O$ we see how to enumerate O^Y in a higher Y -c.e. fashion. \square

Porism 2.8. In Proposition 2.7 we may replace the definition of uniformity of $\Delta_1^1(Y) = \Delta_1^1 \oplus Y$ by the apparently weaker condition that $Y^{(\alpha)} = \Psi(Y, \emptyset^{g(\alpha)}, \alpha)$ for all $\alpha \in O$ (rather than all $\alpha \in O^Y$). Spector showed (see [Sac90, II2.4]) that

there is a Turing functional Γ such that for all $\alpha \in O^Y$, $O_\alpha^Y = \Gamma(Y^{(\alpha+1)})$. If $\alpha \in O$ then $\alpha + 1 \in O$. In the proof of (2) \implies (1) we apply g to $\alpha + 1$.

2.3. The behaviour of generics for various forcing notions. We discuss Proposition 2.7 in the context of Cohen genericity, randomness and Sacks genericity. The ideas here are certainly not new, but some are hard to find in print in the form below.

2.3.1. *Cohen generics.* It is well-known, via the analysis of Cohen forcing, that if G is Cohen generic then $\Delta_1^1(G) = \Delta_1^1 \oplus G$ uniformly. We will employ the following direct definition of the class of Σ_α^0 sets; see for example [AK00].

Definition 2.9. For $\alpha \in O$ we define the class of Σ_α^0 sets (of numbers and of reals) and indices for these sets. For $\alpha = 1$, the Σ_1^0 sets are the c.e. sets (and c.e. open sets of reals), with $(e, 1)$ being the index of the e^{th} such set in some effective listing.

Let $\alpha > 1$. A set is $\Sigma_{<\alpha}^0$ if it is Σ_β^0 for some $\beta <_O \alpha$. A $\Sigma_{<\alpha}^0$ -index for such a set is a Σ_β^0 -index for some $\beta <_O \alpha$. A set A is Σ_α^0 if it is the effective union of $\Pi_{<\alpha}^0$ sets. That is, if there is a c.e. set W such that A is the union of the complements of the sets whose $\Sigma_{<\alpha}^0$ -indices are in W . The Σ_α^0 -index for this union is (e, α) , where W is the e^{th} c.e. set. Note that we do not require that all elements of W are $\Sigma_{<\alpha}^0$ indices, so (e, α) is a Σ_α^0 code for all $e < \omega$.

Let $Y \in 2^\omega$. Note that $Y^{(\omega)}$ is not a $\Sigma_\omega^0(Y)$ -complete set, as it is only $\Delta_\omega^0(Y)$ (a uniform disjoint union of $\Sigma_n^0(Y)$ sets for n unbounded). For this reason, we use alternative notation (used in [GMS13], following ideas from [AK00]) denoting $\Sigma_\alpha^0(Y)$ -complete sets. For $n < \omega$ let $Y_{(n)} = Y^{(n)}$. For infinite $\alpha \in O$ let $Y_{(\alpha)} = Y^{(\alpha+1)}$. Also if α is limit let $Y_{(\alpha-1)} = Y^{(\alpha)}$, whereas for α successor $Y_{(\alpha-1)}$ keeps its obvious meaning. For all $\alpha \geq 1$, a subset of ω is $\Sigma_\alpha^0(Y)$ if and only if it is c.e. in $Y_{(\alpha-1)}$. For all $\alpha \geq 1$, $Y_{(\alpha)}$ is recursively isomorphic to the set of numbers e such that Y belongs to the e^{th} Σ_α^0 set of reals. The isomorphism is uniform in α .

When discussing open and closed sets we run into an annoying fact: there is an open set U which is a Σ_2^0 set of reals, but for which the predicate $[\sigma] \subseteq U$ is not Σ_2^0 . The fact that such a set is Σ_2^0 will not be too helpful for us. For this reason we call an open set Σ_α^0 -open if the set of cylinders contained in it is a Σ_α^0 set of numbers; equivalently, if it is $\Sigma_1^0(\emptyset_{(\alpha-1)})$. The complement of such a set is called Π_α^0 -closed.

The following is the effective version of the fact that all Borel sets have the property of Baire. Recall that for an open set V we let $\partial V = \bar{V} - V$, the boundary of V , be the set-theoretic difference between the closure of V and V itself.

Proposition 2.10. *Suppose that A is a Σ_α^0 set of reals. Then there is a Σ_α^0 -open set U such that the symmetric difference $A \Delta U$ is contained in the union $\bigcup \partial V_n$ where each V_n is a $\Sigma_{<\alpha}^0$ -open set. Indices for U and each V_n can be obtained effectively from an index for A .*

Proof. If the proposition holds for α then for every Π_α^0 set B there is a $\Sigma_{\alpha+1}^0$ -open set W such that the symmetric difference $B \Delta W$ is contained in the union $\bigcup \partial V_n$ where each V_n is Σ_α^0 -open; if U is the open set given for the complement of B then W is the complement of the closure of U , and to the list of sets V_n we add the set U . Once this is known, the proposition follows by induction on α , using the fact that $(\bigcup A_n) \Delta (\bigcup V_n) \subseteq \bigcup (A_n \Delta V_n)$. \square

Let $\alpha \in O$. A real $G \in 2^\omega$ is called $< \alpha$ -Cohen generic if it does not lie on the boundary of any $\Sigma_{< \alpha}^0$ -open set. For example n -genericity is $< (n + 1)$ -genericity and arithmetical genericity is $< \omega$ -genericity. Proposition 2.10 implies that if G is $< \alpha$ -Cohen generic then $G_{(\alpha)}$ is c.e. in $G \oplus \emptyset_{(\alpha-1)}$. This implies that $G_{(\alpha-1)}$ is computable in $G \oplus \emptyset_{(\alpha-1)}$. Unravelling the notation, this means:

- If $n < \omega$ and G is n -generic, then $G^{(n)} \equiv_T G \oplus \emptyset^{(n)}$;
- If $\alpha \geq \omega$ and G is $< \alpha$ -generic, then $G^{(\alpha)} \equiv_T G \oplus \emptyset^{(\alpha)}$.

The equivalence is uniform in α .

In [GM] we show that a Δ_1^1 -Cohen generic set preserves ω_1^{ck} if and only if it is Σ_1^1 -generic. Porism 2.8 implies that if G is Σ_1^1 -generic then a set is $\Pi_1^1(G)$ if and only if it is higher G -c.e.

2.3.2. *Random reals.* It is well-known that if Z is 2-random then Z is generalised low: $Z' \equiv_T Z \oplus \emptyset'$.

The following is the effective version of the fact that all Borel sets are Lebesgue measurable. It is treated in the theses of Kurtz and Kautz (for the arithmetic hierarchy); see [DH10, Thm 6.8.3].

Proposition 2.11. *Let $\alpha \in O$. For any Σ_α^0 set of reals A and positive $q \in \mathbb{Q}$ there are:*

- a Σ_α^0 -open set $U \supseteq A$ such that $\lambda(U - A) \leq q$; and
- a $\Pi_{< \alpha}^0$ -closed set $F \subseteq A$ such that $\lambda(A - F) \leq q$.

An index for U can be obtained effectively from an index for A and from q , using the oracle $\emptyset_{(\alpha-1)}$. An index for F can be obtained effectively from an index for A and from q , using the oracle $\emptyset_{(\alpha)}$. All calculations are uniform in α .

A real is called α -random if it avoids all nested tests $\langle A_n \rangle$ where A_n are uniformly Σ_α^0 sets (not necessarily open). We require that $\lambda(A_n) \leq 2^{-n}$. Proposition 2.11 implies that a real is α -random if and only if it is ML-random relative to $\emptyset_{(\alpha-1)}$. Uniformly in α we have a universal ML-test $\langle U_n^\alpha \rangle$ relative to $\emptyset_{(\alpha-1)}$. An α -randomness deficiency of an α -random real Z is some n such that $Z \notin U_n^\alpha$. If $\langle V_n \rangle$ is any ML-test relative to $\emptyset_{(\alpha-1)}$ (so the sets V_n are uniformly Σ_α^0 -open) then from an α -randomness deficiency of an α -random real Z and an index for the sequence $\langle V_n \rangle$ we can effectively find some m such that $Z \notin V_m$. If $\beta < \alpha$ and Z is α -random then of course it is also β -random, and a β -randomness deficiency of Z can be effectively found from an α -randomness deficiency of Z .

Chong and Yu [CY15] observed that $\Delta_1^1(Z) = \Delta_1^1 \oplus Z$ uniformly for any Δ_1^1 -random real Z which preserves ω_1^{ck} . We prove a more precise version of this result.

Proposition 2.12. *Let $\alpha \geq 2$. If Z is α -random then $Z_{(\alpha-1)} \leq_T Z \oplus \emptyset_{(\alpha-1)}$. An index for the reduction can be found effectively from an α -randomness deficiency of Z . This is uniform in α .*

In short, for all $\alpha \geq 1$, if Z is ML-random relative to $\emptyset^{(\alpha)}$ then $Z^{(\alpha)} \equiv_T Z \oplus \emptyset^{(\alpha)}$. Note the difference at infinite levels compared with Cohen genericity. For example, if G is arithmetically Cohen generic then $G^{(\omega)} \equiv_T G \oplus \emptyset^{(\omega)}$. In contrast, by forcing with arithmetical sets with positive measure one obtains an arithmetically random set Z for which the equation fails.

Proof. We show this in two steps. First we consider successor ordinals α . Suppose that $\alpha = \beta + 1$. We need to show that if Z is $\beta + 1$ -random (ML random relative

to $\mathcal{O}_{(\beta)}$) then $Z_{(\beta)} \leq_T Z \oplus \mathcal{O}_{(\beta)}$. Given a Σ_β^0 set of reals A we want to decide whether $Z \in A$ or not. Using $\mathcal{O}_{(\beta)}$ we find sequences $\langle U_n \rangle$ and $\langle F_n \rangle$ such that U_n is Σ_β^0 -open, F_n is $\Pi_{<\beta}^0$ -closed, $F_n \subseteq A \subseteq U_n$ and $\lambda(U_n - F_n) \leq 2^{-n}$. The sequence $\langle U_n - F_n \rangle$ is a $\mathcal{O}_{(\beta)}$ -ML test, and so we can find some n such that $Z \notin (U_n - F_n)$. Thus $Z \in A$ if and only if $Z \in F_n$. To determine whether $Z \in F_n$ we employ a similar process. F_n is Π_γ^0 -closed for some $\gamma < \beta$. Relativising the case $\alpha = 1$ to $\mathcal{O}_{(\gamma)}$ we obtain a $\mathcal{O}_{(\gamma)}$ -computable sequence $\langle C_m \rangle$ of clopen supersets of F_n such that $\lambda(C_m - F_n) \leq 2^{-m}$. Again this is a $\mathcal{O}_{(\gamma)}$ -test and so we can find some m such that $Z \notin (C_m - F_n)$. We conclude that $Z \in A$ if and only if $Z \in C_m$, and this can of course be checked directly with the oracle Z .

Next we consider limit ordinals α . If Z is α -random (ML-random relative to $\mathcal{O}^{(\alpha)}$) then uniformly in $\gamma <_O \alpha$ it is γ -random (by this we mean that we can, uniformly in γ , compute an upper bound on the γ -randomness deficiency of Z). As $Z_{(\alpha-1)} = Z^{(\alpha)}$ is the effective join $\bigoplus_{\gamma <_O \alpha} Z^{(\gamma)}$, to compute $Z^{(\alpha)}$ it suffices to compute each $Z^{(\gamma)}$, and we may restrict ourselves to successor ordinals γ . However with oracle $\mathcal{O}^{(\alpha)}$ we uniformly obtain $\mathcal{O}^{(\gamma)}$ and we have already shown that $Z^{(\gamma)} \leq_T Z \oplus \mathcal{O}^{(\gamma)}$ uniformly. \square

Remark 2.13. The components of the $\mathcal{O}_{(\beta)}$ -ML tests described in the proof of Proposition 2.12 are all Σ_β^0 rather than $\Sigma_{\beta+1}^0$. These are equivalent to weak β -tests (generalized $\mathcal{O}_{(\beta-1)}$ -ML tests). It would seem that we could relax the randomness requirement. However the key is the uniformity in A : for each A we have a different test, and the full $\beta + 1$ -randomness deficiency of Z is used to find components of these tests that Z avoids. Indeed, Lewis, Montalbán and Nies [LMN07] showed that there is a weakly 2-random set which is not generalized low.

Stern [Ste75] and independently Chong, Nies and Yu [CNY08] showed that a Δ_1^1 -random real is Π_1^1 -random if and only if it preserves ω_1^{ck} . Suppose that Z is Π_1^1 -random. Then it is Π_1^1 -ML random (higher ML-random). From a hyperarithmetic index for a ML-test relative to some hyperarithmetic oracle we can effectively find an index for this test as a sequence of uniformly Π_1^1 open sets. Hence from a randomness deficiency for Z as a higher ML-random real we can uniformly in $\alpha \in O$ find an α -randomness deficiency for Z . Consequently, $\Delta_1^1(Z) = \Delta_1^1 \oplus Z$ uniformly. Hence, if Z is Π_1^1 -random, then a set is $\Pi_1^1(Z)$ if and only if it is higher Z -c.e.

Remark 2.14. Chong and Yu [CY15] proved an analogue of Demuth's theorem: If X is Π_1^1 -random, $Y \leq_h X$ and Y is not hyperarithmetic, then $\text{deg}_h(Y)$ contains a Π_1^1 -random sequence. The structure of their argument follows that of Demuth's theorem; this can be further clarified using Higher Turing reducibility. In the first step we already know that $Y \leq_T X \oplus H$ for some hyperarithmetic set H . Further, being Π_1^1 -random, X is Δ_1^1 -dominated: every $\Delta_1^1(X)$ function is bounded by a hyperarithmetic one. Applying this to the use of the reduction, we see that Y is higher truth-table reducible to X . This implies that Y is higher ML-random for the image measure, and since it is not hyperarithmetic, it is not an atom of this measure. The second step of the proof is now identical to the classical one: if Y is higher ML-random for some hyperarithmetic measure and is not an atom of this measure, then the higher Turing degree of Y contains a higher ML-random sequence. Being Π_1^1 -random, X preserves ω_1^{ck} , and so Y preserves ω_1^{ck} as well, which implies that any higher ML-random sequence in $\text{deg}_{\omega_1^{\text{ck},T}}(Y)$ is in fact Π_1^1 -random.

2.3.3. *Sacks generics.* We consider sets which are generic for forcing with perfect hyperarithmetic closed sets. Sacks (see [Sac90, IV.5]) showed that if G is sufficiently generic for this notion of forcing then G preserves ω_1^{ck} and has minimal hyperdegree. The proof shows that $\Delta_1^1(G) = \Delta_1^1 \oplus G$. However, this is not uniform. We thank Adam Day for pointing this out.

Proposition 2.15. *If G is sufficiently generic for hyperarithmetic Sacks forcing, then O^G is not higher G -c.e.*

Proof. In fact we prove more: we prove that, given a countable collection of enumeration functionals $\langle \Gamma_i \rangle$ (with no assumption on their effectivity), if G is generic enough, then $\Gamma_i^G \neq O^G$ for all i . Consider a given perfect hyperarithmetic closed set, represented by a perfect tree T and an enumeration functional Γ . It is easy to construct a hyperarithmetic set of nodes $D \subseteq T$, open in T , which is dense in T but such that the (hyperarithmetic) tree $T - D$ is perfect. Since D is hyperarithmetic, there exists an n such that for every real X , X has a prefix in D if and only if $n \notin O^X$. If there are no paths X in T such that $n \in \Gamma^X$, then the tree $T - D$, which refines T , forces that $\Gamma^G \neq O^G$. Otherwise there is some $\sigma \in D$ such that $n \in \Gamma^\sigma$. Then the “full subtree” T_σ of nodes in T comparable with σ forces that $\Gamma^G \neq O^G$. \square

3. CONTINUITY AND RANDOMNESS

As discussed in the introduction, when trying to establish analogues of familiar theorems of algorithmic randomness, we sometimes need to work around the usage of time tricks. As a first example we consider van-Lambalgen’s theorem. The proof of one direction: if X is higher ML-random, and Y is higher X -ML-random, then $X \oplus Y$ is higher ML-random — is identical to the analogous “lower” proof. The other direction usually uses a uniform universal ML test, and as discussed in the introduction, no such uniform universal test exists in the higher setting. Given an enumeration operator U , we cannot transform every U^X to an open set with some fixed measure bound. But we show that we can do this for *most* oracles X , and then argue that this suffices.

In the following lemma and below we think of operators enumerating open sets given oracles as open subsets of the plane; if $U \subset (2^\omega)^2$ is open then U^X is the X -section of U .

Lemma 3.1. Let $U \subseteq (2^\omega)^2$ be higher effectively open. For every $\varepsilon > 0$ there is a higher effectively open set $V \subseteq (2^\omega)^2$ such that:

- (1) If $\lambda(U^X) \leq \varepsilon$ then $U^X = V^X$; and
- (2) For all but a set of measure ε -many oracles X , $\lambda(V^X) \leq \varepsilon$.

An index for V can be obtained uniformly from ε and an index for U .

For the proof we use the *projectum function* $p: \omega_1^{\text{ck}} \rightarrow \omega$: this is an ω_1^{ck} -computable injective function.

Proof. We enumerate V . For $s < \omega_1^{\text{ck}}$ we let V_s be the open set enumerated by stage s . Suppose that we see the cylinder $[\sigma, \tau]$ enumerated into U_{s+1} . Let P_s be the set of $X \in [\sigma]$ such that $\lambda(V_s^X \cup [\tau]) > \varepsilon$. We find a clopen set $C_s \subseteq [\sigma]$ which is close to the complement $[\sigma] - P_s$ of P_s inside $[\sigma]$:

- $C_s \cup P_s = [\sigma]$; and

- $\lambda(C_s \cap P_s) \leq \varepsilon \cdot 2^{-p(s)}$.

We then let $V_{s+1} = V_s \cup (C_s \times [\tau])$.

We have $V \subseteq U$, and the desired property (1) holds. To see (2), let $B = \{X \in 2^\omega : \lambda(V^X) > \varepsilon\}$. We claim that $B \subseteq \bigcup_{s < \omega_1^{\text{ck}}} (C_s \cap P_s)$. Let $X \in B$. For limit ordinals $s \leq \omega_1^{\text{ck}}$, $V_s = \bigcup_{t < s} V_t$ (here we let $V_{\omega_1^{\text{ck}}} = V$) and so there is some $s < \omega_1^{\text{ck}}$ such that $\lambda(V_s^X) \leq \varepsilon$ but $\lambda(V_{s+1}^X) > \varepsilon$. But then $X \in C_s \cap P_s$. Now

$$\lambda(B) \leq \lambda\left(\bigcup_{s < \omega_1^{\text{ck}}} P_s \cap C_s\right) \leq \varepsilon \cdot \sum_{s < \omega_1^{\text{ck}}} 2^{-p(s)} \leq \varepsilon$$

as p is injective. □

Remark 3.2. We apply the notational convention used in the previous proof throughout this paper. If X is any object which is approximated or enumerated in ω_1^{ck} many steps then we let $X_{\omega_1^{\text{ck}}} = X$. For example if U is a c.e. open set and $\langle U_s \rangle_{s < \omega_1^{\text{ck}}}$ is an ω_1^{ck} -effective enumeration of U then we write $U_{\omega_1^{\text{ck}}}$ for U ; if $\langle f_s \rangle$ is an ω_1^{ck} -computable approximation of a function f then we let $f_{\omega_1^{\text{ck}}} = f$.

We can now prove a the higher version of van Lambalgen's theorem.

Proof of Theorem 1.4. As discussed above, the proof of one direction has no new ingredients, and so we omit it. In the other direction we are given a pair (X, Y) and assume that Y is not higher X -ML random; and need to show that the pair (X, Y) is not higher ML-random.

Let $\langle U_n^X \rangle$ be a higher X -ML-test which captures Y . By Lemma 3.1 we may assume that for all n , the measure of $B_n = \{Z \in 2^\omega : \lambda(U_n^Z) > 2^{-n}\}$ is at most 2^{-n} ; X is not in any B_n and Y is captured by the X -test after applying the transformation of that lemma. So $(X, Y) \in \bigcap_n U_n$. A calculation (essentially Fubini's theorem) shows that $\lambda(U_n) \leq (1 - 2^{-n}) \cdot 2^{-n} + 2^{-n}$ which converges to 0 (computably). □

3.1. Pulling back strong tests. The argument of Miller and Yu's, sketched in the introduction, relies on the consistency of the given functional. Recall that a *continuous semi-measure* is a function m which assigns to every finite binary string a non-negative real number, such that for all $\sigma \in 2^{<\omega}$, $m(\sigma \hat{0}) + m(\sigma \hat{1}) \leq m(\sigma)$. A continuous semi-measure is *higher c.e.* if the real $m(\sigma)$ is higher left-c.e., uniformly in σ . If Φ is a consistent functional then the function $\sigma \mapsto \lambda(\Phi^{-1}[\sigma])$ is a continuous semi-measure. In the higher setting not all functionals can be made continuous. However as above, given a functional Ψ and some $\varepsilon > 0$ we can transform Ψ to a functional Φ such that $\Phi(X) = \Psi(X)$ if $\Psi(X)$ is consistent, and such that $\Phi(X)$ is inconsistent for at most ε -many (in the sense of measure) oracles. In fact we can combine all the ε -modifications in one to get the following.

Lemma 3.3. For every higher Turing functional Ψ there is a higher Turing functional Φ such that:

- (1) for all X for which $\Psi(X)$ is consistent, $\Phi(X) = \Psi(X)$; and
- (2) the function $\tau \mapsto \lambda(\Phi^{-1}[\tau])$ is bounded by a higher c.e. continuous semi-measure.

Proof. Fix a function $q: 2^{<\omega} \rightarrow \mathbb{Q}^+$ such that $\sum_{\tau \in 2^{<\omega}} q(\tau) \leq 1$ and such that $\tau_1 \leq \tau_2$ implies $q(\tau_1) \geq q(\tau_2)$ (for example let $q(\tau) = 2^{-3|\tau|}$). We enumerate a functional Φ .

Suppose that we see the axiom (σ_s, τ_s) enumerated into Ψ_{s+1} . We let P_s be the set of $X \in [\sigma_s]$ such that $\Phi_s(X)$ is inconsistent with τ_s . Let C_s be a clopen subset of $[\sigma_s]$ close to the complement $[\sigma_s] - P_s$; we mean that $P_s \cup C_s = [\sigma_s]$ and $\lambda(P_s \cap C_s) \leq 2^{-p(s)} \cdot q(\tau_s)$, where as above p is the projection function. We then declare that $\Phi_{s+1}(X) \geq \tau_s$ for all $X \in C_s$.

Inductively, for all s and X , $\Phi_s(X) \leq \Psi_s(X)$, and so if $X \in P_s$ then $\Psi(X)$ is inconsistent. This establishes (1).

For (2) we let

$$m(\tau) = \lambda(\Phi^{-1}[\tau]) + \sum_{\rho \succ \tau} q(\rho).$$

For $\tau \in 2^{<\omega}$ let $B(\tau) = \Phi^{-1}[\tau \hat{\ } 0] \cap \Phi^{-1}[\tau \hat{\ } 1]$. So

$$\lambda(\Phi^{-1}[\tau \hat{\ } 0]) + \lambda(\Phi^{-1}[\tau \hat{\ } 1]) \leq \lambda(\Phi^{-1}[\tau]) + \lambda(B(\tau)).$$

If $X \in B(\tau)$ then there is some stage $s < \omega_1^{\text{ck}}$ such that $X \in P_s \cap C_s$ and τ_s extends either $\tau \hat{\ } 0$ or $\tau \hat{\ } 1$. Since $q(\tau) \geq q(\tau_s)$, the argument of Lemma 3.1 shows that $\lambda(B(\tau)) \leq q(\tau)$. A calculation now shows that m is a continuous semi-measure. \square

Lemma 3.3 allows us to show that strong randomness notions are downwards closed in the ω_1^{ck} -degrees of higher ML-random sets. In particular we get Theorem 1.2.

Theorem 3.4. *Suppose that X and Y are higher ML-random and that $X \leq_{\omega_1^{\text{ck}}} Y$. If Y is higher weakly-2-random (higher difference random, higher Z-ML-random for some $Z \in 2^\omega, \dots$) then so is X .*

Proof. By Lemma 3.3 we get a higher Turing functional Φ such that $\Phi(Y) = X$ and $\lambda(\Phi^{-1}[\tau]) \leq m(\tau)$ for some higher c.e., continuous semi-measure. Since X is higher ML-random, $m(X \upharpoonright_n) \leq c \cdot 2^{-n}$ for some constant c . We can then enumerate a functional $\Psi \subseteq \Phi$ such that $\Psi(Y) = X$ and $\lambda(\Psi^{-1}[\tau]) \leq c \cdot 2^{-|\tau|}$ for all τ : we enumerate Ψ . At stage s say an axiom (σ, τ) appears in Φ_{s+1} . If $\lambda([\sigma] \cup \Psi_s^{-1}[\rho]) > c \cdot 2^{-|\rho|}$ for some $\rho \leq \tau$ then we let $\Psi_{s+1} = \Psi_s$; otherwise we let $\Psi_{s+1} = \Psi_s \cup \{(\sigma, \tau)\}$. In the first case $\lambda(\Phi^{-1}[\rho]) > c \cdot 2^{-|\rho|}$ and so ρ is not an initial segment of X ; so σ is not an initial segment of Y .

If $\langle U_n \rangle$ is any strong test capturing X then $\langle \Phi^{-1}[U_n] \rangle$ is a strong test capturing Y . The point is that $\lambda(\Phi^{-1}[U_n]) \leq c \cdot \lambda(U_n)$. There may not be any higher c.e. (higher Z-c.e.) antichain generating U_n ; but for the measure calculation we do not need effectiveness: the inequality is obtained by considering the antichain of minimal strings (maximal intervals) in U_n . \square

4. K-TRIVIALITY

Hjorth and Nies defined in [HN07] the notion of higher prefix-free Kolmogorov complexity, based on the concept of universal Π_1^1 prefix-free machine. We denote this complexity function by K , as we will not be using the traditional “lower” complexity. Armed with this concept Hjorth and Nies defined the class of higher K -trivial sets, those sets $A \in 2^\omega$ satisfying $K(A \upharpoonright_n) \leq^+ K(n)$.

Hjorth and Nies proved that there are higher K -trivial sets which are not hyperarithmetical (arguing that Solovay's proof applies in the higher setting) and also that every higher K -trivial is Turing reducible to Kleene's O . As described in the introduction, since they use discontinuous relativisations, their notions of higher lowness for K , higher bases for randomness and higher lowness for MLR coincide with being hyperarithmetical. Continuous relativisations yield analogues of familiar equivalences.

In addition to Theorem 1.5, we also show that a set is higher K -trivial if and only if it is higher low for K . As mentioned above, defining the notion is not completely straightforward because there are oracles A for which there is no optimal prefix-free complexity; so K^A is not well-defined for all A . Further complication is due to the potential failure of the equivalence between prefix-free complexity and discrete c.e. measures. Recall that a *discrete measure* (often called a discrete semi-measure, but it is a measure) is simply a measure on ω (equivalently, on any computable set); such a measure is of course determined by the measures of its atoms. A discrete measure μ is called (higher) c.e. if $\mu(n)$ is a (higher) left-c.e. real, uniformly in n . Nies and Hjorth showed that the higher analogue of the Kraft-Chaitin theorem holds, from which follows the higher analogue of the coding theorem, which says that every higher c.e. discrete measure can be realised as the measure derived from a higher prefix-free machine ($\mu_M = 2^{-K_M}$). Thus 2^{-K} is an optimal higher c.e. discrete measure.

We do not know whether the coding theorem can be continuously relativised to every oracle. Thus given an oracle A we can investigate both higher A -computable prefix-free machines (their graphs are higher A -c.e.) and their associated complexities; and higher A -c.e. discrete measures. This gives two definitions of lowness:

- an oracle A is *low for higher K* if for every higher A -computable prefix-free machine M , $K \leq^+ K_M$;
- an oracle A is *low for higher c.e. discrete measures* if for every higher A -c.e. discrete measure ν , $\mu \geq^{\times} \nu$ where μ is the optimal higher c.e. discrete measure.

A-priori the second notion is stronger. We will show that both of these concepts coincides with higher K -triviality. On the other hand, since the concept of K -triviality itself does not involve relativisation, it can be characterised using discrete measures: a set A is K -trivial if and only if $\mu(A \upharpoonright_n) \geq^{\times} \mu(n)$.

4.1. Approximations of K -trivial sets. The following is implicit in [HN07].

Proposition 4.1. *Every nonhyperarithmetical higher K -trivial set has a collapsing approximation.*

In fact if A is higher K -trivial then there is an increasing approximation $\langle \mu_s \rangle$ of μ and a collapsing approximation $\langle A_s \rangle$ of A such that for some constant $\delta > 0$, $\mu_s(A \upharpoonright_n) \geq \delta \cdot \mu_s(n)$ for all $n < \omega$ and all $s < \omega_1^{\text{ck}}$.

Proof. We start with an arbitrary enumeration $\langle \mathbb{U}_s \rangle$ of the universal higher-c.e. prefix-free machine \mathbb{U} , and let $K_s = K_{\mathbb{U}_s}$. As usual we assume that the enumeration of \mathbb{U} is continuous, i.e. $\mathbb{U}_s = \bigcup_{t < s} \mathbb{U}_t$ for every limit ordinal $s \leq \omega_1^{\text{ck}}$. Hence $K_s = \lim_{t \rightarrow s} K_t$ for every limit ordinal s .

There is an ω_1^{ck} -computable sequence of trees $\langle T_s \rangle_{s \leq \omega_1^{\text{ck}}}$ such that:

- For all limit $s \leq \omega_1^{\text{ck}}$, $T_s = \lim_{t \rightarrow s} T_t$; and

- A is the unique path of $T_{\omega_1^{\text{ck}}}$.

For let b be a K -triviality constant for A . There are only finitely many K -trivial sequences with constant b . For $s \leq \omega_1^{\text{ck}}$ let S_s be the tree of finite binary strings which are K_s -trivial with constant b . Let σ be a string on $S_{\omega_1^{\text{ck}}}$ which isolates A on $S_{\omega_1^{\text{ck}}}$. We let T_s be the restriction of S_s to strings comparable with σ .

In [HN07], Hjorth and Nies show that there is an ω_1^{ck} -computable closed and unbounded set $C \subseteq \omega_1^{\text{ck}}$ such that for all $s \in C$, the tree T_s has only finitely many paths. A similar argument shows that after thinning to a possibly smaller set of stages we may assume that for all $s \in C$, T_s has a path (for all n , if T_t contains a string of length n for all t in some set B of stages, then by continuity $T_{\sup B}$ also contains a string of length n .) We define the approximation $\langle A_s \rangle$ for $s \in C$ by letting A_s be the leftmost path in T_s . Then $A = \lim_{s \in C} A_s$. This approximation is collapsing: if $A \upharpoonright_n \in T_{s(n)}$ and $s(\omega) = \sup_n s(n)$ then A is a path in $T_{s(\omega)}$; if $s(\omega) < \omega_1^{\text{ck}}$ then $T_{s(\omega)}$ is hyperarithmetic, and so each of its finitely many paths is hyperarithmetic.

Finally we renumber our approximations using the increasing ω_1^{ck} -computable enumeration of C , and let $\mu_s = 2^{-K_s}$. \square

The fact that a set A has a collapsing approximation allows us to relativise to A many familiar techniques, with arguments along the lines of that of Proposition 2.2. In the language of [BGHM], it is a “good oracle”. For example:

Lemma 4.2. Suppose that A has a collapsing approximation. Then there is an optimal higher A -c.e. discrete measure μ^A , and a sequence Z is higher A -ML-random if and only if $\mu^A(Z \upharpoonright_n) \leq 2^{-n}$. Further, there is a universal higher A -c.e. prefix-free machine \mathbb{U}^A and $\mu^A = \times 2^{-K^A}$.

Proof. To get a universal higher A -c.e. prefix-free machine we show that we can uniformly transform a given enumeration functional W to an enumeration functional V such that V^A is the graph of a function with prefix-free domain (indeed this is true for every oracle), and if W^A is a graph of such a function then $V^A = W^A$. As in the proof of Proposition 2.2, if we see that $\sigma = A_s \upharpoonright_n$ is not an initial segment of A_t for any $t < s$, and W_s^τ is the graph of a function with prefix-free domain, then we let $V_s^\tau = W_s^\tau$.

In the same way we get μ^A ; if $\mu \subseteq 2^{<\omega} \times \omega \times \mathbb{Q}^+$ then we let, for each $n < \omega$ and $X \in 2^{<\omega}$, $\mu^X(n) = \sup \{q \in \mathbb{Q}^+ : (\sigma, n, q) \in \mu \text{ for some } \sigma \leq X\}$; and let $\mu^X(\omega) = \sum_{n \in \omega} \mu^X(n)$. We can transform each higher c.e. such μ into some ν such that $\nu^A(\omega) \leq 1$ and if $\mu^A(\omega) \leq 1$ then $\nu^A = \mu^A$: when we see a “fresh” $\tau < A_s$, we copy μ_s^τ , provided that $\mu_s^\tau(\omega) \leq 1$.

The key step in the standard (“lower”) proof of the Levin-Schnorr theorem (the equivalence of discrete measures and tests in capturing ML-randomness) is taking an effectively open set U and obtaining a c.e. prefix-free set generating U . In the higher setting this is impossible; using the projectum function and approximations of closed sets from above by clopen sets, we can get a set of strings generating U whose *weight* is bounded by $\lambda(U) + \varepsilon$ for any prescribed $\varepsilon > 0$. However working relative to an oracle A with a collapsing approximation makes the situation *easier*: in some sense the collapsing approximation brings us closer to ω -computability. If A has a collapsing approximation and U^A is higher A -effectively open then there is a higher A -c.e. prefix-free set of strings W^A generating U : if $\tau < A_s$ is fresh then

we enumerate into W_{s+1}^τ all strings σ of length $|\tau|$ such that $[\sigma] \subseteq U_s^A$ but $[\sigma]$ is disjoint from $[W_s^\tau]$.

In a similar way, relative to A we can follow the standard proof of the Kraft-Chaitin / coding theorem without having to resort to the necessary complications of the proof of the unrelativised theorem in the higher setting (see [HN07]).⁵ \square

Suppose that A is low for higher K . Then it is higher K -trivial. With Lemma 4.2 we can then conclude that it is also low for higher c.e., discrete measures, and low for higher ML-randomness.

4.2. Hungry sets. We next show that if A is a base for higher randomness ($A \leq_{\omega_1^{\text{ckT}}} Z$ for some higher A -ML-random set Z) then A is higher K -trivial.

- We could modify the argument to obtain lowness for higher K . We will later show though that higher K -triviality implies lowness for higher K .
- The higher version of the Kučera-Gács theorem shows that if A is low for higher ML-randomness then it is a base for higher randomness. So we also conclude that lowness for higher ML-randomness implies higher K -triviality and therefore lowness for higher K . A more direct argument is likely possible but for brevity we omit it.

We need to carry out the “hungry sets” construction of [HNS07]. In [HN07] the authors claim that the proof carries over with only notational changes; they ignore the typical topological problems. These problems are present even if one assumes that the reduction of A to Z is a fin-h reduction; the problems increase slightly when inconsistent functionals are admitted. Here we discuss these problems and show how to overcome them.

We recall the structure of the proof. Suppose that $\Phi(Z) = A$ where Φ is a higher Turing functional and Z is higher A -ML-random. We fix $\varepsilon > 0$. We enumerate “hungry sets” $C^\alpha = C^\alpha(\varepsilon)$ for every finite binary string α ; we ensure that $C^\alpha \subseteq \Phi^{-1}[\alpha]$. An attempt to show that A is K -trivial is made by ensuring that $\alpha \mapsto \lambda(C^\alpha)$ is a higher-c.e. discrete measure, and attempting to show that $\lambda(C^{\uparrow n}) \geq \mu(n)$. So we aim to ensure three things:

- (1) the measure of $\bigcup_{\alpha < A} C^\alpha$ is bounded by ε ;
- (2) either for all $\alpha < A$, $\lambda(C^\alpha) = \varepsilon \mu(|\alpha|)$, or $Z \in \bigcup_{\alpha < A} C^\alpha$; and
- (3) the sum $\sum_{\alpha \in 2^{<\omega}} \lambda(C^\alpha)$ is finite.

We ensure that for all s and α , $\lambda(C_s^\alpha) \leq \varepsilon \mu_s(|\alpha|)$; this ensures (1). In the standard proof, (3) is obtained by ensuring that the hungry sets are pairwise disjoint. The usual topological reasons preclude this from happening in the higher setting; at an infinite stage s , $\Phi_s^{-1}[\alpha] - C_s^\alpha$ may have positive measure but no interior. Further, if Φ is inconsistent then we do not automatically get that C^α and C^β are

⁵Another way to understand the situation is to observe that a collapsing approximation of A gives us an ω_1^{ck} - A -computable ω -sequence $\langle \alpha_n \rangle$ cofinal in ω_1^{ck} . From this we get a relation $E \leq_{\omega_1^{\text{ckT}}} A$ such that $(\omega, E) \cong (L_{\omega_1^{\text{ck}}}, \varepsilon)$ (and as is the situation with O , we can make the map $n \mapsto n^{(\omega, E)}$ computable). This means that the higher A -c.e. sets are precisely those which are Σ_1 -definable in the structure $(L_{\omega_1^{\text{ck}}}, \varepsilon, A)$. So when designing higher A -c.e. sets we don't have to consider other oracles, as is usually the case with designing oracle-c.e. sets; and we can enumerate such sets using a recursion of length ω along the sequence $\langle \alpha_n \rangle$. All familiar constructions can be performed this way. For example when enumerating a higher A -effectively open set U we may assume that by stage α_n , only strings of length n have been enumerated into U .

disjoint if α and β are incomparable. As above, we remedy this by allowing overlap, but ensuring that it is small.

Fix positive rational numbers δ_α for all strings $\alpha \in 2^{<\omega}$, so that $\sum_{\alpha \in 2^{<\omega}} \delta_\alpha$ is finite. For notational simplicity at each stage of the construction we consider a single string α (at stage $t + n$, t limit, consider the n^{th} finite binary string). Let $C_s = \bigcup_{\beta \in 2^{<\omega}} C_s^\beta$. We find a clopen $B_s \subseteq C_s$ such that $\lambda(C_s - B_s) \leq \delta_\alpha 2^{-p(s)}$. We now consider:

- $G_s = \Phi_s^{-1}[\alpha] - B_s$ — this is potential fodder;
- $q_s = \varepsilon \mu_s(|\alpha|) - \lambda(C_s^\alpha)$ — this is the amount we would like to add to C^α .

If $\lambda(G_s) \leq q_s$ then we let $C_{s+1}^\alpha = C_s^\alpha \cup G_s$. If $\lambda(G_s) > q_s$ we find some hyperarithmetic open set $U_s \subseteq G_s$ of measure exactly q_s and let $C_{s+1}^\alpha = C_s^\alpha \cup U_s$. It is easy to check that the bound $\lambda(C^\alpha) \leq \varepsilon \mu(|\alpha|)$ is maintained at stage $s + 1$; that if $\lambda(G_s) > q_s$ then $\varepsilon \mu_s(|\alpha|) - \lambda(C_{s+1}^\alpha) \leq \delta_\alpha 2^{-p(s)}$; and that $\lambda(E_s^\alpha) \leq \delta_\alpha 2^{-p(s)}$, where $E_s^\alpha = (C_{s+1}^\alpha - C_s^\alpha) \cap C_s$.

Suppose that $Z \notin \bigcup_{\beta < A} C^\beta$; since Φ is consistent on Z and $C^\beta \subseteq \Phi^{-1}[\beta]$, $Z \notin C = \bigcup_{\beta \in 2^{<\omega}} C^\beta$. Let $\alpha < A$, and suppose for a contradiction that $\lambda(C^\alpha) < \varepsilon \mu(|\alpha|)$; let $\lambda(C^\alpha) < r < q < \varepsilon \mu(|\alpha|)$ be rational numbers. For all but a bounded set of stages s we have $\varepsilon \mu_s(|\alpha|) > q$, $\lambda(C_s^\alpha) < r$, and $\delta_\alpha 2^{-p(s)} < q - r$. Suppose that s is a late stage at which α is considered; so $Z \in \Phi^{-1}[\alpha]$. The fact that $Z \notin C$ implies that $\lambda(G_s) > q_s$, but then enough measure is added to C_{s+1}^α to bring it to within $\delta_\alpha 2^{-p(s)}$ of $\varepsilon \mu_s(|\alpha|)$; this is a contradiction, which yields (2).

It remains to verify (3). For each α let $E^\alpha = \bigcup_s E_s^\alpha$; so $\lambda(E^\alpha) \leq \delta_\alpha$. The sets $C^\alpha - E^\alpha$ are pairwise disjoint: a real $X \in C^\alpha - E^\alpha$ enters C^α before it enters any other C^β . Hence

$$\sum_{\alpha \in 2^{<\omega}} \lambda(C^\alpha) = \sum_{\alpha \in 2^{<\omega}} \lambda(C^\alpha - E^\alpha) + \sum_{\alpha \in 2^{<\omega}} \lambda(E^\alpha) \leq 1 + \sum_{\alpha \in 2^{<\omega}} \delta_\alpha$$

which is finite.

4.3. The main lemma. Unlike the hungry sets construction, there are no major topological complications associated with the golden run argument. The proof translated to the higher setting without many modifications. Proposition 4.1 gives a useful approximation with which to run the construction. In the standard construction we assume that the given enumeration is first sped-up so that at every stage s , $A_s \upharpoonright_s$ is K_s -trivial; here we can assume that A_s in its entirety is K_s -trivial. When drip-feeding measure we are instructed to put some weight on a fresh number n , and this usually means larger than any number chosen so far. This of course we cannot do. However we can choose a number as large as necessary (larger than the length of some initial segment of A which we are trying to certify) without needing to re-use followers; at stage s we choose from the $p(s)^{\text{th}}$ column of ω .

This allows us to prove the higher version of the main lemma [Nie09, Lemma 5.5.1]. Suppose that $\langle A_s \rangle_{s < \omega_1^{\text{ck}}}$ is an ω_1^{ck} -computable approximation of a set A . For $s < \omega_1^{\text{ck}}$ let $A_s \wedge A_{s+1}$ be the longest common initial segment of A_s and A_{s+1} . Let μ^A be a higher A -c.e. discrete measure. If $\langle A_s \rangle$ is a collapsing approximation then we may assume that we have an enumeration $\langle \mu_s \rangle$ of μ such that for all $s < \omega_1^{\text{ck}}$, $\mu_s^{A_s}$ is a higher c.e. discrete measure as well (in fact as discussed above we may assume that μ^X is a discrete measure for all oracles X). Recall that for a

discrete measure ν we let $\nu(\omega) = \sum_n \nu(n)$. The quantity

$$\mu_s^{A_s}(\omega) - \mu_s^{A_s \wedge A_{s+1}}(\omega)$$

is the total mass assigned by $\mu_s^{A_s}$ which was believed at stage s but thought to be incorrect at stage $s + 1$.

Proposition 4.3. *Let A be higher K -trivial, and suppose that μ^A is a higher A -c.e. discrete measure. Then there is an approximation $\langle A_s \rangle$ of A such that the sum*

$$\sum_{s < \omega_1^{\text{ck}}} (\mu_s^{A_s}(\omega) - \mu_s^{A_s \wedge A_{s+1}}(\omega))$$

is finite.

Further, we may assume that if $\langle A_s \rangle$ is a given collapsing approximation of A and $\langle \mu_s \rangle$ is an enumeration of μ such that for all s , $\mu_s^{A_s}(\omega) \leq 1$, then there is an ω_1^{ck} -computable closed unbounded set $C \subseteq \omega_1^{\text{ck}}$ such that

$$\sum_{s \in C} (\mu_s^{A_s}(\omega) - \mu_s^{A_s \wedge A_{s+1}}(\omega)) < \infty,$$

where $s^+ = \min(C - (s + 1))$.

We obtain familiar corollaries:

- Every higher K -trivial set is low for higher K ; this completes the proof of Theorem 1.5.
- Every higher K -trivial set is higher Turing reducible to a higher c.e., higher K -trivial set.
- Every higher K -trivial set is higher ω -c.a.

5. HIGHER WEAK 2-RANDOMNESS

Recall that a higher weak 2-test (a generalised higher ML test) is a sequence $\langle U_n \rangle$ of uniformly Π_1^1 open sets (higher c.e. open sets) whose intersection is null. Note that we can suppose that the U_n are nested, i.e., $U_{n+1} \subseteq U_n$ for all n (indeed, if they are not, one can consider $V_n = \bigcap_{k \leq n} U_k$ and observe that the V_n are nested and that their intersection is the same as $\bigcap_n U_n$).

A sequence is higher weak 2-random if it avoids all higher weak 2-tests. In this section we find alternative, Demuth-like characterisations of higher weak 2-randomness; we consider their Borel rank through an effective lens; and we investigate the interaction with classes of higher Δ_2^0 sequences. These considerations will culminate in a separation of Π_1^1 randomness from higher weak 2-randomness.

5.1. Compact approximations and higher weak 2-randomness. Definition 1.10 describes compact approximations. We recall the notational convention discussed in Remark 3.2: if $\langle f_s \rangle_{s < \omega_1^{\text{ck}}}$ is an ω_1^{ck} -computable approximation of a function f then we write $f_{\omega_1^{\text{ck}}}$ for f .

Proposition 5.1. *No sequence $X \in 2^\omega$ with a higher closed approximation is higher weakly 2-random.*

Proof. Let $\langle X_s \rangle_{s < \omega_1^{\text{ck}}}$ be a closed approximation of $X = X_{\omega_1^{\text{ck}}}$. Let $C = \{X_s : s \leq \omega_1^{\text{ck}}\}$. We let $U_n = \bigcup_{s < \omega_1^{\text{ck}}} [X_s \upharpoonright_n]$. The sequence $\langle U_n \rangle$ is uniformly higher effectively open. Certainly $X \in \bigcap_n U_n$. If $Y \in U_n$ then the distance of Y from C is

at most 2^{-n} . Hence if $Y \in \bigcap_n U_n$ then the distance of Y from C is 0. Since C is closed, this implies that $\bigcap_n U_n \subseteq C$.

The set C is countable, and so null. This shows that $\bigcap_n U_n$ is null, and so is a higher weak 2-test. \square

Even if $\langle X_s \rangle$ is a higher left-c.e. approximation, we do not know how to directly show that the measure of the sets U_n tends to 0.

A generalisation of Proposition 5.1 gives a Demuth-style characterisation of higher weak 2-randomness, a weakening of the class higher $\text{MLR}[O]$ (introduced later in Section 7). In the lower setting of course weak 2-randomness is equivalent to $\text{MLR}[\emptyset]$. Recall that we let W_e denote the e^{th} higher c.e. open set.

Proposition 5.2. *The following classes of tests precisely capture higher weak 2-tests.*

- (1) *Nested tests of the form $\langle W_{f(n)} \rangle$ where $\lambda(W_{f(n)}) \leq 2^{-n}$ and f has a finite-change approximation.*
- (2) *Nested tests of the form $\langle W_{f(n)} \rangle$ where $\lambda(W_{f(n)}) \leq 2^{-n}$ and f has a compact approximation.*

Proof. Every function which has a finite-change approximation also has a compact approximation (Lemma 1.14). So we need to show that:

- (a) Every weak 2-test can be covered by a test with a finite-change index function (as in (1)).
- (b) Every test with a compact index function (as in (2)) can be covered by a weak 2-test.

For (a), let $\langle U_n \rangle$ be a higher weak 2-test; let $U_{n,s}$ be a uniform enumeration of U_n . For $s \leq \omega_1^{\text{ck}}$ let $f_s(k)$ be the least n such that $\lambda(U_{n,s}) \leq 2^{-k}$. Since the measures of $U_{n,s}$ are non-decreasing, the functions $f_s(k)$ are non-decreasing in s , and converge to a limit since for all k there is an n such that $\lambda(U_n) < 2^{-k}$. So $\langle f_s \rangle$ is a finite-change approximation of $f = f_{\omega_1^{\text{ck}}}$. Passing to canonical indices we get a test with a finite-change index function which covers the test $\langle U_n \rangle$.

For (b), the argument is inspired by that of Proposition 5.1. Let $\langle f_s \rangle_{s < \omega_1^{\text{ck}}}$ be a compact approximation of a function f such that $\lambda(W_{f(n)}) \leq 2^{-n}$ and $\langle W_{f(n)} \rangle$ is nested.

A priori, the sets $W_{f_s(n)}$ (for a fixed s) may not be nested. We replace $W_{f_s(n)}$ by $\bigcap_{m \leq n} W_{f_s(m)}$. This changes the index function. However the first n values of the new index function g_s are determined by the first n bits of f_s . In particular, the map $f_s \mapsto g_s$ is continuous, and hence the set $\{g_s : s \leq \omega_1^{\text{ck}}\}$ is compact (and of course $g = \lim_s g_s$). Thus, without loss of generality, we may assume that each test $\langle W_{f_s(n)} \rangle$ is nested. We may also assume that $\lambda(W_{f_s(n)}) \leq 2^{-n}$ for all s and n .

Let $U_n = \bigcup_{s < \omega_1^{\text{ck}}} W_{f_s(n)}$. Since $U_n \supseteq W_{f(n)}$, the test $\langle U_n \rangle$ covers the given test $\langle W_{f(n)} \rangle$; and the sets U_n are uniformly Π_1^1 open. We show that $\bigcap_n U_n$ is null.

For each $s \leq \omega_1^{\text{ck}}$, let $A_s = \bigcap_n W_{f_s(n)}$, and let $A = \bigcup_{s \leq \omega_1^{\text{ck}}} A_s$. Each A_s is null; since $\omega_1^{\text{ck}} + 1$ is countable, A is null. We show that $\bigcap_n U_n \subseteq A$. For let $Y \in \bigcap_n U_n$. For each n there is some $s(n)$ such that $Y \in W_{f_{s(n)}(n)}$. Since the set $\{f_t : t \leq \omega_1^{\text{ck}}\}$ is compact, the set $\{f_{s(n)} : n < \omega\}$ has a limit point, and that limit point equals f_t for some $t \leq \omega_1^{\text{ck}}$. Then $Y \in A_t$: to see this, let $n < \omega$. There is some $k > n$ such that $f_t \upharpoonright_{n+1} = f_{s(k)} \upharpoonright_{n+1}$. Then $Y \in W_{f_{s(k)}(k)} \subseteq W_{f_{s(k)}(n)} = W_{f_t(n)}$ as required. \square

5.2. A short proof of a theorem of Chong and Yu's. Chong and Yu [CY15] showed that every hyperdegree above that of Kleene's O contains a higher ML-random set which is not higher weak 2-random. The above results give us a short proof of this fact. Let $Y \geq_h O$. There is some $X \equiv_h Y$ such that $X \geq_{\omega_1^{\text{ck}}\text{T}} O$, for example $X = Y \oplus O$. By the higher Kučera-Gács theorem there is some $Z \equiv_{\omega_1^{\text{ck}}\text{T}} X$ which is higher ML-random. Since $Z \geq_{\omega_1^{\text{ck}}\text{T}} \Omega$ and higher Ω is not higher weak 2-random, neither is Z (Theorem 1.2). And $Z \equiv_h Y$.

5.3. The effective Borel rank of higher weak 2-randomness. Every higher null weak 2-set is G_δ , and so the set of higher weak 2-random sequences is Π_3^0 . Yu showed that this is sharp. There is a natural higher lightface version of the Borel hierarchy. For example a set is higher Π_2^0 if it is the uniform intersection of Π_1^1 open sets (so the higher null weak 2-sets are precisely the null higher Π_2^0 sets). A set is higher Σ_3^0 if it is the uniform union of higher Π_2^0 sets, and so on. We investigate this hierarchy in detail in [GM]. Here we show that the set of higher weakly 2-random sequences is not higher Π_3^0 . Thus, picking out the null higher Π_2^0 sets requires an oracle. This follows from Proposition 5.1 and Lemma 1.14 and the following proposition.

Proposition 5.3. *Every higher Π_3^0 set of measure 1 contains a sequence which has a finite-change approximation.*

Proof. Let F be a higher Π_3^0 set of measure 1. So $F = \bigcap_{e < \omega} F^e$, where F^e are uniformly higher Σ_2^0 , and since $F \subseteq F^e$, each F^e has measure 1. We write $F^e = \bigcup_k F^{e,k}$ where $\langle F^{e,k} \rangle_{k < \omega}$ is an increasing sequence of uniformly higher effectively closed sets, namely, Σ_1^1 closed sets.

We define a real $x \in F$ by recursion on $e < \omega$. To ensure that $x \in F$ we will, for each e , pick one of the closed sets $F^{e,k}$ and ensure that $x \in F^{e,k}$. We denote the index k chosen by $c(e)$. We define $x \upharpoonright_e$ and $c \upharpoonright_e$ by simultaneous recursion. At step $e < \omega$, given $x \upharpoonright_e$ and $c \upharpoonright_e$, let $H^e = \bigcap_{d < e} F^{d, c(d)}$. For $e = 0$ we have $H^e = 2^\omega$. Inductively we ensure that $\lambda(H^e \upharpoonright x \upharpoonright_e) \geq 2^{-e}$. We then choose:

- $x(e) \in \{0, 1\}$ to be the least so that $\lambda(H^e \upharpoonright x \upharpoonright_{e+1}) \geq 2^{-e}$.
- Since F^e has measure 1, $\lambda(H^e \cap F^e \upharpoonright x \upharpoonright_{e+1}) \geq 2^{-e}$, and so there is some $k < \omega$ such that $\lambda(H^e \cap F^{e,k} \upharpoonright x \upharpoonright_{e+1}) \geq 2^{-(e+1)}$. We let $c(e)$ be the least such k .

For all $e < \omega$ and $d \geq e$, $H^e \cap [x \upharpoonright_d] \supseteq H^d \cap [x \upharpoonright_d]$ are not null and so nonempty. Since H^e is closed, $x \in H^e$. And $H^e \subseteq F^d$ for all $d < e$, and so $\bigcap H_e \subseteq F$. Thus $x \in F$.

It remains to show that x has a finite-change approximation. To do so, we approximate the set F and the sets it is built up from. The sets $F^{e,k}$ have (uniform) co-enumerations $F_s^{e,k}$ for $s < \omega_1^{\text{ck}}$, each $F_s^{e,k}$ is hyperarithmetic and if $s < t$ then $F_s^{e,k} \supseteq F_t^{e,k}$. We also assume that these co-enumerations are continuous: for limit $s < \omega_1^{\text{ck}}$, $F_s^{e,k} = \bigcap_{t < s} F_t^{e,k}$. We let $F_s^e = \bigcup_k F_s^{e,k}$. We then repeat the construction above at each stage $s < \omega_1^{\text{ck}}$: we define $x_s \in 2^\omega$ and $c_s \in \omega^\omega$ coding choices of indices so that letting $H_s^e = \bigcap_{d < e} F_s^d$ we have:

- (1) $\lambda(H_s^e \upharpoonright x_s \upharpoonright_e) \geq 2^{-e}$;
- (2) $x_s(e)$ is least such that $\lambda(H_s^e \upharpoonright x_s \upharpoonright_{e+1}) \geq 2^{-e}$; and
- (3) $c_s(e)$ is the least k such that $\lambda(H_s^e \cap F_s^{e,k} \upharpoonright x_s \upharpoonright_{e+1}) \geq 2^{-(e+1)}$.

We will show that $\langle x_s \rangle$ is a finite-change approximation of x . To begin, we note that if $e < \omega$, $s < t \leq \omega_1^{\text{ck}}$ and $c_s \upharpoonright_e = c_t \upharpoonright_e$ then $H_s^e \supseteq H_t^e$. This implies the following:

- (*) Suppose that $c_s \upharpoonright_e = c_t \upharpoonright_e$ and $x_s \upharpoonright_e = x_t \upharpoonright_e$. Then $x_s(e) \leq x_t(e)$.
- (**) Suppose that $c_s \upharpoonright_e = c_t \upharpoonright_e$ and $x_s \upharpoonright_{e+1} = x_t \upharpoonright_{e+1}$. Then $c_s(e) \leq c_t(e)$.

The following claim shows that we cannot cycle through infinitely many values of $c_r(e)$ while $c_r \upharpoonright_e$ remains stable. We use the following notation. If $I \subseteq \omega_1^{\text{ck}}$ is an interval of stages and $x_r \upharpoonright_e$ is constant for all $r \in I$, then we denote this constant value by $x_I \upharpoonright_e$; similarly for c , or $x(e)$, etc.

Claim 5.3.1. Let $e < \omega$. Let $I \subseteq \omega_1^{\text{ck}}$ be an interval of stages on which $c_r \upharpoonright_e$ and $x_r \upharpoonright_e$ are constant. Then $c_{\sup I} \upharpoonright_e = c_I \upharpoonright_e$ and $x_{\sup I} \upharpoonright_e = x_I \upharpoonright_e$.

Proof. By induction on e . Assume we know this for e . Let $s = \sup I$. We assume that $c_r \upharpoonright_{e+1}$ and $x_r \upharpoonright_{e+1}$ are constant on I ; we need to show that $x_s(e) = x_I(e)$ and $c_s(e) = c_I(e)$. By induction and continuity of the co-enumeration of the closed sets $F^{e,k}$, $H_s^e = \bigcap_{r \in I} H_r^e$. For all $r \in I$, $c_r(e)$ is the least $i \in \{0, 1\}$ such that $\lambda(H_r^e \mid x_I \upharpoonright_e \hat{\ } i) \geq 2^{-e}$. By induction, $x_s \upharpoonright_e = x_I \upharpoonright_e$, and by continuity, $\lambda(H_s^e \mid x_I \upharpoonright_e \hat{\ } i) = \inf_{r \in I} \lambda(H_r^e \mid x_I \upharpoonright_e \hat{\ } i)$ and so is at least 2^{-e} . On the other hand, if $i = 1$, then $\lambda(H_r^e \mid x_I \upharpoonright_e \hat{\ } 0) < 2^{-e}$ for all $r \in I$, and so $\lambda(H_s^e \mid x_s \upharpoonright_e \hat{\ } 0) < 2^{-e}$. Overall we see that $x_s(e) = x_I(e)$. The argument for $c_s(e)$ is the same. \square

We show that $\langle x_s \rangle$ changes only finitely often on each input. Claim 5.3.1 would then imply that $x = \lim_{s \rightarrow \omega_1^{\text{ck}}} x_s$. By induction on e we show that $\omega_1^{\text{ck}} + 1$ can be partitioned into finitely many closed intervals of stages on which both $x_s \upharpoonright_e$ and $c_s \upharpoonright_e$ are constant. Suppose that this has been shown for e ; let I be a closed interval of stages on which $x_s \upharpoonright_e$ and $c_s \upharpoonright_e$ are constant. For $i < 2$ let I_i be the set of stages $s \in I$ at which $x_s(e) = i$. By (*), both I_0 and I_1 are intervals, with $I_0 < I_1$. Claim 5.3.1 shows that they are closed. Now fix $i < 2$; let $t = \max I_i$ and let $k = c_t(e)$. For $m \leq k$ let $I_{i,m}$ be the set of stages at which $c_s(e) = m$. By (**), each $I_{i,m}$ is an interval with $I_{i,0} < I_{i,1} < \dots < I_{i,k}$, and $\bigcup_{m \leq k} I_{i,m} = I_i$. Claim 5.3.1 shows that each $I_{i,m}$ is closed. This concludes the proof of Proposition 5.3. \square

5.4. Separating Π_1^1 randomness from higher weak 2-randomness. In this section we construct a sequence $x \in 2^\omega$ which is higher weak 2-random but not Π_1^1 random. This sequence will be O -computable. The construction is an elaboration on that of the previous section. Here too we need to build an element of a $\mathbf{\Pi}_3^0$ set of measure 1 which is the intersection of higher Σ_2^0 sets, namely all of the ones of measure 1. To ensure that x is not Π_1^1 random we need to show that it collapses ω_1^{ck} , as in the presence of higher weak 2-randomness (and in fact Δ_1^1 randomness), being Π_1^1 random is equivalent to preserving ω_1^{ck} . So we will ensure that we can give x a collapsing approximation. On the other hand, Proposition 5.1 shows that we cannot give x a compact approximation, let alone a finite-change one. The difficulty of course compared to the previous construction is that we cannot effectively enumerate all of the higher Σ_2^0 sets of measure 1. In the indices of such sets, the property of having measure 1 is higher Π_1^0 but not decidable.

Technically, it is the key Claim 5.3.1 which may fail: if F^e does not really have measure 1, then it is possible that at every stage r in an interval I , F_r^e has measure 1, but for $s = \sup I$, F_s^e does not have measure 1. (For example, let $I = \omega$, and $F_r^{e,k} = 2^\omega$ when $r < k$ and empty when $r \geq k$; then $F_r^e = 2^\omega$ for all $r < \omega$ but

$F_\omega^e = \emptyset$). It is then possible that $c_r(e)$ cycles through all of ω during the stages in I . At stage s we know that we didn't need to ensure that $x \in F^e$. But by then it is too late, the approximation changed infinitely often.

Thus, we devise a wider class of approximations which is compatible with being higher weakly 2-random, but still implies collapsing ω_1^{ck} .

Definition 5.4. An ω_1^{ck} -computable approximation $\langle f_s \rangle$ of a function f is *finite-change along true initial segments* if for no n is there an increasing infinite sequence $\langle t(k) \rangle$ of stages such that $f_{t(k)} \upharpoonright_n = f \upharpoonright_n$ for all k but $f_{t(k+1)}(n) \neq f_{t(k)}(n)$ for all k .

To see that such an approximation is collapsing we isolate another notion.

Definition 5.5. An ω_1^{ck} -computable approximation $\langle f_s \rangle$ of a function f is a *club approximation* if for all n , the set of stages s such that $f_s \upharpoonright_n = f \upharpoonright_n$ is a closed set of stages.

Lemma 5.6. Every function which has a finite-change-along-true-initial-segments approximation also has a club approximation. If $\langle f_s \rangle$ is a club approximation of $f \notin \Delta_1^1$ then $\langle f_s \rangle$ is a collapsing approximation.

Proof. Suppose that $\langle f_s \rangle$ is an ω_1^{ck} -computable approximation of f which is finite-change along true initial segments. By induction on $n < \omega$ we see that if s is a limit stage and for unboundedly many $t < s$, $f_t \upharpoonright_n = f \upharpoonright_n$, then $f \upharpoonright_n = \lim_{t \rightarrow s} f_t \upharpoonright_n$. Similarly to what we did with finite-change approximations, we can make the approximation partially continuous by requiring, for every limit stage $s < \omega_1^{\text{ck}}$ and $n < \omega$, that if $\lim_{t \rightarrow s} f_t(n)$ exists, then it equals $f_s(n)$. This makes it a club approximation.

If $\langle f_s \rangle$ is a club approximation of f and s is least such that f lies in the closure of $\{f_t : t < s\}$ then $f = f_s$. Hence if $s < \omega_1^{\text{ck}}$ then f is hyperarithmetic. \square

The separation of Π_1^1 randomness from higher weak 2-randomness then follows from the following proposition.

Proposition 5.7. *There is a sequence x which is higher weak 2-random and has an ω_1^{ck} -computable approximation which changes finitely along true initial segments.*

The rest of this section is devoted to the proof of Proposition 5.7.

5.4.1. *Discussion.* The new idea is to “banish” strings which would contradict the property of the approximation being finite-change along true initial segments. That is, if I is an interval of stages, $x_r \upharpoonright_e$ is constant on I , but we see $x_r(e)$ changes infinitely often on I , then we require that $x_I \upharpoonright_e$ is not an initial segment of x . We simply do not allow any future x_t to extend $x_I \upharpoonright_e$. The construction is dynamic: rather than defining x and c a priori and then giving them approximations, we first define the approximation and then show it converges and has the desired properties.

We need to show that the construction can actually be carried out: at every stage there are non-banished strings that can be chosen to construct x_s , particularly non-banished strings relative to which we can make the sets H_s^e not too small.

This is done as follows:

1. At each length, we will banish at most one string. The continuity properties of the approximations to our sets will ensure that if we do see $c_r(e)$ cycle through all possible values in ω on an interval I of stages, then this will witness that F^e does

not have measure 1. Once we see that, we no longer need to force x to enter F^e (we can replace F^e by 2^ω). After this event there will be no need to banish another string of length e .

2. Nonetheless, even if just one string is banished, it is possible that this was the string on which H^e was large. I.e., it is possible that one string of length $e + 1$ is banished and the other is useless. To counter this we rely on a measure-theoretic observation which is the basis of Kučera's coding technique [Kuč85]. We spread out the levels of the construction, adding more than one bit between step e and $e + 1$. If the levels are sufficiently spread out, then every good string at level e has at least two good strings at the next level. So if one of them is banished, the other can still be used.

These are the ideas needed for the construction. We can now give the formal details.

5.4.2. Construction. We start with an effective enumeration $\langle F^e \rangle$ of all higher Σ_2^0 sets. So $F^e = \bigcup_k F^{e,k}$, an increasing sequence, with each $F^{e,k}$ a closed Σ_1^1 set. Each of these have co-enumerations $\langle F_s^{e,k} \rangle_{s < \omega_1^{ck}}$. We let $F_s^e = \bigcup_{k < \omega} F_s^{e,k}$. If s is a limit ordinal then $F_s^{e,k} = \bigcap_{t < s} F_t^{e,k}$.

We require that $F^{0,k} = 2^\omega$ for all k .

Let $\langle \ell(e) \rangle_{e < \omega}$ and $\langle \varepsilon^e \rangle_{e < \omega}$ be computable sequences such that:

- $\langle \ell(e) \rangle$ is an increasing sequence of natural numbers with $\ell(0) = (0)$.
- $\langle \varepsilon^e \rangle$ is a decreasing sequence of positive rational numbers with $\varepsilon^0 = 1$.
- For any $e < \omega$, for any measurable set A , and for any string σ of length $\ell(e)$, if $\lambda(A \mid \sigma) \geq \varepsilon^e/2$ then there are at least two extensions τ of σ of length $\ell(e + 1)$ such that $\lambda(A \mid \tau) \geq \varepsilon^{e+1}$.

If $\lambda(F^e) < 1$ then we may define during the construction a string ρ^e of length $\ell(e)$; this will be the “banished” string of length $\ell(e)$. We will ensure that the real we build does not extend ρ^e . [We required F^0 to have measure 1 to ensure that ρ^0 is never defined, as we would not have been able to avoid it.]

At every stage s we will define:

- A sequence $x_s \in 2^\omega$;
- A sequence of closed sets $\langle H_s^e \rangle$;
- A function $c_s \in (\omega + 1)^\omega$ which codes our choices which define the closed sets H_s^e . [A choice $k = c_s(e) < \omega$ indicates as before the choice of $F_s^{e,k}$; $c_s(e) = \omega$ indicates that $\lambda(F_s^e) < 1$.]

At a limit stage $s < \omega_1^{ck}$ we first see if we need to banish some strings. Let $e < \omega$ and suppose that $\lambda(F_s^e) < 1$ but that there is some final segment $I = [s_0, s)$ of s such that

- $\lambda(F_r^e) = 1$ for all $r \in I$;
- $c_r \upharpoonright_e$ is constant on the interval I ; and
- The string $x_r \upharpoonright_{\ell(e)}$ is constant on I .

Then we define $\rho^e = x_I \upharpoonright_{\ell(e)}$. We do this for all e for which this is needed. Note that $\lambda(F_t^e)$ is nonincreasing in t , and so for all e there may be at most one stage at which we want to define ρ^e .

We then define x_s , our choice function c_s and the closed sets H_s^e . To start, we let $H_s^0 = 2^\omega$. At step e we already have $x_s \upharpoonright_{\ell(e)}$, $c_s \upharpoonright_e$ and H_s^e . By induction, $\lambda(H^e \mid x \upharpoonright_{\ell(e)}) \geq \varepsilon^e [s]$.

At step e of stage s we first define $c_s(e)$ and H_s^{e+1} :

- If $\lambda(F_s^e) = 1$ then we let $c_s(e)$ be the least $k < \omega$ such that $\lambda(H^e \cap F^{e,k} \mid x \upharpoonright_{\ell(e)}) \geq \varepsilon^e/2 [s]$. We then let $H_s^{e+1} = H_s^e \cap F_s^{e,c_s(e)}$.
- If $\lambda(F_s^e) < 1$ then we let $c_s(e) = \omega$ and $H_s^{e+1} = H_s^e$.

We then define $x_s \upharpoonright_{\ell(e+1)}$:

- If ρ^{e+1} is undefined then we let $x_s \upharpoonright_{\ell(e+1)}$ be the leftmost extension σ of $x_s \upharpoonright_{\ell(e)}$ of length $\ell(e+1)$ such that $\lambda(H_s^{e+1} \mid \sigma) \geq \varepsilon^{e+1}$.
- If ρ^{e+1} is defined then we let $x_s \upharpoonright_{\ell(e+1)}$ be the leftmost extension σ of $x_s \upharpoonright_{\ell(e)}$ of length $\ell(e+1)$ other than ρ^{e+1} such that $\lambda(H_s^{e+1} \mid \sigma) \geq \varepsilon^{e+1}$.

This concludes the construction.

5.4.3. *Verification.* As above, if $e < \omega$, $s < t \leq \omega_1^{\text{ck}}$ and $c_s \upharpoonright_e = c_t \upharpoonright_e$ then $H_t^e \subseteq H_s^e$. This implies:

- (*) Suppose that $c_s \upharpoonright_e = c_t \upharpoonright_e$ and $x_s \upharpoonright_{\ell(e)} = x_t \upharpoonright_{\ell(e)}$. Then $c_s(e) \leq c_t(e)$.
- (**) Suppose that $c_s \upharpoonright_{e+1} = c_t \upharpoonright_{e+1}$ and $x_s \upharpoonright_{\ell(e)} = x_t \upharpoonright_{\ell(e)}$. Then $x_s \upharpoonright_{\ell(e+1)} \leq x_t \upharpoonright_{\ell(e+1)}$ (lexicographically).

For (**) note that if ρ^{e+1} is first defined between stages s and t , this only pushes $x_t \upharpoonright_{\ell(e+1)}$ further to the right. For (*) again note that if $\lambda(F_t^e) = 1$ then $\lambda(F_s^e) = 1$.

The following claim shows that banishing conforms to our original intention. Suppose that $c_r \upharpoonright_e$ and $x_r \upharpoonright_{\ell(e)}$ are constant on an interval I of stages. Suppose that $c_r(e) < \omega$ for all $r \in I$. By (*), $\sup_{r \in I} c_r(e) = \omega$ if and only if $c_r(e)$ changes infinitely often on I (there is an infinite increasing sequence $\langle t(k) \rangle$ of stages in I such that $c_{t(k+1)}(e) \neq c_{t(k)}(e)$).

Claim 5.7.1. Let $s < \omega_1^{\text{ck}}$ be a limit stage. Let $e < \omega$. Suppose that both $c_r \upharpoonright_e$ and $x_r \upharpoonright_{\ell(e)}$ are constant on a final segment I of s . Suppose that $c_r(e) < \omega$ for all $r \in I$ but that $\sup_{r \in I} c_r(e) = \omega$. Then at stage s we define $\rho^e = x_I \upharpoonright_{\ell(e)}$.

Proof. Let $\sigma = x_I \upharpoonright_{\ell(e)}$. If $r < t$ are in I then $H_t^e \subseteq H_r^e$. Let $H_{<s}^e = \bigcap_{t \in I} H_t^e$.

If $t \in I$ and $k < c_t(e)$ then $\lambda(H_t^e \cap F_t^{e,k} \mid \sigma) < \varepsilon^e/2$, and so $\lambda(H_{<s}^e \cap F_s^{e,k} \mid \sigma) < \varepsilon^e/2$. It follows that $\lambda(H_{<s}^e \cap F_s^e \mid \sigma) \leq \varepsilon^e/2$.

On the other hand, for all $t \in I$, $\lambda(H_t^e \mid \sigma) \geq \varepsilon^e$ and so $\lambda(H_{<s}^e \mid \sigma) \geq \varepsilon^e$. This shows that $\lambda(F_s^e) < 1$. The conditions for defining $\rho^e = \sigma$ at stage s are fulfilled. \square

Since each string of length $\ell(e)$ has only finitely many extensions of length $\ell(e+1)$, (**) and Claim 5.7.1 together imply:

Claim 5.7.2. Let $s < \omega_1^{\text{ck}}$ be a limit stage. Let $e < \omega$. Suppose that both $c_r \upharpoonright_e$ and $x_r \upharpoonright_{\ell(e)}$ are constant on a final segment I of s . Suppose that $x_r \upharpoonright_{\ell(e+1)}$ changes infinitely often on I (but not on a proper initial segment of I). Then at stage s we define $\rho^e = x_I \upharpoonright_{\ell(e)}$.

By induction on e we can show that eventually each $x_s \upharpoonright_{\ell(e)}$ and $c_s \upharpoonright_e$ are constant. We can let $x = \lim_{s \rightarrow \omega_1^{\text{ck}}} x_s$ and $c = \lim_{s \rightarrow \omega_1^{\text{ck}}} c_s$.

Claim 5.7.3. x is higher weak 2-random.

Proof. Let $e < \omega$, and suppose that $\lambda(F^e) = 1$. We show that $x \in F^e$. Let I be a final segment of ω_1^{ck} on which $c_t \upharpoonright_{e+1}$ is constant. Since $\lambda(F^e) = 1$, $k = c(e) < \omega$. For all $t \in I$ and all $d \geq e$, $H_t^d \subseteq F_t^{e,k}$; and $[x_t \upharpoonright_{\ell(d)}] \cap H_t^d$ is not null, and so nonempty. It follows that $[x \upharpoonright_{\ell(d)}] \cap F^{e,k}$ is nonempty. We then use the fact that $F^{e,k}$ is closed. \square

The proof of 5.7 is concluded by showing that $\langle x_s \rangle$ is an approximation which changes finitely often along true initial segments. To see this, it suffices to show that for no $e < \omega$ is there an increasing sequence $\langle t(k) \rangle$ of stages such that $x_{t(k)} \upharpoonright_{\ell(e)} = x \upharpoonright_{\ell(e)}$ for all $k < \omega$, but that $x_{t(k+1)} \upharpoonright_{\ell(e+1)} \neq x_{t(k)} \upharpoonright_{\ell(e+1)}$ for all $k < \omega$ (to verify Definition 5.4 for an arbitrary n , consider the greatest e such that $\ell(e) \leq n$). Suppose that such a sequence $\langle t(k) \rangle$ is given; let $s = \sup_k t(k)$. Let d be the greatest such that both $\lim_{r \rightarrow s} x_r \upharpoonright_{\ell(d)}$ and $\lim_{r \rightarrow s} c_r \upharpoonright_{\ell(d)}$ exist. So $d \leq e$. Either the conditions of Claim 5.7.1 or Claim 5.7.2 hold at stage s for d , so at stage s we define $\rho^d = x \upharpoonright_{\ell(d)}$. However the construction ensures that for all $d \geq 1$, x does not extend ρ^d (and that ρ^0 is never defined).

6. CLASSES OF HIGHER Δ_2^0 FUNCTIONS

Motivated by their usage in investigating higher weak 2-randomness, we study the classes of higher Δ_2^0 functions which we introduced above. We first consider the higher limit lemma.

6.1. The higher limit lemma. The proof of the higher limit lemma (Proposition 1.8) is not complicated. The equivalence of $f \leq_{\text{T}} O$ and $f \leq_{\omega_1^{\text{ckT}}} O$ was established in Proposition 2.1. If $f = \Phi(O)$ (where Φ is either c.e. or higher c.e.) then we can give f an ω_1^{ck} -computable approximation by letting $f_s = \Phi_s(O_s)$, where $\langle O_s \rangle$ is an ω_1^{ck} -computable enumeration of O . And if $\langle f_s \rangle$ is an ω_1^{ck} -computable approximation of f then the graph of f is Δ_2 over $L_{\omega_1^{\text{ck}}}$; since a set is Σ_2 over $L_{\omega_1^{\text{ck}}}$ if and only if it is c.e. in O , we see that f is O -computable.

In fact, the higher limit lemma relativises to every oracle. Recall that a subset X of $L_{\omega_1^{\text{ck}}}$ is A - ω_1^{ck} -computable (where $A \in 2^\omega$) if there is an ω_1^{ck} -c.e. $\Phi \subseteq 2^{<\omega} \times L_{\omega_1^{\text{ck}}}$ such that $X = \Phi(A)$. Also recall that we let J^A be the higher jump of A , the effective join of all subsets of ω which are higher A -c.e.

Proposition 6.1. *Let $A \in 2^\omega$. The following are equivalent for $f: \omega \rightarrow \omega$.*

- (1) $f \leq_{\omega_1^{\text{ckT}}} J^A$.
- (2) f has an A - ω_1^{ck} -computable approximation $\langle f_s \rangle_{s < \omega_1^{\text{ck}}}$.

Proof. Recall that we use an ω_1^{ck} -computable projection function $p: \omega_1^{\text{ck}} \rightarrow \omega$.

Assume (2); Let $m: \omega \rightarrow \omega_1^{\text{ck}}$ be the modulus of the sequence $\langle f_s \rangle_{s < \omega_1^{\text{ck}}}$: The value $m(n)$ is the least s such that for all $t > s$ we have $f_t(n) = f_s(n)$. Let $W = \{(n, p(s)) : s < m(n)\}$; the set W is higher A -c.e.: to enumerate $(n, p(s))$ into W , what we need from A is the value $f_s(n)$ and a different value $f_t(n)$ for some $t > s$; both are given with finitely much use of A . So $W \leq_{\omega_1^{\text{ckT}}} J^A$. Now, from one pair $(n, p(s)) \notin W$ and finitely much of A we output $f(n) = f_s(n)$. So $f \leq_{\omega_1^{\text{ckT}}} A \oplus W \leq_{\omega_1^{\text{ckT}}} J^A$.

Assume (1). Recall that we regard J as a higher enumeration functional. The sequence $\langle J_s^{A \upharpoonright_{p(s)}} \rangle$ is an A - ω_1^{ck} -computable approximation of J^A (using the fact

that for all n there is some $t < \omega_1^{\text{ck}}$ such that $p(s) \geq n$ for all $s \geq t$). Note that the sequence $\langle J_s^A \rangle$ is *not* A - ω_1^{ck} -computable.

If Ψ is a higher Turing functional then $\langle \Psi_s \left(J_s^{A \upharpoonright p(s)} \right) \rangle$ is an A - ω_1^{ck} -computable approximation of $\Psi(A)$.

We remark that it is not the case that for all A , J^A has an A - ω_1^{ck} -computable enumeration (an ω_1^{ck} -computable sequence $\langle A_s \rangle_{s < \omega_1^{\text{ck}}}$ such that $A_s \subseteq A_t$ for $s \leq t$). \square

6.2. Equivalent characterisations of classes. A couple of classes we defined have equivalent characterisations, some related to the limit lemma.

6.2.1. Higher ω -computably approximable functions. These were defined in Definition 1.13: functions approximable by finite-change approximations which moreover have hyperarithmetic bounds on the number of changes. In complete analogy with the lower case, this notion can be characterised by using strong reducibilities.

- Let $X, Y \in 2^\omega$. We say that X is higher truth-table reducible to Y if there is a hyperarithmetic sequence $\langle F_n \rangle$ of finite subsets of $2^{<\omega}$ such that $X(n) = 1$ if and only if Y extends some string in F_n . Nerode's argument shows that X is higher truth-table reducible to Y if and only if $X = \Phi(Y)$ for some higher Turing functional Φ which is total and consistent on all oracles.
- Let $f, g \in \omega^\omega$. We say that f is higher weak truth-table reducible to g if there is a higher Turing functional Φ such that $\Phi(g) = f$ and there is a hyperarithmetic function h such that for all axioms $(\tau, \sigma) \in \Phi$, $|\tau| \leq h(|\sigma|)$.

The lower-case arguments carry over to show that $X \in 2^\omega$ is higher ω -c.a. if and only if it is higher truth-table reducible to O ; and that $f \in \omega^\omega$ is higher ω -c.a. if and only if it is higher weak truth-table reducible to O .

6.2.2. Finite-change approximations. As discussed above, a finite-change approximation can be made continuous at limit stages. Hence, $f \in \omega^\omega$ has a finite-change approximation if and only if it has an approximation $\langle f_s \rangle$ such that for all limit $s < \omega_1^{\text{ck}}$, $f_s = \lim_{t \rightarrow s} f_t$.

We give a characterisation using a strong variant of the limit lemma.

Proposition 6.2. *The following are equivalent for $f \in \omega^\omega$:*

- (1) f has a finite-change approximation.
- (2) f is higher O -computable by a higher Turing functional Φ which is total (and consistent) on every subset of O .

Proof. (2) \implies (1): Let $\langle O_s \rangle$ be an ω_1^{ck} -computable enumeration of O . For $s < \omega_1^{\text{ck}}$ let $f_s = \Phi(O_s)$. For a limit $s < \omega_1^{\text{ck}}$, $O_s = \bigcup_{t < s} O_t$. Since Φ uses only finitely much of an oracle, $f_s = \lim_{t \rightarrow s} f_t$, so $\langle f_s \rangle$ is a finite-change approximation of f .

(1) \implies (2): this is a modification of the argument that every function which is higher ω -c.a. is higher weak truth-table reducible to O . Let $\langle f_s \rangle$ be a finite-change approximation of f . For all n and k we can compute some $d = d(n, k)$ such that $d \in O$ if and only if there are at least k changes in $\langle f_s(n) \rangle$. We then let $\Phi(X, n) = m$ if m is the k^{th} value of $f_s(n)$ observed, where k is the least such that $d(n, k) \notin X$. In other words, the procedure Φ queries an oracle X as if it were O , asking successively whether $\langle f_s(n) \rangle$ changes once, twice, thrice, ... until it finds X 's opinion on the number of changes; and outputs the corresponding value. If $X = O$

the answer is correct. If $X \subseteq O$ then the answer could be smaller than the actual number of changes but not larger, so the search for the k^{th} value will terminate. \square

6.2.3. Compact approximations.

Lemma 6.3. The following are equivalent for $x \in 2^\omega$:

- (1) x has a closed approximation.
- (2) x has an ω_1^{ck} -computable approximation $\langle x_s \rangle$ such that the closure of the set $\{x_s : s \leq \omega_1^{\text{ck}}\}$ is countable.

Proof. The idea is similar to that of the proof of Lemma 1.11. In the nontrivial direction, we first note that if y is a limit point of $\{x_s : s \leq \omega_1^{\text{ck}}\}$ other than x then there is an increasing sequence $\langle t(k) \rangle$ of stages such that $y = \lim_{k \rightarrow \omega} x_{t(k)}$. Further, for all limit $s < \omega_1^{\text{ck}}$, since the closure of $\{x_t : t < s\}$ is countable, this closure can be effectively obtained (again using the Cantor-Bendixon analysis). We now fatten the approximation $\langle x_s \rangle$ by inserting, for each limit $s < \omega_1^{\text{ck}}$, between $\langle x_t \rangle_{t < s}$ and x_s , all the limit points of $\{x_t : t < s\}$ which were not previously inserted. If $x_t \upharpoonright_n$ has stabilised before s , then all limit points extend this string, and so the fattened approximation still approximates x . \square

6.2.4. *Club approximations.* The class of approximations given by Definition 6.5 is mostly a tool which we use later, because it is easier to deal with than club approximations. To motivate that definition we first consider a ‘‘pointwise version’’.

Definition 6.4. An approximation $\langle f_s \rangle$ is *almost finite-change* if for all $n < \omega$, if $\langle t(i) \rangle$ is an increasing sequence of stages such that $f_{t(i+1)}(n) \neq f_{t(i)}(n)$ for all $i < \omega$, then $f_t(n)$ is constant on $[\sup_i t(i), \omega_1^{\text{ck}})$.

Suppose that an approximation $\langle x_s \rangle$ consists of elements of Cantor space and that it is partially continuous: for all $n < \omega$ and limit $s < \omega_1^{\text{ck}}$, if $\lim_{t \rightarrow s} f_t(n)$ exists then it equals $f_s(n)$. Then the approximation is almost finite-change if and only if for all $n < \omega$, for all $s < \omega_1^{\text{ck}}$ and $i < 2$, if the set $\{t < s : f_t(n) = i\}$ is not a closed subset of s , then $f_s(n) \neq i$.

Definition 6.5. An approximation $\langle f_s \rangle$ is *locally almost finite-change* if for all $n < \omega$ and all strings $\sigma \in \omega^n$, if $\langle t(i) \rangle$ is an increasing sequence of stages such that $f_{t(i)} \upharpoonright_n = \sigma$ and $f_{t(i+1)}(n) \neq f_{t(i)}(n)$ for all $i < \omega$, then $f_t(n)$ is constant on the stages $t \geq \sup t(i)$ at which $\sigma < f_t$.

Call an approximation $\langle f_s \rangle$ *locally continuous* if for all $n < \omega$ and all $\sigma \in \omega^n$, the function $f_t(n)$ is continuous on the set of stages t at which $\sigma < f_t$ (using the subspace topology). Namely, letting F_σ be that set of stages, if s is a limit point of F_σ which is also in F_σ , and $f_t(n)$ is constant on a final segment of $s \cap F_\sigma$, then $f_s(n)$ equals that constant value.

Lemma 6.6. Let $\langle x_s \rangle$ be a locally continuous approximation consisting of elements of Cantor space. Then the approximation is locally almost finite-change if and only if for all strings $\sigma \in \omega^{<\omega}$ and all $s < \omega_1^{\text{ck}}$, if the set $\{t < s : \sigma < f_t\}$ is not a closed subset of s , then $\sigma \not\prec f_s$.

Lemma 6.7. Every locally almost finite-change approximation is a club approximation. If $x \in 2^\omega$ has a club approximation then it has a locally almost finite-change approximation.

Proof. Let $\langle x_s \rangle$ be a club approximation of $x \in 2^\omega$. We may assume it is locally continuous (making it so does not change it being a club approximation). We define a locally continuous sequence $\langle y_s \rangle$ by recursion. At stage s we have already defined $\langle y_t \rangle_{t < s}$. For any string σ let F_σ be the set of stages t at which $\sigma < y_t$.

We call a string σ *forbidden* at stage s if the set $F_\sigma \cap s$ is not a closed subset of s . Otherwise a string is *permitted* at stage s . By induction, for all $t < s$, every initial segment of y_t is permitted at stage t .

The empty string is always permitted. Every string which is permitted at stage s has an immediate extension which is also permitted. To see this, suppose that σ is permitted but suppose, for a contradiction, that both $\sigma \hat{\ }0$ and $\sigma \hat{\ }1$ are forbidden at stage s . For $i < 2$ let r_i be the least stage $r < s$ which is a limit point of $F_{\sigma \hat{\ }i}$ but is not in $F_{\sigma \hat{\ }i}$. Since $y_{r_i}(|\sigma|)$ has just two possible values, $r_0 \neq r_1$. Say $r_0 < r_1$. But this means that $\sigma \hat{\ }0$ is forbidden at stage r_1 , so by induction we cannot have $\sigma \hat{\ }0 < y_{r_1}$, a contradiction.

We define y_s by induction. Suppose that $\sigma = y_s \upharpoonright_n$ is defined; by induction this string is permitted at stage s . We then act as follows:

- (1) If one extension $\sigma \hat{\ }i$ is forbidden at stage s then we let $y_s(n) = 1 - i$.
- (2) Otherwise, we let $y_s(n) = x_s(n)$.

The fact that $\langle x_t \rangle$ is locally continuous at s implies that so is $\langle y_t \rangle$. Hence, by the construction and by Lemma 6.6, the sequence $\langle y_s \rangle$ is locally almost finite-change.

By induction on $s < \omega_1^{\text{ck}}$ we observe that: (a) no initial segment of x is forbidden at s ; and (b) if σ is an initial segment of both x and x_s , then $\sigma < y_s$. We conclude that $x = \lim y_s$. \square

6.3. Enumerating approximations. In the next subsection we will prove non-implications between classes we defined above. When trying to diagonalise against a class of higher Δ_2^0 functions we need to enumerate an effective list of approximations. We discuss here when this is possible.

A *partial approximation* is a sequence $\langle f_t \rangle_{t < s}$ for some $s \leq \omega_1^{\text{ck}}$.

Lemma 6.8. There is an effective ω -enumeration of all ω_1^{ck} -computable partial approximations. That is, there is a partial array $\langle f_t^n \rangle$ for $n < \omega$ and $t < \omega_1^{\text{ck}}$ such that the function $(n, t) \mapsto f_t^n$ is partial ω_1^{ck} -computable, and every ω_1^{ck} -computable partial approximation equals $\langle f_t^n \rangle_{t < s}$ for some $n < \omega$.

Proof. There is a universal partial ω_1^{ck} -computable function. This allows us to devise an array $\langle f_t^\alpha \rangle$ for $\alpha, t < \omega_1^{\text{ck}}$ such that every ω_1^{ck} -computable partial approximation is $\langle f_t^\alpha \rangle$ for some $\alpha < \omega_1^{\text{ck}}$. Now renumber using the projection function p . \square

Uniformly we can totalise approximations: transform a given ω_1^{ck} -computable partial approximation $\langle g_s \rangle$ into an ω_1^{ck} -computable approximation $\langle f_s \rangle_{s < \omega_1^{\text{ck}}}$ such that if $\langle g_s \rangle$ is total and converges to some g , then $\lim_s f_s = g$ as well. This is similar to how it is done in lower computability, with care taken at limit stages. Namely, we define a non-decreasing function $t(s)$ which indicates the next expected g_t . At a successor stage s , if $g_{t(s-1)}$ is revealed by stage s , we let $f_s = g_{t(s-1)}$ and let $t(s) = t(s-1) + 1$; otherwise we let $t(s) = t(s-1)$ and $f_s = f_{s-1}$. At a limit stage s we let $t(s) = \sup_{r < s} t(r)$ and let $f_s(n) = \lim_{t \rightarrow s} f_t(n)$ when the limit exists, and 0 otherwise.

Thus, we can give an ω -list of total sequences $\langle f_s \rangle$, not all of which converge but for which the convergent ones list all higher Δ_2^0 functions. In some cases we can

do better. For example, as in the lower case, we can enumerate all higher ω -c.a. functions:

Lemma 6.9. There is a (total) ω_1^{ck} -computable array $\langle f_t^n \rangle_{n < \omega, t < \omega_1^{\text{ck}}}$ such that:

- For every n , $\langle f_t^n \rangle_{t < \omega_1^{\text{ck}}}$ is a higher ω -computable approximation of a function f^n .
- Every higher ω -c.a. function equals f^n for some n .

The construction is as expected. There is an ω_1^{ck} -list of all hyperarithmetic functions. Using the projection function p we can give a partial ω_1^{ck} -computable function $n \mapsto h^n$ which enumerates all hyperarithmetic functions. In fact by coupling it with partial approximations we can get a partial ω_1^{ck} -computable array $\langle h^n, g_t^n \rangle$ which lists all pairs $(h, \langle g_t \rangle)$ of hyperarithmetic functions and partial approximations.

We totalise as above, so we assume that each $\langle g_t^n \rangle$ is total. We then produce a total approximation $\langle f_t^n \rangle$. If h^n is not yet defined at stage t then f_t^n is the zero function. If h^n is defined at stage t and for no k have we seen more than $h^n(k)$ many changes on $\langle g_r^n(k) \rangle_{r \leq t}$ then we let $f_t^n = g_t^n$. Otherwise f_t^n is again the zero function.

6.3.1. Enumerating other approximations. On the other hand, many of the classes we listed above cannot be enumerated with corresponding approximations. For example, if $\langle f_t^n \rangle$ is a list of finite-change approximations, then it does not contain all functions with a finite-change approximation, as direct diagonalisation would verify. Informally, when we try to enumerate only finite change approximations, we track a sequence $\langle g_t \rangle$ up to a limit stage s at which we first see infinitely many changes on some input. At each stage $t < s$ we have only seen finitely many changes so we copy g_t . By stage s we have seen infinitely many changes but it is too late to go back and change the sequence.

A different difficulty is met when we try to enumerate almost finite-change or locally almost finite-change approximations (Definitions 6.4 and 6.5). Again diagonalisation shows we cannot list such approximations yielding all functions with these approximations. When we totalise approximations as above, starting with an almost-finite-change approximation we might inadvertently ruin this property. Take such an approximation $\langle g_t \rangle$ and suppose that the totalising process yields $\langle f_t \rangle$. Let s be a limit stage and suppose that at stage s we have seen $f_t(k)$ change infinitely often. We need to define f_s but since we are working uniformly, we cannot rely on the fact that $\langle g_t \rangle$ is total; we cannot wait to see what $g_s(k)$ is; the procedure above has us declare an arbitrary value for $f_s(k)$. When we later see that $g_s(k)$ is different it is too late. Either we change a later value of $f_t(k)$; this means that $\langle f_t \rangle$ is no longer an almost finite-change approximation. Or we can stick with the value $f_s(k)$; in this case $\langle f_t \rangle$ is an almost finite-change approximation, but $\lim_t f_t \neq \lim_t g_t$.

Luckily, for our purposes, we do not need tight restrictions on the kind of approximations we list. We will use the following two listings.

Lemma 6.10. There is a total ω_1^{ck} -computable array $\langle x_t^n \rangle$ of elements of Cantor space such that:

- For all n , $\langle x_t^n \rangle$ converges to a real x^n ; and

- If a real $x \in 2^\omega$ has an approximation which changes finitely along true initial segments, then there is some n such that $x = x_n$ and $\langle x_n \rangle$ changes finitely along true initial segments.

Proof. Given a partial approximation, we totalise it to a sequence $\langle y_s \rangle_{s \leq \omega_1^{\text{ck}}}$ as above, except that at limit stages we make the approximation locally continuous (for limit s we inductively define $y_s(n)$ to be the limit of $y_t(n)$ over the stages $t < s$ at which $y_t \upharpoonright_n = y_s \upharpoonright_n$, 0 if the limit does not exist). If the original approximation changes finitely along true initial segments, so does $\langle y_s \rangle$. We can then transform the approximation to be locally almost finite-change, in particular ensuring it has a limit. This follows the construction in the proof of Lemma 6.7, tracking forbidden strings. Again, if $\langle y_s \rangle$ changes finitely along true initial segments, so does the new approximation. \square

Lemma 6.11. There is a total ω_1^{ck} -computable array $\langle x_t^n \rangle$ of elements of Cantor space such that:

- For all n , $\langle x_t^n \rangle$ converges to a real x^n .
- Every real $x \in 2^\omega$ which has a club approximation equals x^n for some n .

Proof. The idea is to transform partial approximations $\langle x_t \rangle$ into “nearly” locally almost finite-change total approximations. Totalise as above, making the approximation $\langle y_s \rangle$ locally continuous. Once we have seen, for some $\sigma \in 2^n$, infinitely many changes in $y_t(n)$ on the set of stages at which $\sigma < y_t$, we set $y_s(n) = 0$, but later allow one last change, if we see the value 1 show up in the approximation $\langle x_t \rangle$. \square

6.4. Separations between classes. None of the classes we defined in the previous sections coincide. For a summary see Fig. 1. All implications were discussed above. In this section we show that no other implications hold. In fact, all separations are made in Cantor space.


6.4.1. *A real with a finite-change approximation which is not ω -c.a.* This is a simple diagonalisation argument, using Lemma 6.9, but working in Cantor space. Let $\langle x_t^n \rangle$ be as given by the lemma (with $x_t^n \in 2^\omega$). Define $y \in 2^\omega$ by letting $y(n) = 1 - x^n(n)$. Then $\langle x_t^n(n) \rangle$ is a finite-change approximation of y .

6.4.2. *A real with a finite-change approximation along true initial segments, but no compact approximation.* An example for such a real is given by Proposition 5.7 (using Proposition 5.1).

6.4.3. *A higher Δ_2^0 real which collapses ω_1^{ck} but has no collapsing approximation.* In [BGHM] we construct a higher Δ_2^0 real y below which higher Turing and fin-h reducibility differ. By Proposition 2.3, the real y collapses ω_1^{ck} . By Proposition 2.2, y does not have a collapsing approximation.

6.4.4. *A real with a club approximation but no approximation which is finite-change along true initial segments.* This is a slightly finer diagonalisation argument. Let $\langle x_t^n \rangle$ be the array given by Lemma 6.10. We build an approximation $\langle y_t \rangle_{t < \omega_1^{\text{ck}}}$ and diagonalise against each x^n by showing that $y \upharpoonright_{n+1} \neq x^n \upharpoonright_{n+1}$, provided $\langle x_t^n \rangle$ changes finitely along true initial segments.

To ensure that y has a club approximation we follow the construction of the proof of Lemma 6.7. As in that construction, define the sets F_σ , and the notion of a string being permitted or forbidden at stage s . We again ensure that all initial

FIGURE 1. Classes of higher Δ_2^0 reals

segments of each x_t are permitted at stage t and that the approximation is locally continuous.

At stage s , given $\langle y_t \rangle_{t < s}$, define y_s by recursion. We are given $\sigma = y_s \upharpoonright_n$, which by induction is permitted at stage s . Then:

- (1) If an immediate extension $\sigma \hat{\ } i$ of σ is forbidden at stage s , then we let $y_s(n) = 1 - i$.
- (2) If s is a limit stage, F_σ is cofinal in s and x_t^n is a constant i on a final segment of $F_\sigma \cap s$, then we let $y_s(n) = i$ (note that the assumption implies that $\sigma \hat{\ } i$ is permitted at every stage $t < s$, and so also at s).
- (3) Otherwise, we consider the set $A_\sigma = \{t < \omega_1^{\text{ck}} : \sigma < x_t^n\}$. If $x_s^n(n)$ changes infinitely along the stages in $A_\sigma \cap s$ (there is an increasing sequence $\langle t(i) \rangle$ of stages $t(i) \in A_\sigma \cap s$ such that $x_{t(i+1)}^n(n) \neq x_{t(i)}^n(n)$ for all $i < \omega$) then we let $y_s(n) = 0$. Otherwise, $y_s(n)$ is a constant i on a final segment of $A_\sigma \cap s$;⁶ we let $y_s(n) = 1 - i$.

By construction, the sequence $\langle y_s \rangle$ is locally almost finite-change, and so $y = \lim_s y_s$ has a club approximation. Let $n < \omega$ such that $\langle x_t^n \rangle$ changes finitely along true initial segments. Let $\sigma = y \upharpoonright_n$. If $\sigma \neq x^n \upharpoonright_n$ we are done, so we assume that $\sigma < x^n$ as well. The value $x_t^n(n)$ changes finitely often on A_σ . By induction on $s < \omega_1^{\text{ck}}$ we see that the value $y_t(n)$ changes only finitely often on F_σ and that both $\sigma \hat{\ } 0$ and $\sigma \hat{\ } 1$ are permitted at s . We then succeed in ensuring that $y(n) \neq x^n(n)$.

⁶This includes the case that $A_\sigma \cap s$ has a greatest element t ; then $i = x_t^n(n)$.

6.4.5. *A real with a compact approximation but no club approximation.* To show that there are no more implications in Fig. 1, it remains to show that there is a real $x \in 2^\omega$ which has a closed approximation but not a club approximation. Note that this also shows that there is a real which has a collapsing approximation but not a club approximation.

To construct a real with a closed approximation we use the following.

Lemma 6.12. Let $\langle x_s \rangle$ be an ω_1^{ck} -computable approximation of $x \in 2^\omega$. Suppose that for all limit $s < \omega_1^{\text{ck}}$ there are at most finitely many n such that $\lim_{t \rightarrow s} x_t(n)$ does not exist. Then x has a closed approximation.

Proof. We use Lemma 6.3. Since ω_1^{ck} is countable, it suffices to show that for all limit $s < \omega_1^{\text{ck}}$ there are at most countably many y which are the limit $\lim_{i \rightarrow \omega} x_{t(i)}$ where $\langle t(i) \rangle$ is increasing and $s = \sup_k t(i)$. But the condition implies that for a fixed s , all such y differ on only finitely many bits. \square

We in fact show the following.

Proposition 6.13. *No uniform listing of higher Δ_2^0 elements of Cantor space contains all reals with closed approximations. That is, if $\langle x_t^n \rangle$ is an ω_1^{ck} -computable array such that for all n , $\langle x_t^n \rangle$ converges to a real x^n , then there is some $y \in 2^\omega$ with a closed approximation which equals none of the x^n .*

We then use Lemma 6.11 to obtain the desired separation.

To prove Proposition 6.13 we will in fact build an approximation $\langle y_t \rangle$ such that for all limit $s < \omega_1^{\text{ck}}$ there is at most one $k < \omega$ such that $\lim_{t \rightarrow s} y_t(k)$ does not exist.

The naïve approach, letting $y_t(n) = 1 - x_t^n(n)$, will of course not work, since it is likely that for some limit $s < \omega_1^{\text{ck}}$, $x_t^n(n)$ change infinitely often up to s for more than one n . However we can choose other witnesses k to diagonalise y against x^n . Adding bounded injury to the argument makes it work.

In detail, along with $\langle y_t \rangle$ we also define a sequence of witnesses k_t^n for all $n < \omega$ and $t < \omega_1^{\text{ck}}$. Witnesses for different n are distinct; this is achieved by requiring that $k_t^n \in \omega^{[n]}$ (the n^{th} column of ω) for all n . Once the witnesses k_t^n are defined, y_t is determined by letting:

- $y_t(k_t^n) = 1 - x_t^n(k_t^n)$ for all $n < \omega$; and
- $y_t(k) = 0$ if $k \neq k_t^n$ for all n .

The idea is that if we see y change on k_t^n then we discard k_t^m for $m > n$. In detail: at stage s , we need to define a new witness k_s^n in case either

- (1) s is a limit stage and k_t^n is not stable below s (k_t^n is not constant on a final segment of s); or
- (2) for some $m < n$, it is not the case that $k = k_{<s}^m$ and $i = x_{<s}^m$ are well-defined and $x_s^m(k) = i$. In other words, either
 - s is a successor stage and for $k = k_{s-1}^m$ we have $x_{s-1}^m(k) \neq x_s^m(k)$; or
 - s is a limit stage, and either k_t^m is not stable below s ; or it is, with value k , but $x_t^m(k)$ is not stable below s ; or it is, with value i , but $x_s^m(k) \neq i$.⁷

⁷We could omit the very last case by requiring that $\langle x_t^n \rangle$ is partially continuous.

In all cases, we let k_s^n be the $p(s)^{\text{th}}$ element of the column $\omega^{[n]}$. If none of these cases hold, then we let $k_s^n = k_{<s}^n$, where as usual this means k_{s-1}^n if s is a successor stage, or the stable value k_I^n for some final segment I of s if s is a limit stage.

This concludes the construction. By induction on n we see that each k_t^n reaches a limit k^n and that $y(k^n) \neq x^n(k^n)$. It remains to show the condition which implies compactness. Let $s < \omega_1^{\text{ck}}$ be a limit stage. Suppose that there is $n < \omega$ such that k_t^n is stable on a final segment I of s (with value k_s^n), but that $\lim_{t \rightarrow s} x_t^n(k_s^n)$ does not exist. For all $m < n$, both k_t^m and $x_t^m(k_t^m)$ are stable on I . If $k \neq k_s^m$ for all $m \leq n$ then on a final segment of s , $k \neq k_t^m$ for all $m < \omega$ (if it is ever chosen, it is discarded before stage s), and so $y_t(k) = 0$ on a final segment of s . This shows that $\lim_{t \rightarrow s} y_t(k)$ exists for all $k \neq k_s^n$.

If no such n exists, then by induction on n we see that both k_t^n and $x_t^n(k_t^n)$ are stable below s (though likely there is no single final segment I of s on which they are all stable). Thus if $k = k_s^n$ for some n , then $\lim_{t \rightarrow s} y_t(k)$ exists. Suppose that $k \neq k_s^n$ for all n . Say $k \in \omega^{[n]}$. Then $k \neq k_t^m$ for all $m \neq n$ and all t ; and $k \neq k_t^n$ on a final segment of s , so again $y_t(k) = 0$ on a final segment of s .

6.5. A remark on club approximations. We can weaken Definition 5.5 as follows.

Definition 6.14. An ω_1^{ck} -computable sequence $\langle f_s \rangle_{s < \omega_1^{\text{ck}}}$ is a *club quasi-approximation* of a function f if for all $n < \omega$, the set of stages s at which $f \upharpoonright_n = f_s \upharpoonright_n$ is a closed and unbounded subset of ω_1^{ck} .

The point is that we do not require that $f = \lim_s f_s$. If $\langle f_s \rangle$ is a club approximation of any function, then this function is determined uniquely: for each string σ , $\{s : \sigma < f_s\}$ is an ω_1^{ck} -computable set, and the intersection of finitely many ω_1^{ck} -computable club subsets of ω_1^{ck} is a club subset of ω_1^{ck} .

For elements of Cantor space we get nothing new: if $x \in 2^\omega$ has a club quasi-approximation then it has a club approximation, in particular it is higher Δ_2^0 . However there are elements of Baire spaces which have club quasi-approximations but are not higher Δ_2^0 .

To see this, following the discussion in Section 6.3, fix a total ω_1^{ck} -computable array $\langle f_t^n \rangle$ of functions which contains all ω_1^{ck} -computable approximations. We define a sequence $\langle g_t \rangle_{t < \omega_1^{\text{ck}}}$ which is a club quasi-approximation of $g \in \omega^\omega$, ensuring that if $\langle f_t^n \rangle$ converges to some $f^n \in \omega^\omega$ then $g(n) \neq f^n(n)$. In fact we will ensure a stronger property than required: for all n , the set of stages $t < \omega_1^{\text{ck}}$ such that $g_t(n) = g(n)$ is closed and unbounded. The definition is simple: at a limit stage s we let $g_s(n) = \lim_{t \rightarrow s} g_t(n)$ if the limit exists, and 0 otherwise. At a successor stage s we compare $g_{s-1}(n)$ and $f_s^n(n)$. If they are distinct we let $g_s(n) = g_{s-1}(n)$. If they are equal to a nonzero value, we let $g_s(n) = 0$. If they are both equal to 0 then we let $g_s(n) = p(s)$, where as usual $p: \omega_1^{\text{ck}} \rightarrow \omega$ is ω_1^{ck} -computable and injective. Now the point is that for all $k \neq 0$, the set of stages $\{t < \omega_1^{\text{ck}} : g_t(n) = k\}$ is an interval of stages and so closed; and that the set of stages $\{t < \omega_1^{\text{ck}} : g_t(n) = 0\}$ is closed. By admissibility of ω_1^{ck} , one of these sets must be unbounded.

Finally we remark that the proof of the second part of Lemma 5.6 (that every club approximation of $f \notin \Delta_1^1$) shows that every club quasi-approximation of $f \notin \Delta_1^1$ is “quasi collapsing” in that the sequence of stages $s(n)$ at which we first observe $f \upharpoonright_n$ is unbounded in ω_1^{ck} . Hence if $f \notin \Delta_1^1$ has a club quasi-approximation then $\omega^f > \omega_1^{\text{ck}}$, even if f is not higher Δ_2^0 .

7. THE CLASS $\text{MLR}[O]$

It is not very hard to prove that one can characterize weak 2 randomness using a restricted relativisation of ML-randomness to \emptyset' . Define an $\text{MLR}[A]$ -test to be a nested test $\langle U_n \rangle$ satisfying $\lambda(U_n) \leq 2^{-n}$, where each U_n is *effectively* open (not A -effectively open), but an index for each U_n is given by A . That is, $U_n = W_{f(n)}$ where $\langle W_e \rangle$ enumerates effectively open sets and $f \leq_T A$. Weak 2 randomness is equivalent to $\text{MLR}[\emptyset']$ -randomness.

One direction is straightforward; given a weak 2 test $\langle V_n \rangle$, \emptyset' can find the least m such that $\lambda(V_m) \leq 2^{-n}$. The other direction requires a time-trick: if that $\langle W_{f(n)} \rangle$ is a test as described then we cover it with the null Π_2^0 set $\bigcap_{n,t} \bigcup_{s>t} W_{f_s(n)}$. Trying to lift the argument to the higher setting fails since the intersection would be over $\omega \times \omega_1^{\text{ck}}$ -many higher open sets, and we have no way to effectively convert this to an ω -list.

We shall indeed prove that the notion of higher Martin-Löf randomness, where Kleene's O can be used for the index of each component is much stronger than higher weakly 2-randomness, and even stronger than Π_1^1 -randomness. We now let $\langle W_e \rangle$ enumerate the *higher* effectively open sets.

Definition 7.1. Let $A \in 2^\omega$. A *higher* $\text{MLR}[A]$ -test is a nested sequence $\langle W_{f(n)} \rangle$ where $f \leq_{\omega_1^{\text{ck}}T} A$ and $\lambda(W_{f(n)}) \leq 2^{-n}$. The null set determined by such a test is $\bigcap_n W_{f(n)}$. A sequence is in $\text{MLR}[A]$ if it is not captured by any $\text{MLR}[A]$ -test.

Of course for Kleene's O the index-function f can be taken to be O -computable (Proposition 2.1); however the building blocks are still higher effectively open sets.

We start by giving an alternate characterisation of $\text{MLR}[O]$. A *long (higher) ML-test* is a sequence $\langle U_\alpha \rangle_{\alpha < \omega_1^{\text{ck}}}$ of uniformly higher effectively open sets such that $\bigcap_\alpha U_\alpha$ is null. No assumption is made about nesting.

Lemma 7.2. Higher $\text{MLR}[O]$ tests and long ML-tests capture the same null sets.

Proof. One direction follows the failed time trick: if $\bigcap_n W_{f(n)}$ is an $\text{MLR}[O]$ test then for $n < \omega$ and $s < \omega_1^{\text{ck}}$ we let $V_{n,t} = \bigcup_{s>t} W_{f_s(n)}$. We can reorder the array $\langle V_{n,s} \rangle$ effectively in ordertype ω_1^{ck} using an effective bijection between $\omega \times \omega_1^{\text{ck}}$ and ω_1^{ck} . If t is sufficiently late then $V_{n,t} = W_{f(n)}$.

In the other direction let $\langle U_s \rangle_{s < \omega_1^{\text{ck}}}$ be a long ML-test. Using O , for each n we can find a finite set $F \subset \omega_1^{\text{ck}}$ such that $\lambda(\bigcap_{s \in F} U_s) \leq 2^{-n}$ (the measure of a higher effectively open set is O -computable, uniformly). \square

Hirschfeldt and Miller (see [DH10]) showed that a ML-random sequence is weak 2 random if and only if it forms a minimal pair with \emptyset' ; the witness for failure of this property can be taken to be c.e. The situation is more complicated in the higher setting. Higher weak 2 randomness does not seem to align with such a property. In [GM] the authors show that Π_1^1 -randomness partly corresponds to this property: a higher ML-random sequence X is Π_1^1 -random if and only if there is no higher-c.e., non hyperarithmetic set higher Turing reducible to X . However, not every Π_1^1 -random sequence forms a minimal pair with Kleene's O in the higher Turing degrees; by the Gandy basis theorem, there is a Π_1^1 -random sequence computable from O .

Higher $\text{MLR}[O]$ gives a certain analogue of the Hirschfeldt-Miller property. Recall that we extended the notion of higher Turing reducibility to subsets of ω_1^{ck} in the obvious way.

Proposition 7.3. *The following are equivalent for a higher ML-random sequence X :*

- (1) $X \notin \text{MLR}[O]$.
- (2) X higher Turing computes an ω_1^{ck} -c.e. subset of ω_1^{ck} which is not ω_1^{ck} -computable.
- (3) X higher Turing compute a Δ_2 subset of ω_1^{ck} which is not ω_1^{ck} -computable.
- (4) There is some higher Δ_2^0 subset of ω which is not higher c.e., but is higher c.e. in X .

We note that the lower setting analogue of property (4) does characterise weak 2 randomness, a fact which has not been observed so far.

Proof. (1) \rightarrow (2): the lowercase argument can be copied to the higher setting. Let $\langle V_\alpha \rangle$ be a long ML-test capturing X . Using an indexing of all finite subsets of ω_1^{ck} (and taking finite intersections) we may assume that for all $\varepsilon > 0$, there are unboundedly many α such that $\lambda(V_\alpha) < \varepsilon$. We enumerate an ω_1^{ck} -c.e. subset $A \subseteq \omega_1^{\text{ck}}$, attempting to meet the requirements P_β : the complement of A is not W_β , where $\langle W_\beta \rangle$ is an ω_1^{ck} -effective sequence of all ω_1^{ck} -c.e. subsets of ω_1^{ck} . Suppose that a requirement P_β has not been initialised since stage $t < \omega_1^{\text{ck}}$, is not yet met at stage $s > t$, and that at stage $s > t$ we see that some $\alpha \in W_{\beta,s}$ and $\lambda(V_{\alpha,s}) \leq 2^{-p(\beta)}$ for some $\alpha \in \omega_1^{\text{ck}[\beta]}$. Then we enumerate α into A_{s+1} and initialise every requirement P_γ where $p(\gamma) > p(\beta)$. We also let $G_\alpha = V_{\alpha,s}$. If $\alpha \notin A$ then we let $G_\alpha = \emptyset$. Then $\langle G_\alpha \rangle$ is a higher Solovay test, and if X is not captured by this test then $A \leq_{\omega_1^{\text{ckT}}} X$.

(3) \rightarrow (4): Say $B \leq_{\omega_1^{\text{ckT}}} O$ is not ω_1^{ck} -computable and that $B \leq_{\omega_1^{\text{ckT}}} X$. Then $p[B]$ is higher X -c.e. but is not hyperarithmetic.

(4) \rightarrow (1): Let $C \subset \omega$ be O -computable, not higher c.e., but higher X -c.e. The usual majority-vote argument shows that the set of oracles Y such that C is higher Y -c.e. is null. Let $\langle C_s \rangle_{s < \omega_1^{\text{ck}}}$ be an ω_1^{ck} -computable approximation of C , and let Γ be a higher enumeration functional. For $n, k < \omega$ and $t < \omega_1^{\text{ck}}$ let $V_{n,k,t}$ be the set of $Y \in 2^\omega$ such that for some $s > t$, either:

- $n \in C_s$ and $n \in \Gamma_s^X$; or
- $n \notin C_s$ and $n \notin \Gamma_s^{X \upharpoonright k}$.

Then $\langle V_{n,k,t} \rangle$ is a long ML-test which captures the oracles Y such that $\Gamma^Y = C$. \square

Finally we show that higher $\text{MLR}[O]$ -randomness is strictly stronger than Π_1^1 -randomness.

Proposition 7.4. *Higher $\text{MLR}[O]$ -randomness is strictly stronger than Π_1^1 -randomness.*

Proof. As mentioned before, there is an O -computable Π_1^1 -random sequence; no higher $\text{MLR}[O]$ -random sequence can be O -computable.

Suppose that X is not Π_1^1 -random; we show it is not higher $\text{MLR}[O]$ -random. We assume that X is higher ML-random. By [Mon14], there exists a uniformly higher effectively open sequence $\langle U_n \rangle$ such that $X \in \bigcap_n U_n$ but X is not an element of

any higher effectively closed set $F \subseteq \bigcap_n U_n$. The set of canonical indices of higher effectively closed subsets of $\bigcap_n U_n$ is higher c.e.; this gives us a sequence $\langle P_\alpha \rangle_{\alpha < \omega_1^{\text{ck}}}$ which enumerates the higher effectively closed subsets of $\bigcap_n U_n$. Then the sequence $\langle U_n \rangle$ together with the sequence of the complements of the P_α 's gives a long ML-test which captures X . \square

8. HIGHER OBERWOLFACH RANDOMNESS (WITH DAN TURETSKY)

Oberwolfach randomness [BGK+16] is the notion of randomness which captures computing all K -trivials: a ML-random sequence computes all K -trivial sets if and only if it is not Oberwolfach random. The higher analogue holds.

Definition 8.1. A *higher Oberwolfach test* is a pair $(\langle G_\sigma \rangle, \alpha)$ where:

- For $\sigma \in 2^{<\omega}$, G_σ is (uniformly) higher effectively open, and $\lambda(G_\sigma) \leq 2^{-|\sigma|}$;
- The array is nested, in the sense that if $\sigma \preceq \tau$ then $G_\tau \subseteq G_\sigma$; and
- $\alpha \in 2^\omega$ is a higher left-c.e. sequence.

The null set determined by the test is $\bigcap_{n < \omega} G_{\alpha \upharpoonright n}$. A sequence is higher Oberwolfach random if it is not captured by any higher Oberwolfach test.

Proposition 5.2 shows that every higher weak 2 random sequence is higher Oberwolfach random; higher difference randomness can be characterised using “version-disjoint” higher Oberwolfach tests and so higher Oberwolfach randomness implies higher difference randomness (this follows from the proof of one of the implications in Proposition 1.8, and is identical to the lower setting). In fact both implications are strict. It is not difficult to build a higher Oberwolfach random sequence with a compact approximation, and then appeal to Proposition 5.1 to separate between higher weak 2 randomness and higher Oberwolfach randomness. To separate between higher Oberwolfach randomness and higher difference randomness we need to appeal to the forcing used by Day and Miller [DM15] to construct a difference random set which is not a density one point in effectively closed sets; the argument can be performed in the higher setting without change, both constructing such a random and showing that such a random cannot be higher Oberwolfach random.

The characterisation of higher Oberwolfach randomness in terms of computing K -trivial sets consists of two steps:

Theorem 8.2. *If X is higher ML-random but not higher Oberwolfach random, then it higher Turing computes every higher K -trivial set.*

Theorem 8.3. *There is a higher K -trivial set which is not higher computable from any higher Oberwolfach random sequence.*

A set A as given by Theorem 8.3 is called a “smart” K -trivial set: any higher ML-random sequence which higher computes A , must higher compute all higher K -trivial sets.

The usual proof of the lower-setting analogue of Theorem 8.2 passes through a characterisation of Oberwolfach randomness in terms of weak 2 tests which are bounded by *additive cost functions*. These are weak 2 tests $\langle U_n \rangle$ whose measure is bounded by $\alpha - \alpha_n$, where $\langle \alpha_n \rangle$ is an increasing approximation of a left-c.e. real α . By their very definition these use a time-trick. We can emulate the time trick by working over a K -trivial oracle.

First proof of Theorem 8.2. Let X be higher ML-random but not higher Oberwolfach random. Since X is not higher weak 2 random, the Hirschfeldt-Miller argument shows that there is some non-hyperarithmetic, higher c.e. set B which is higher Turing reducible to X (in fact this is true for any higher ML-random which is not Π_1^1 -random). We may assume that X is higher difference random, and so the Hirschfeldt-Nies-Stephan argument shows that B is higher K -trivial. The idea is to work relative to B and emulate the proof in [BGK⁺16].

Since B has a collapsing approximation, working relative to B we can revert to computability of length ω (see Lemma 4.2 and its footnote). Let $\langle g(n) \rangle$ be an increasing, cofinal sequence in ω_1^{ck} which is ω_1^{ck} -computable from B ; let $\langle \langle G_\sigma \rangle, \alpha \rangle$ be a higher Oberwolfach test capturing X . We let $U_n = \bigcup_{s \geq g(n)} G_{\alpha_s \upharpoonright k}$. Then $\langle U_n \rangle$ is nested and uniformly higher B -c.e.; and $\lambda(U_n) \leq 2^{-n} + (\alpha - \alpha_{g(n)})$. By delaying the approximation of U_n we can also suppose $\lambda(U_{n,g(m)}) \leq 2^{-n} + (\alpha_{g(m)} - \alpha_{g(n)})$ for each n and m .

Let $\mathbf{c}(k, s) = \alpha_s - \alpha_{g(k)}$. The aim is to find a higher B -computable approximation $\langle A_n \rangle_{n < \omega}$ of A such that letting $k(n) = |A_{n-1} \wedge A_n|$ (the least k such that $A_n(k) \neq A_{n-1}(k)$), we have $\sum_{n < \omega} (\alpha_{g(n)} - \alpha_{g(k(n))})$ is finite (we may assume that $k(n) \leq n$; otherwise we replace $\alpha_{g(n)} - \alpha_{g(k(n))}$ by 0). Once we have such an approximation we can define a higher B -Solovay test $\langle G_k \rangle$ by letting $G_k = U_{k,g(n)}$ if n is the greatest such that $k = k(n)$ (and $G_k = \emptyset$ if there is no such n). Since B is higher K -trivial, X cannot be captured by this test, and then the usual argument builds a higher B -c.e. functional Φ such that $\Phi(X) = A$. Since $B \leq_{\omega_1^{\text{ckT}}} X$ we get $A \leq_{\omega_1^{\text{ckT}}} X \oplus B \leq_{\omega_1^{\text{ckT}}} X$ as required.

To obtain the required approximation $\langle A_n \rangle$ we can operate in two ways. We define the higher B -c.e. oracle discrete measure $\mu^\tau(n) = \alpha_{g(n+1)} - \alpha_{g(n)}$ (for all strings τ of length n). One way is to use the fact that A is higher K -trivial relative to B ; we repeat the proof of the main lemma in $(L_{\omega_1^{\text{ck}}}; \epsilon, B)$ and use it for the measure μ^A . Another way is to directly use the unrelativised main lemma (Proposition 4.3). Recall that we can let $g(n)$ be the least s such that $B_s \upharpoonright n = B \upharpoonright n$ for some fixed higher enumeration $\langle B_s \rangle_{s < \omega_1^{\text{ck}}}$ of B . For $t < \omega_1^{\text{ck}}$ we let $g_t(n)$ be the least $t \leq s$ such that $B_s \upharpoonright n = B_t \upharpoonright n$. Note that $\sup_n g_s(n) = s$ and that $g_s(n) \leq g(n)$. For all τ of length n we let $\mu_s^{B_s \upharpoonright n \oplus \tau}(n) = \alpha_{g_s(n+1)} - \alpha_{g_s(n)}$. Let $\langle A_s \rangle$ be a collapsing approximation of A . The main lemma gives us an ω_1^{ck} -computable closed and unbounded set $C \subseteq \omega_1^{\text{ck}}$, such that the sum $\sum_{s \in C} (\alpha_s - \alpha_{g_s(k(s))})$ is finite; here $k(s) = |A_s(k) \wedge A_{s^+}(k)|$, where s^+ is the next element of C beyond s . We define the required B -computable approximation of A by letting $\hat{A}_n = A_{s(n)}$ for some $s(n) \in C$, $s(n) \geq g(n)$ (for example $s(n) = \min(C - g(n))$). Let $k = |\hat{A}_{n-1} \wedge \hat{A}_n|$. Then there is some $s \in [s(n), s(n+1))$ such that $k \geq |A_s \wedge A_{s^+}|$. Since $s \geq g(n)$, $\alpha_{g(n)} - \alpha_{g(k)} \leq \alpha_s - \alpha_{g_s(k)}$. \square

We can however eliminate the time trick, with an argument which also works in the lower setting. Rather than use additive cost functions, we use cost functions which in the lower setting are “subadditive”. If μ is a discrete measure then we let $\mathbf{c}_\mu(n) = \sum_{m \geq n} \mu(m)$. If $\langle \mu_s \rangle$ is an increasing enumeration of a left-c.e. discrete measure μ then we let $\mathbf{c}_\mu(n, s) = \sum_{m \geq n} \mu_s(m)$. We say that an approximation $\langle A_s \rangle_{s < \omega_1^{\text{ck}}}$ of a set A witnesses that A obeys \mathbf{c}_μ if the sum $\sum_{s < \omega_1^{\text{ck}}} \mathbf{c}(|A_s \wedge A_{s+1}|, s)$ is finite. If μ is the optimal left-c.e. discrete measure then any set obeying \mathbf{c}_μ must

be higher K -trivial. If A is higher K -trivial then the main lemma (Proposition 4.3) shows that A obeys \mathbf{c}_μ for any left-c.e. discrete measure μ .

A \mathbf{c}_μ -bounded test is a higher weak 2 test $\langle U_n \rangle$ such that $\lambda(U_n) \leq^{\times} \mathbf{c}_\mu(n)$; if $\langle U_n \rangle$ is such a test then we may assume that $\lambda(U_{n,s}) \leq^{\times} \mathbf{c}_\mu(n, s)$ (where of course the multiplicative constant is the same for all s). The usual argument shows that if X is a higher ML-random sequence which is captured by some \mathbf{c}_μ -bounded test and A obeys \mathbf{c}_μ then $A \leq_{\omega_1^{\text{ck}} \Gamma} X$. So Theorem 8.2 follows from:

Proposition 8.4. *A sequence is higher Oberwolfach random if and only if it is not captured by any \mathbf{c}_μ -bounded test.*

Proof. In one direction, let $(\langle G_\sigma \rangle, \alpha)$ be a higher Oberwolfach test. For all $n < \omega$ and $s < \omega_1^{\text{ck}}$, let:

- $k_{n,s} = \# \{ \alpha_t \upharpoonright_n : t \leq s \}$ and
- $m_{n,s}$ be the integer m such that $m2^{-n} \leq \alpha_s < (m+1)2^{-n}$.

We define a higher left-c.e. discrete measure ν with the aim that $\mathbf{c}_\nu(n, s) = 2^{-n}k_{n,s} + (\alpha_s - 2^{-n}m_{n,s})$. We would then let $U_n = \bigcup_{s < \omega_1^{\text{ck}}} G_{\alpha_s \upharpoonright_n}$; $\lambda(U_n) = 2^{-n}k_{n, \omega_1^{\text{ck}}} \leq \mathbf{c}_\nu(n)$. The measure ν is not difficult to define. We may assume that for limit s , $\alpha_s = \lim_{t \rightarrow s} \alpha_t$ and so we can let $\nu_s = \sup_{t < s} \nu_t$. Let $\sigma = \alpha_s \wedge \alpha_{s+1}$. We may assume that $\alpha_{s+1} = \sigma 10^\omega$. We then let $\nu_{s+1}(n) = \nu_s + 2^{-n}$ if $n > |\sigma| + 1$ and $\alpha_s(n) = 0$; otherwise we let $\nu_{s+1}(n) = \nu_s(n)$.

In the other direction let $\langle U_n \rangle$ be a \mathbf{c}_μ -bounded test; say $\lambda(U_n) \leq d \cdot \mathbf{c}_\mu(n)$. Let $\mu = d \cdot \mathbf{c}_\mu$ (so $\lambda(U_n) \leq \mathbf{c}_\mu(n)$). By taking a tail of the measure μ (and of the test) and renumbering, we may assume that $\mu(\omega) < 1$. We let $\alpha_s = \mathbf{c}_\mu(0, s) = \mu_s(\omega)$. We define indices $k_s(n)$ for $n < \omega$ and $s < \omega_1^{\text{ck}}$; we let $G_{\alpha_s \upharpoonright_n, s} = U_{k_s(n), s}$. To keep the sets G_σ nested we ensure that $k_s(n)$ is increasing in n . We redefine $k_s(n)$ if $\alpha_s \upharpoonright_n \neq \alpha_{s-1} \upharpoonright_n$. To redefine it we pick a new value k such that $\mathbf{c}_\mu(k, s) \leq 2^{-n}$. Let $t < \omega_1^{\text{ck}}$ and let $\sigma = \alpha_t \upharpoonright_n$; let s be the least stage such that $\sigma < \alpha_s$; let $k = k_s(n) = k_t(n)$. We claim that $\lambda(G_{\sigma, t}) \leq 2^{-(n-1)}$. For $G_{\sigma, t} = U_{k, t}$ and $\lambda(U_{k, t}) \leq \mathbf{c}_\mu(k, s)$; if this is greater than $2 \cdot 2^{-n}$ then as $\mathbf{c}_\mu(k, s) \leq 2^{-n}$ we have $\alpha_t - \alpha_s \geq \mathbf{c}_\mu(k, t) - \mathbf{c}_\mu(k, s) > 2^{-n}$; this implies that $\alpha_t \upharpoonright_n \neq \alpha_s \upharpoonright_n$. \square

The proof in [BGK⁺16] constructing a smart K -trivial set works with subadditive, rather than only with additive cost functions. This proof can be adapted to the higher setting using the usual techniques for overcoming topological problems. However to prove Theorem 8.3 we use a streamlined argument by Turetsky.

Proof of Theorem 8.3. Let Γ be a “universal” higher Turing functional; $\Gamma(0^\epsilon 1X) = \Phi_e(X)$. Since higher Oberwolfach randomness is invariant under the shift, it suffices to enumerate a higher K -trivial c.e. set A and a \mathbf{c}_μ -bounded test $\langle U_n \rangle$ which captures every sequence X such that $\Gamma(X) = A$. In this proof let $\mathbf{c} = \mathbf{c}_\mu$.

We may assume that for all n , $\mathbf{c}(n, 0) > 0$. We enumerate A and $\langle U_n \rangle$ as follows. At each stage we have a “follower” $x_{n,s}$; the sequence $\langle x_{n,s} \rangle$ increases with n . We also enumerate a global error set \mathcal{E}_s ; \mathcal{E}_s is the set of oracles X such that $\Gamma_s(X)$ lies to the left of A_s . Let

$$G_{n,s} = \{ X : \Gamma_s(X) \geq A_s \upharpoonright_{x_{n,s}+1} \}.$$

We will have $U_{n,s} \subseteq \mathcal{E}_s \cup G_{n,s}$. We will change $x_{n,s}$ only finitely many times (for each n), and so at limit stages we can take limits of all objects. We ensure that $\lambda(U_{n,s} - \mathcal{E}_s) \leq \mathbf{c}(n, s)$. Let s be a stage and let $n < \omega$. If $\lambda(G_{m,s} - \mathcal{E}_s) \leq \mathbf{c}(m, s)$ for

all $m \leq n$ then we let $U_{n,s+1} = U_{n,s} \cup G_{n,s}$. If n is least such that $\lambda(G_{n,s} - \mathcal{E}_s) > \mathbf{c}(n, s)$ then we enumerate $x_{n,s}$ into A_{s+1} ; we cancel $x_{m,s}$ for all $m > n$; for all $m \geq n$, we choose unused $x_{m,s+1} > m$ for $m \geq n$, and let $U_{m,s+1} = U_{m,s}$. Note that the enumeration of $x_{n,s}$ into A_{s+1} means that $U_{m,s+1} \subseteq \mathcal{E}_{s+1}$ for all $m \geq n$.

The fact that $G_{n+1,s} \subseteq G_{n,s}$ (as $x_{n+1,s} > x_{n,s}$) ensures that $U_{n+1,s} \subseteq U_{n,s}$ for all n (and all s). If $x_{n,s}$ is enumerated into A_{s+1} then $\lambda(\mathcal{E}_{s+1} - \mathcal{E}_s) > \mathbf{c}(n, s) \geq \mathbf{c}(n, 0)$. This shows that $x_{n,s}$ is enumerated into A_{s+1} at only finitely many stages s . In turn this shows that $x_{n,s+1} \neq x_{n,s}$ for only finitely many stages s .

The enumeration $\langle A_s \rangle$ witnesses that A obeys \mathbf{c} , and so is higher K -trivial. To see this, suppose that $x = x_{n,s}$ is enumerated into A_{s+1} . Then $\mathbf{c}(x, s) \leq \mathbf{c}(n, s)$; this shows that the total cost paid along this enumeration is bounded by $\lambda(\mathcal{E})$.

Finally we need to show that $\lambda(U_n) \leq^{\times} \mathbf{c}(n)$. We enumerate a left-c.e. measure ν , with the aim of having $\lambda(\mathcal{E}_s \cap U_{n,s}) \leq \mathbf{c}_\nu(n, s)$ for all n and s . We would then have $\lambda(U_n) \leq \mathbf{c}_\nu(n) + \mathbf{c}(n) \leq^{\times} \mathbf{c}(n)$ as required. At stage s we need to have

$$\mathbf{c}_\nu(n, s+1) - \mathbf{c}_\nu(n, s) \geq \min\{\mathbf{c}(n, s), \lambda(\mathcal{E}_{s+1} - \mathcal{E}_s)\};$$

this suffices since $U_{n,s+1} \cap (\mathcal{E}_{s+1} - \mathcal{E}_s) \subseteq U_{n,s} - \mathcal{E}_s$. Since $\mathbf{c}(n, s) \rightarrow 0$ as $n \rightarrow \omega$ we can distribute a total of $\lambda(\mathcal{E}_{s+1} - \mathcal{E}_s)$ among the natural numbers (so that $\nu_{s+1}(\omega) \leq \nu_s(\omega) + \lambda(\mathcal{E}_{s+1} - \mathcal{E}_s)$) to achieve the desired increase in $\mathbf{c}_\nu(n, s+1)$. Of course ν is indeed a discrete measure since $\nu(\omega) = \lambda(\mathcal{E})$. \square

REFERENCES

- [AK00] Chris J. Ash and Julia F. Knight. *Computable structures and the hyperarithmetical hierarchy*, volume 144 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 2000.
- [BGHM] Laurent Bienvenu, Noam Greenberg, Denis Hirschfeldt, and Benoit Monin. Bad oracles. In preparation.
- [BGK⁺16] Laurent Bienvenu, Noam Greenberg, Antonín Kučera, André Nies, and Dan Turetsky. Coherent randomness tests and computing the K -trivial sets. *J. Eur. Math. Soc. (JEMS)*, 18(4):773–812, 2016.
- [BMN16] Vasco Brattka, Joseph S. Miller, and André Nies. Randomness and differentiability. *Trans. Amer. Math. Soc.*, 368(1):581–605, 2016.
- [CNY08] Chi Tat Chong, André Nies, and Liang Yu. Lowness of higher randomness notions. *Israel J. Math.*, 166(1):39–60, 2008.
- [CY15] C. T. Chong and Liang Yu. Randomness in the higher setting. *J. Symb. Log.*, 80(4):1131–1148, 2015.
- [DH10] Rodney Downey and Denis Hirschfeldt. *Algorithmic randomness and complexity*. Theory and Applications of Computability. Springer, 2010.
- [DM15] Adam R. Day and Joseph S. Miller. Density, forcing, and the covering problem. *Math. Res. Lett.*, 22(3):719–727, 2015.
- [FHM⁺15] Santiago Figueira, Denis R. Hirschfeldt, Joseph S. Miller, Keng Meng Ng, and André Nies. Counting the changes of random Δ_2^0 sets. *J. Logic Comput.*, 25(4):1073–1089, 2015.
- [FN11] Johanna Franklin and Keng Meng Ng. Difference randomness. *Proceedings of the American Mathematical Society*, 139(1):345–360, 2011.
- [GM] Noam Greenberg and Benoit Monin. Higher randomness and genericity. Submitted.
- [GMS13] Noam Greenberg, Antonio Montalbán, and Theodore A. Slaman. Relative to any non-hyperarithmetical set. *J. Math. Log.*, 13(1):1250007, 26, 2013.
- [HN07] Greg Hjorth and André Nies. Randomness via effective descriptive set theory. *Journal of the London Mathematical Society*, 75(2):495–508, 2007.
- [HNS07] Denis Hirschfeldt, André Nies, and Frank Stephan. Using random sets as oracles. *Journal of the London Mathematical Society*, 75(3):610–622, 2007.
- [Jec08] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer, 2008.

- [Kuč85] Antonín Kučera. Measure, Π_1^0 -classes and complete extensions of PA. In *Recursion theory week*, pages 245–259. Springer, 1985.
- [Kun11] Kenneth Kunen. *Set theory*. Studies in Logic: Mathematical Logic and Foundations. College Publications, 2011.
- [LMN07] Andrew Lewis, Antonio Montalbán, and André Nies. A weakly 2-random set that is not generalized low. In *Proceedings of the CiE 2007*, volume 4497, pages 474–477. Springer, 2007.
- [ML66] Per Martin-Löf. The definition of random sequences. *Information and Control*, 9:602–619, 1966.
- [Mon14] Benoit Monin. Higher randomness and forcing with closed sets. In *LIPICs-Leibniz International Proceedings in Informatics*, volume 25. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2014.
- [MY08] Joseph Miller and Liang Yu. On initial segment complexity and degrees of randomness. *Transactions of the American Mathematical Society*, 360(6):3193–3210, 2008.
- [Nie05] André Nies. Lowness properties and randomness. *Advances in Mathematics*, 197(1):274–305, 2005.
- [Nie09] André Nies. *Computability and randomness*. Oxford Logic Guides. Oxford University Press, 2009.
- [Sac90] Gerald Sacks. *Higher recursion theory*. Perspectives in Mathematical Logic. Springer, 1990.
- [Ste75] Jacques Stern. Some measure theoretic results in effective descriptive set theory. *Israel J. Math.*, 20(2):97–110, 1975.
- [Ste78] John R Steel. Forcing with tagged trees. *Annals of Mathematical Logic*, 15:55–74, 1978.
- [vL87] Michiel van Lambalgen. *Random sequences*. PhD dissertation, University of Amsterdam, Amsterdam, 1987.
- [Yu06] Liang Yu. Lowness for genericity. *Archive for Mathematical Logic*, 45(2):233–238, 2006.
- [Yu11] Liang Yu. A new proof of Friedman’s conjecture. *Bull. Symbolic Logic*, 17(3):455–461, 2011.

LIAFA, UNIVERSIT PARIS DIDEROT, PARIS, FRANCE
E-mail address: laurent.biennu@computability.fr
URL: <http://www.liafa.jussieu.fr/~lbiennu/>

SCHOOL OF MATHEMATICS, STATISTICS AND OPERATIONS RESEARCH, VICTORIA UNIVERSITY OF WELLINGTON, WELLINGTON, NEW ZEALAND
E-mail address: greenberg@msor.vuw.ac.nz
URL: <http://homepages.mcs.vuw.ac.nz/~greenberg/>

SCHOOL OF MATHEMATICS, STATISTICS AND OPERATIONS RESEARCH, VICTORIA UNIVERSITY OF WELLINGTON, WELLINGTON, NEW ZEALAND
E-mail address: benoit.monin@computability.fr
URL: <http://www.liafa.univ-paris-diderot.fr/~benoitm/>